



Occurrence of *Bacillus thuringiensis* from different plant areas on South Kalimantan, Indonesia

Akhmad Rizali*

Department of Agroecotechnology, Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru, Indonesia

Article published on December 27, 2017

Key words: *B. thuringiensis*, Different plant areas, South Kalimantan, Indonesia

Abstract

B. thuringiensis is a gram-positive soil bacterium, and produce a crystalline inclusion body during sporulation. Therefore, many biological control of insects have been investigated. Currently, researches on the use pathogenic microorganisms to control insect pests are increasing. Microbial pest control is practiced in different parts of the world though utilization of pathogen like fungi, bacteria, viruses and nematodes.. It was a study of flacherie of the silkworm, *bombxmori* in this report on the discovery of *sotto bacillus*, which causes the disease to silkworm larvae. Five 1-g soil samples were separately suspended to 9 ml of distilled water. After allowing the suspension to stand for 5 minute, 3-4 ml of the suspension were taken. One half of the suspension was transferred to a test tube and heated in a water bath of 80°C for 15 minutes, so that all microorganisms were killed except *Bacillus* and other spore forming bacteria, then allowed to cool at room temperature. Ten-fold serial dilutions of the heated suspension in sterile distilled water were placed on nutrient agar (NA-pH 7.5). After two days of incubation at 28°C, *Bacillus* colonies were recorded. After 2 to 3 days incubation, crystalliferous spore forming bacteria were determined in phase contrast microscope. Isolation from six soil samples yielded about 50 isolates; only one was identified as *B. thuringiensis*. Observations on *B. thuringiensis* isolated from citrus areas on *C. binotalis* showed that infected larvae turned yellowish at the middle and hind part of the abdomen and dead larvae become shrunken and later turned black with putrid odor.

*Corresponding Author: Akhmad Rizali ✉ arizali25@yahoo.com

Introduction

Cabbage moth, *C. binotalis* Zell. (Lepidoptera: Pyraustidae) is considered the most important limiting factor for a successful production of cruciferous vegetable not only in the Indonesia but in other country in the world. The larva feeds on foliage from seedling to harvest causing 100% yield loss if not control (Rejesus and Sayaboc, 1990).

Numerous chemical insecticides have been used in order to control pests, which damage for agriculture. While chemical insecticides have knock down effect to the insect pests, they are too expensive in the developing countries and harmful to both human and the environment. In addition, target insect pests rapidly develop biological resistance especially at higher rates of application. The chemical insecticides are still contributing to human life enormously, but they have been distributed in ecological system of organisms including human beings because of their low specific toxicity to any organism and their low specific toxicity to any organism and their slight decomposition in nature (Shorey and Hall, 1962). Therefore, many biological control of insects have been investigated. Currently, researches on the use pathogenic microorganisms to control insect pests are increasing. Microbial pest control is practiced in differen parts of the world though utilization of pathogen like fungi, bacteria, viruses and nematodes. Bacterial research causing disease in insects began in the late nineteenth century. It was a study of flacherie of the silkworm, *bombxmori* (Burges and Hussey, 1971; Burges, 1981). Ishiwata (1901) in this report on the discovery of *sotto bacillus*, reffered briefly to occurrence of sotto bacillus-like organism, which causes the disease to silkworm larvae.

Berliner (1911) proposed the name of *B. thuringiensis* for a species of bacillus which was isolated from the diseased larvae of the Mediterranean flour moth *Anagasta (Ephestia) kuhniella* Zell. Later, Berliner (1915) noted infection of the larvae after the ingestion of the bacillus or its spore, described and named it *B. thuringiensis*. Mattes (1927) isolated the same bacillus from the same insect host, which Berliner had found earlier.

B. thuringiensis is a gram-positive soil bacterium, and produce a crystalline inclusion body during sporulation (Bulla *et al.*, 1980). This parasporal body is composed of proteins termed “delta-endotoxin”, and specifically toxic to insects. In addition, *B. thuringiensis* produce another toxins namely: alpha-toxin, beta-exotoxin, and gamma-exotoxin. All of the toxic substance may not present in the bacterium (Heimpel, 1967). In another hand, Krieg (1961) has defined various toxic substance produced *B. thuringiensis* as follow: (a) thermolabileendotoxic; (b) thermostable exotoxin; (c) bacillogenic antibiotic; (d) lecithinase; (e) proteinase.

Most strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to lepidopteran insects such as moth (Dulmage *et al.*, 1970). Recently, however several researches have shown that *B. thuringiensis* is also widely distributed in natural soils of various area. Delucca *et al.*, (1982) reported that *B. thuringiensis* made up less than 0.5% of more than 46,000 bacterial isolates recovered from various soils in the United States. *Crocidolomia binotalis* larvae feeds on foliage from seedling to harvest causing 100% yield loss if not control.so that very important to select *B. thuringiensis* to control the larvae.

The objective of the studies to survey, collect and determine the distribution of *B. thuringiensis* in selected diverse crop-growing area.

Materials and methods

Isolation of *B. thuringiensis*

Soil samples were collected in areas planted to vegetables, rice, citrus, peanut and corn in South Kalimantan (Indonesia) following multistage random sampling. Soil samples were collected at random in a 1- hectare area for a total of 5 kg. The soil sample were taken from the top 1 cm of the soil layer. The 5-kg soil samples were mixed thoroughly and composite sample of 1 kg was taken from which isolation were made for as long as one month. The samples were labeled denoting date, place of collection and crops planted.

Five 1-g soil samples were separately suspended to 9 ml of distilled water. After allowing the suspension to stand for 5 minute, 3-4 ml of the suspension were taken. One half of the suspension was transferred to a test tube and heated in a water bath of 80°C for 15 minutes, so that all microorganisms were killed except bacillus and other spore forming bacteria, then allowed to cool at room temperature. Ten-fold serial dilutions of the heated suspension in sterile distilled water were placed on nutrient agar (NA-pH 7.5). After two days of incubation at 28°C, colonies were

recorded. After 2 to 3 days incubation, crystalliferous spore forming bacteria were determined in phase contrast microscop.

Results and discussion

Isolation and distribution of B. thuringiensis in different plants growing areas

Fifty *Bacillus sp.* were isolated from 6 soil samples collected from diverse crop growing areas in the South Kalimantan, Indonesia (Table 1).

Table 1. South Kalimantan Isolates with different plants areas source.

Isolates	Crop planted	Soil Source	
		Great Soil	pH
AA	Rice	Typicalciborolls	6.5
AA1.1			
AA1.2			
AA2.1 (1)			
AA2.2			
AA2.3			
AA4.1			
AA2.5			
AA2.5 (1)			
AA2.5 (2)			
BB	Vegetable	Typictropudults	4.8
BB 1.1			
BB 1.2			
BB 2.3			
BB 1.4			
BB 3.1			
CC	Corn	„	5.2
CC 1.2			
CC 2.1			
CC 2.3			
CC 2.4			
CC 3.2			
CC 5.1			
CC 5.1 (1)			
DD	Citrus	„	5.7
DD 1.2 (1)			
DD 1.3			
EE	Peanut	„	4.7
EE 1			
EE 2			
EE 1.3			
EE 1.4			
EE 3.1			
EE 4.3			
F	Cabbage	„	5.1
FF 1.2			
FF 2.1			
FF 2.2 (1)			
FF 2.3			
FF 2.4			
FF 4.1			
FF 3.1			
FF 3.2			

Information : From AA 1.1 to FF 5.1 code of soil sample was collected from different growing areas.

The different isolates were obtained from the same area planted with diverse crop. Two soil samples great were found in this area namely: typiccalciborrols with pH 6.5 was planted to rice and typictropudults pH ranging from 4.8 – 5.7 planted to vegetable, corn, citrus, peanut and cabbage. Out of the 50 *Bacillus sp.* isolates, only one (2%) was identified as *B. thuringiensis* based on phase contrast microscope examination for the presence of parasporal inclusion bodies. Only the soil sample from citrus yielded *B. thuringiensis*. The possible reasons for the low incidence of *B. thuringiensis* isolated from the samples taken in the areas surveyed are the small number of samples size from which the isolation were made, the area where sampling was done and also the difference in the physic-chemical characteristic of the soil where samples were taken. The low incidence of *B. thuringiensis* was also reported in Japan by Ohba and Aizawa (1966) in soil sample from non-agricultural areas. Out of 6910 isolates only 189 (2.7%) isolates were identified as *B. thuringiensis*.



Fig. 1. Dead larvae of *C. binotalis* showing the progress of symptoms of *B. thuringiensis* isolated from citrus areas.

Symptomatology

Observations on of *B. thuringiensis* isolated from citrus areas on *C. binotalis* showed that infected larvae turned yellowish at the middle and hind part of the abdomen (Fig.1). The integument also turned brown to black as the infection progressed and the body became sticky because of oral and anal discharges. Dead larvae become shrunken and later turned black with putrid odor.

Conclusion

Soil samples were collected in areas planted to vegetables, rice, citrus, peanut and corn in South Kalimantan (Indonesia) following multistage random sampling. Soil samples were collected at random in a 1- hectare area for a total of 5 kg. The soil sample were taken from the top 1 cm of the soil layer.

Six soil samples were collected from different crop growing areas in South Kalimantan (Indonesia). Isolation from six soil samples yielded about 50 isolates, only one was identified as *B. thuringiensis*. The toxicity of *B. thuringiensis* isolated from citrus areas on *C. binotalis* showed that infected larvae turned yellowish at the middle and hind part of the abdomen and Dead larvae become shrunken and later turned black with putrid odor. This result suggest that *B. thuringiensis* is rare and not widespread in the places where sampling was done and isolates of *B. thuringiensis* will screen to the cabbage worm (*Crociodomia binotalis*).

References

- Abdel-Hameed A, Carlberg G, El-Tayeb, OM** 1990. Studies on *Bacillus thuringiensis* H-14 strains isolated in Egypt. Screening for active strains. World Journal of Microbiology and Biotechnology. 6: 299-304.
- Asano S, Bando H, Iizuka T.** 1993. Amplification and identification of *cryII* genes from *Bacillus thuringiensis* by PCR procedures. Journal Sericulture. Science. Japan. **62**, 223-227.
- Baba F, Asano S, Iizuka T.** 1990. Purification of crystals from *Bacillus thuringiensis* by using Percoll. Journal Sci. Jpn **59**, 487-489.
- Balarman K, Hoti SL, Manonmani LM.** 1981. An Idigenous virulent strain of *Bacillus thuringiensis*, highly pathogenic and specific to mosquitoes. Current Science **50**, 199-200.
- Berliner E.** 1915. Ober dieschalaffsucht der mehlmottenraupe (*Ephestia kuhniella* Zeller) und ihreerreger *Bacillus thuringiensis* n. sp. Zangue. Entomol. **2**, 29-56.

- Bulla LA, faust RM, Andrews R, Goodman N.** 1985. Insecticidal bacilli, pp. 185-209. In: the Molecular Biology of the Bacilli, Vol. II. D. A. Dubnau (Ed.). Academic Press, New York.
- Burges HD.** 1981. Microbial control of pest and plant disease. 1970-1980. Academic Press. 949 p.
- Burges HD, Hussey NW.** 1971. Microbial control of insects and mites. Academic Press. 876 p.
- Bulla LA, Jr, Kramer KJ, Davidson LI.** (1977). Characterization of the entomocidal parasporal crystal of *Bacillus thuringiensis*. Journal Bacteriology. **130**, 375-383.
- Delucca AJ, Simonson JG, Larson AD.** 1981. *Bacillus thuringiensis* distribution in soils of the United States. Canadian Journal Microbiology. **27**, 865-870.
- Dulmage HT.** 1992. Insecticidal activity of *Bacillus thuringiensis* and their potential for pest control in Microbial control for pests and plant diseases and plant diseases 1970-1980 (Ed. H.D. Burges). Academic Press. N.Y. P.
- Golberg LJ, Margalit J.** 1977. A bacterial spore demonstrating rapid larvicidal activity against *Anopheles sergentii*, *Uranotaenia unguiculata*, *Culex univittatus*, *Aedes aegypti* and *Culex pipiens*, Mosq. News **37**, 355-358.
- Heimpel AM.** 1967. A critical review of *Bacillus thuringiensis* Berl. and other crystalliferous bacteria. Annual Review Entomology **12**, 287-322.
- Held GA, Kawanishi CY, Huang YS.** 1990. Characterization of the parasporal inclusion of *Bacillus thuringiensis* subsp. *kyushuensis*. Journal Bacteriology 481-483.
- Iizuka T, Ishino M, Nakajima T.** 1982. Comparative morphology of Parasporal crystal and characterization of plasmid DNA from various subspecies of entomopathogenic bacteria, *Bacillus thuringiensis*. Journal Faculty Agriculture. Hokkaido University. **13**, 423-431.
- Iizuka T, Yamamoto T.** 1984. Serological properties of the mosquitocidal protein of *Bacillus thuringiensis* and the morphology of its parasporal crystal. Journal Faculty. Hokkaido University **62**, 98-114.
- Iizuka T, Sasaki J, Asano S, Bando H.** 1995. Comparative studies on isolation and identification of *Bacillus thuringiensis*. Biotechnology and Environment. Benefits, **1**, 143-153.
- Ishii T, Ohba M.** 1997. Investigation of mosquito-specific larvicidal activity of a soil isolate of *Bacillus thuringiensis* var. *canadensis*. Current Microbiology. **35**, 40-43.
- Ishiwata S.** 1901. On a kind of severe flacherie (sotto disease). Dainihon Sanshikaiho **114**, 1-5.
- Kalman S, Kiehne KK, Libs JL, Yamamoto T.** 1993. Cloning of novel cryIC-type gene from a strain *Bacillus thuringiensis* subs. *Galleriae*. Applied. Environment. Microbiology **59**, 1131-1137.
- Kawalek MD, Benjamin S, Lee HL, Gill SS.** 1995. Isolation and identification of novel toxin from a new mosquitocidal isolate from Malaysia, *Bacillus thuringiensis* subsp. *Jegathesan*. Applied. Environment. Microbiology. 2965-2969.
- Kim KH, Ohba Aizawa K.** 1984. Purification of the toxic protein from *Bacillus thuringiensis* serotype 10 isolate demonstrating a preferential larvicidal activity to mosquito. Journal Invertebrate. Pathology. **44**, 214-219.
- Krieg A.** 1961. *Bacillus thuringiensis* Berliner. In disease caused by certain sporeforming bacteria. Heimpel and Angus (Eds.). 21-67.
- Mattes O.** 1927. Parasitarekran kheiten der mehnotten larvaen und versucheoberih reverwnd barkeitals biologische becamp fungus mittel. (Zuheich lire beitragszurzytologie de bacgerien). Gesell f. beförd, gedam, Naturw. Sitzber (Marbnog) **62**, 381-417.

- Ohba M, Aizawa K.** 1986. Insect toxicity of *Bacillus thuringiensis* isolated from soils of Japan. *Journal Invertebrate Pathology*. **47**, 12-20.
- Padua LE, Ohba M, Aizawa K.** 1984. Isolation of a *Bacillus thuringiensis* strain (serotype 8a:8b) highly and selectively toxic against mosquito larvae. *Journal Invertebrate Pathology*. **44**, 12-17.
- Rejesus BM, Sayaboc A.** 1990. Management of DBM with *Apanteles plutellae* pros fects in the Philippines. Paper presented in the second International Workshop on Diamondback Moth Management. Dec. 10-15, 1990. Tainna, Taiwan. 17 p.
- Rizali A, Shin-ichiro Asano, Ken Sahara, Hisanori Bando, Bibiana W, Lay, Sugyo Hastowo, Toshihiko Iizuka.** 1998. Novel *Bacillus thuringiensis* serovar aizawai strains isolated from mulberry leaves in Indonesia. *Applied Entomology Zoology*. **33(1)**, 111-114.
- Shorey HH, Hall IM.** 1962. Effect of chemical and microbial insecticides on several insect pests of lettuce in southern California. *Journal Economic Entomology*. **561**, 69-174.
- Yu YM, Ohba M, Gill SS.** 1991. Characterization of mosquitocidal activity of *Bacillus thuringiensis* subsp. *fukuokaensis* crystal proteins. *Applied. Enviroment. Microbiololgy*. 1075-1081.
- Zhang Y, Ku Z, Chan Z, Xu B, Yuan F, Chen G, Zhong T, Ming G.** 1984. A new isolate of *Bacillus thuringiensis* possessing high toxicity against the mosquitoes. *Acta Microbiologica Sinica* **24**, 320-325.