



RESEARCH PAPER

OPEN ACCESS

Identification of most effective sterilization protocol(S) for establishment of *in vitro* culture of endangered medicinal plant *Skimmia laureola*

Samina Siddique¹, Zishan Gul^{*2}, Zafar Jamal¹, Asif ur Rehman²

¹Department of Botany, G. Post Graduate College, Abbottabad, Pakistan

²Hazara Agriculture Research Station, Abbottabad, Pakistan

Key words: *Skimia laureola*, Medicinal plant, Sterilization, Micro propagation

<http://dx.doi.org/10.12692/ijb/12.2.199-209>

Article published on February 26, 2018

Abstract

Skimmia laureola (DC) is an important endangered medicinal plant. The present research was conducted to investigate the effect of sterilizing procedure(s) for establishment of *in vitro* culture of *Skimmia laureola*. The explants were cultured on MS medium containing coconut water as growth promoter instead of any synthetic plant growth hormone. Mercuric chloride (HgCl₂) at 0.5 % concentration for 30 seconds and 1 minute was found effective sterilization procedure when used alone or in combination with fungicide (Ridomil) where 100 % contamination was controlled with 100 % plants survived. Sodium hypochlorite (NaOCl) alone and with fungicide at both concentrations i.e 3% and 5% remained less effective, however, NaOCl + fungicide at 3% concentration for 1 minute showed 20% contamination with 80% survived explants. After 28 days the survived plantlets in HgCl₂ treatment showed average 3.26 cm shoot length with average 5 number of leaves per explant. It has been concluded that HgCl₂ can be used as sterilant for establishment of *Skimmia laureola in vitro* cultures and further work is needed for standardizing HgCl₂ concentrations and time of exposure for sterilization of other endangered medicinal plants.

* Corresponding Author: Zishan Gul ✉ gul.zishan@gmail.com

Introduction

Skimmia laureola (*S. laureola*) (DC.) Decneis an aromatic shrub belongs to family Rutaceae. In Pakistan, *Skimmia laureola* grows at an altitude of 5500- 10000 feet. It is common in the Hazara region, Murree Hills and Kashmir, in Upper Swat and Shangla (Hamayun *et al.*, 2006; Ali and Qaiser, 2007) and Upper and Lower Dir (Barkatullah *et al.*, 2012).

Traditionally, this plant is used as an antitussive, as veterinary anthelmintic, as an insecticide and pesticide. The garlands made from *Skimmia* fresh leaves are considered sacred and smoke of dried leaves is considered useful to ward off evils (Bhattari, 2006; Humayun, 2007). The smoke of the dried leaves is used to treat cold, fever and headache (Qureshi, 2009) and relieves cough (Baart *et al.*, 2004). The essential oils of this plant were used as antioxidant, antimicrobial and antifungal. The presence of high percentages of linalyl acetate and linalool in the oil have been reported for antibacterial and antifungal properties due to their ability to cross cell membranes, penetrating into the interior of the cell and interacting with critical intracellular sites (Trombetta *et al.*, 2005).

Nowadays, Tissue culture technology is used as a mean for germ-plasm conservation of endangered plants, for rapid mass propagation for large-scale revegetation, for genetic manipulation studies and for large scale production of high quality plant based medicines (Debnath, 2006; Yadav *et al.*, 2012).

Microbial contaminations are the major problem to initiate viable *in vitro* cultures. Losses due to contamination in *in vitro* cultures ranged between 3 to 15%. Plant tissues have various microorganisms on their surfaces which can grow in the culture medium, making the culture non sterile. In addition, they compete with the plant tissue for nutrition (CPRI, 1992). A successful tissue culture protocol starts with effective explants sterilization (Dodds and Roberts, 1985). Sterilization is the process of making explants contamination free before *in vitro* establishment of cultures. Various sterilization agents used for this purpose are also toxic to the plant tissues, hence

proper concentration of sterilants, duration of exposure and the sequence of using these sterilants have to be standardized to minimize the injury to the explants (Goswami and Handique, 2013).

Sodium hypochlorite, ethanol, mercuric chloride and fungicides are common sterilants when using field grown tissues. The time of sterilization depends on tissue type (Funguomali *et al.*, 2013, Sharma and Nautiya, 2009). Nurul *et al.* (2012) reported that pre-sterilization using 0.2% Benomyl for 15 minutes reduces the contamination and mortality rate for leaf and nodal explants from *Aquilaria malaccensis*, especially when followed by surface sterilization with 0.1 % mercury chloride for 15 and 30 seconds.

There was no previous reports on *in vitro* sterilization protocols for *Skimmia laureola* micropropagation. Therefore the present study was conducted to test the efficiency of various sterilization procedures against microbial contamination for the primary establishment of *in vitro* culture in order to standardize *in vitro* micropropagation protocol for *Skimmia laureola*.

Materials and methods

Plant material

The present study was conducted in the tissue culture laboratory at Hazara Agriculture Research Station Abbottabad during 2017. The *Skimmia laureola* was collected from naturally grown habitat of Nathia gali hills Abbottabad during the month of March. Plant was identified by using plant identification key 'Flora of Pakistan (Hassan ud din and Ghazanfar, 1980).

Media preparation

Murashige and Skoog (1962) basal medium containing 1 mgL⁻¹Ca-pentothenate, 100mgL⁻¹myoinositol with 30g/L sucrose and 6g/L agar was prepared. The coconut water @ 120ml/L was added to media instead of plant growth hormones. The pH was adjusted to 5.8 before autoclaving at 121 °C and 15 psi pressure for 15 minutes.

Explants surface sterilization

Each sterilizing agent at its respective concentration was tested for 3 different time periods.

There were twenty four experimental treatments (Table 1), each replicated 5 times. The fungicide (Ridomil) was used @ 0.1% (w/v).

After removing the leaves, *Skimmia* explants were washed under running tap water for 30 minutes. Then washed with mild detergent solution for 5 min and rinse thoroughly with distilled water.

The explants were cut into nodal segments. After applying the respective sterilization protocol(s) (Table 1) the nodal segments were rinsed 4 to 5 times with sterilized distilled water in order to remove any traces of sterilants. The test tubes were incubated in growth chamber at 20 °C under 16 h light and 8 h dark photoperiod for about 28 days.

The experimental design used was CRD. Data were analyzed by using computer software Statistix 8.1 and least significance difference test (LSD) at 95% level of significance was used to assess significant differences between various treatments.

Results and discussion

Sterilization is the process of making explants contamination free before establishment of cultures. The use of field grown plants as direct source for *in vitro* cultures presents a major challenge (Webster, *et al.*, 2003). The data of present study revealed that different sterilization protocols (Table 1) showed different effect on the establishment of *in vitro* culture and growth of *Skimmia laureola*.

Table 1. Different sterilization protocols used for *Skimmia* explants sterilization.

Treatments/Sterilization protocols	Sterilizing agents	Concentration%	Exposure time
T1	Mercuric chloride (HgCl ₂)	0.1	30sec
T2			1min
T3			3min
T4	Mercuric chloride (HgCl ₂)	0.5	30sec
T5			1min
T6			3min
T7	Sodium hypochlorite (NaOCl)	3	30sec
T8			1min
T9			3min
T10	Sodium hypochlorite (NaOCl)	5	30sec
T11			1min
T12			3min
T13	Sodium hypochlorite (NaOCl)+fungicide	3	30sec
T14			1min
T15			3min
T16	Sodium hypochlorite (NaOCl)+fungicide	5	30sec
T17			1min
T18			3min
T19	Mercuric chloride (HgCl ₂)+fungicide	0.1	30sec
T20			1min
T21			3min
T22	Mercuric chloride (HgCl ₂)+fungicide	0.5	30sec
T23			1min
T24			3min

Percent contamination

Data regarding the effectiveness of various sterilization procedures (Treatments) on percent contamination of explants after 7 days of culturing

(Table 2) showed that mercuric chloride at both 0.1% and 0.5% was found significantly ($P \leq 0.05$) effective in reducing the contamination of cultured nodes (Table 2).

Table 2. Effect of sterilizing procedures on % contamination, % emergence and % survival rate of *Skimmia laureola* explants after 7 days of culturing.

Treatments	Sterilizing Agent	Concentration %	Time of exposure	Contamination %	Emergence %	Survival Rate %
T1	Mercuric chloride (HgCl ₂)	0.1	30sec	0c	100a	100a
T2			1min	0c	100a	100a
T3			3min	0c	100a	100a
T4	Mercuric chloride (HgCl ₂)	0.5	30sec	0c	100a	100a
T5			1min	0c	100a	100a
T6			3min	0c	100	100a
T7	Sodium hypochlorite (NaOCl)	3	30sec	0c	100a	100a
T8			1min	80ab	20c	20c
T9			3min	0c	100a	100a
T10	Sodium hypochlorite(NaOCl)	5	30sec	40b	60b	60b
T11			1min	0c	100a	100a
T12			3min	80a	20c	20c
T13	Sodium hypochlorite (NaOCl) +fungicide	3	30sec	0c	100a	100a
T14			1min	0c	100a	100a
T15			3min	0c	100a	100a
T16	Sodium hypochlorite (NaOCl) +fungicide	5	30sec	40b	60b	60b
T17			1min	100a	0c	0c
T18			3min	20bc	80ab	80ab
T19	Mercuric chloride (HgCl ₂) +fungicide	0.1	30sec	0c	100a	100a
T20			1min	0c	100a	100a
T21			3min	0c	80ab	80ab
T22	Mercuric chloride (HgCl ₂) +fungicide	0.5	30sec	0c	100a	100a
T23			1min	0c	100a	100a
T24			3min	0c	100a	100a

Means in a column expressed with different alphabets are significantly different ($P \leq 0.05$).

Such response might be due to bleaching action of two chloride atoms and also ions that combines strongly with proteins and causing the death of organisms. There are many reports of surface sterilization in plant tissue culture using HgCl₂ (Naika and Krishna, 2008; Preethi *et al.*, 2011; Sen *et al.*, 2013).

The efficacy of HgCl₂ solution as surface sterilant in *in vitro* cultures of *A. paniculata* had also been reported by Martin, 2004; Purkayastha *et al.*, 2008; Katakya and Handique, 2010) who reported that 0.1% solution of HgCl₂ was sufficient in establishing the explants in culture medium. Whereas sodium hypochlorite at 3% concentration for 30 seconds (T7) and at 5 % concentration for 1 minute (T11) showed 0% contamination. After 14 days of culturing (Table 3) the sterilization procedures (T1, T2, T3, T4, T5, T6) also

showed 0 % contamination while in the treatments 3% and 5% NaOCl (T7) and (T10) each for 30 seconds the contamination increased to 100% (Fig. 1).

In our study the sterilizing agent NaOCl at both concentration i.e 3% and 5% showed contamination for all the tested time periods. Daud *et al.* (2012) reported that NaOCl was ineffective on *A. malaccensis* for leaves and nodal segments explants and concluded that this effect is because the plant materials came directly from natural surroundings and even if the surface is effectively sterilized, the contaminants could come from the inner tissues after dissection of plant materials into small explants after the surface sterilization. After 28 days of inoculation (Table 4) similar trend was observed.

Table 3. Effect of sterilizing procedures on % contamination, % emergence, % survival rate, % death, % discoloration, shoot length and no of leaves of *Skimmia laureola* explants after 14 days of culturing.

Treatments	Sterilizing Agent	Concentration %	Time of exposure	Contamination %	Emergence %	Survival Rate %	Death %	Discoloration %	Shoot length (cm)	No of leaves
T1	Mercuric Chloride(HgCl ₂)	0.1	30sec	0e	80cd	80cd	20ab	20a	0.12cde	0.4fg
T2			1min	0e	60abc	60abc	40bcd	40bc	0.6bcde	2.2abcd
T3			3min	0e	20ab	20ab	80cd	80cd	0.32cde	1.8abcdef
T4	Mercuric Chloride(HgCl ₂)	0.5	30sec	0e	100a	100a	0d	0d	2.24a	3ab
T5			1min	0e	100a	100a	0d	0d	0.7bcde	2abcde
T6			3min	0e	40bcd	40bcd	80ab	80a	0.4cde	0.8defg
T7	Sodium hypochlorite(NaOCl)	3	30sec	100a	0d	0d	100a	0d	0e	0g
T8			1min	80ab	20cd	20cd	80ab	0d	0.32cde	0.8defg
T9			3min	20de	80ab	80ab	20cd	0d	0.48cde	2.4abc
T10	Sodium hypochlorite(NaOCl)	5	30sec	100a	0d	0d	100a	0d	0e	0g
T11			1min	60bc	20cd	40bcd	60abc	0d	0.1de	1cdefg
T12			3min	80ab	80ab	20cd	80ab	0d	0.4cde	0.6efg
T13	Sodium hypochlorite(NaOCl)	3	30sec	100a	0d	0d	100a	0d	0e	0g
T14	+fungicide		1min	20de	80ab	80ab	20cd	0d	0.8bcd	2.4abc
T15			3min	80ab	20cd	20cd	80ab	0d	0.08e	0.4fg
T16	Sodium hypochlorite(NaOCl)	5	30sec	100a	0d	0d	100a	0d	0e	0g
T17	+fungicide		1min	100a	0d	0d	100a	0d	0e	0g
T18			3min	100a	0d	0d	100a	0d	0e	0g
T19	Mercuric Chloride(HgCl ₂)+ Fungicide	0.1	30sec	0e	100a	100a	0d	0d	0.92bcde	3.2a
T20			1min	0e	60abc	60abc	40bcd	40bc	0.84bcde	1.6bcdef
T21			3min	20de	40bcd	40bcd	60abc	40bc	0.3cde	1cdefg
T22	Mercuric Chloride(HgCl ₂)+ fungicide	0.5	30sec	0e	100a	100a	0d	0d	1.3b	3ab
T23			1min	0e	80ab	80ab	40bc	40bc	0.7bcde	1.4cdefg
T24			3min	0e	40bcd	40bcd	80ab	80a	0.6bcde	1.2cdefg

Means in a column expressed with different alphabets are significantly different ($P \leq 0.05$).

Shoot emergence

After seven days of inoculation (Table 2) maximum shoot emergence(100%) was observed in treatment T1, T2, T3, T4, T5 and T6 containing 0.1 % and 0.5 % HgCl₂ alone and 0.1 % and 0.5 % HgCl₂ with fungicide (T19, T20, T21, T22, T23, T24) for three respective time period i.e 30 seconds, 1 minute and 3 minutes (Fig. 2). Similarly Treatment T11(5% NaOCl for 1 minute), treatment T7 and T9 having 3% NaOCl for 30 seconds and 3 minute respectively and treatment T13, T14, T15 (3% NaOCl + fungicide) for three different time periods also showed 100% shoot emergence followed by Treatment T18 (5% NaOCl + fungicide) and T21 (0.1 % HgCl₂+fungicide) for 3 minutes showed 80% emergence.

The minimum shoot emergence i.e 20% was observed in treatment T8 and T12 (NaOCl 3% and 5 % for 1 and 3 minutes, respectively) (Table 2).

Results after 14 days (Table 3) showed that the sterilization procedure 0.5 % HgCl₂ (T4,T5) for 30 seconds and 1 minute and HgCl₂ + fungicide (T19,

T22) at both concentration (0.1 % and 0.5 %) for 30 seconds showed 100% shoot emergence while treatment T9 (3% NaOCl for 3 minutes), treatment T1 (0.1 % HgCl₂), treatment T14 (3% NaOCl + fungicide) and treatment T23(0.5 % HgCl₂+fungicide 0.5 %) for 1 minute showed 80 % shoot emergence. However, NaOCl treatments T7, T10, T13, T16, T17 and T18 showed 0% shoot emergence after 28 days of inoculation (Table 4).

Kiani *et al.*, (2017) obtained maximum percentage of germination (81.33%) of *Melissa officinalis* L. after treating with 2.5% Sodium Hypochlorite for 3 min and (32.66%) at 0.2% Mercuric Chloride for 6 minutes.

Survival rate

Effect of sterilizing agents on explants survival after seven days of inoculation (Table 2) showed 100% survival rate in HgCl₂ treatments. Results of present study is in agreement with other reports (Alam, 2016) who reported that HgCl₂ at 0.1% for 2 min time caused 55.55% of cultures of *Cucumis sativus*

remained survived. 0.1% HgCl₂+fungicide (Ridomil) showed 0% contamination after 7 days of culturing but after 28 days 20% contamination and 80% survival rate was observed at minimum time period

but on increasing time of exposure i.e 3 min, contamination rate remain 20% but survival rate was reduced to 40% (Table 4).

Table 4. Effect of sterilizing procedures on % contamination, % emergence, % survival rate, % death, % discoloration, shoot length and no of leaves of *Skimmia laureola* explants after 28 days of culturing.

Treatments	Sterilizing Agent	Concentration % (w/v)	Time of exposure	Contamination %	Emergence %	Survival Rate %	Death %	Discoloration %	Shoot length (cm)	No of leaves
T1	Mercuric chlorid (HgCl ₂)	0.1	30sec	20c	60cd	60cd	40a	20a	0.16fg	0.60ef
T2			1min	40bc	60abc	60abc	40bcd	od	0.88bcdefg	3.4abcd
T3			3min	20c	20abc	20abc	100bcd	80cd	0.52defg	2.4bcdef
T4	Mercuric chlorid (HgCl ₂)	0.5	30sec	0c	100a	100a	od	od	3.26a	4.6ab
T5			1min	0c	100a	100a	od	od	1.74ab	4.6ab
T6			3min	0c	40bcd	40bcd	80ab	80a	0.58defg	1.2def
T7	Sodium hypochlorite (NaOCl)	3	30sec	100a	od	od	100a	od	0g	0f
T8			1min	80ab	20cd	20cd	80ab	od	0.4efg	1.2def
T9			3min	40bc	60abc	60abc	40bcd	od	0.68cdefg	2.8abcde
T10	Sodium hypochlorite (NaOCl)	5	30sec	100a	od	od	100a	od	0g	0f
T11			1min	60abc	40bcd	40bcd	60abc	od	0.66cdefg	3.6abcd
T12			3min	80ab	20cd	20cd	80ab	od	0.5efg	1.2def
T13	hypochlorite (NaOCl) +fungicide	3	30sec	100a	od	od	100a	od	0g	0f
T14			1min	20c	80ab	80ab	20cd	od	1.2bcde	3.8abc
T15			3min	80ab	20cd	20cd	80ab	od	0.30fg	0.8ef
T16	hypochlorite (NaOCl) +fungicide	5	30sec	100a	od	od	100a	od	0g	0f
T17			1min	100a	od	od	100a	od	0g	0f
T18			3min	100a	od	od	100a	od	0g	0f
T19	Mercuric chlorid (HgCl ₂) +fungicide	0.1	30sec	20c	80ab	80ab	20cd	od	0.86bcdefg	3.4abcd
T20			1min	20c	40bc	40bc	60abc	40bc	0.76cdefg	2.6abcde
T21			3min	20c	40bcd	40bcd	60abc	40bc	0.60defg	1.60cdef
T22	Mercuric chlorid (HgCl ₂) +fungicide	0.5	30sec	0c	100a	100a	od	od	1.50abc	5.00a
T23			1min	0c	60abc	60abc	40bcd	40bc	1.40abcd	3.6abcd
T24			3min	0c	20cd	20cd	80ab	80a	0.92bcdef	2cdef

Means in a column expressed with different alphabets are significantly different ($P \leq 0.05$).

Results similar to present study was also reported by Sen *et al.* (2013) who found that at high concentration of HgCl₂ with fungicide control contamination but cause corrugation to explants.

Similarly Treatment T11, T7 and T9 and treatment T13, T14, T15 having 0.1% NaOCl and 3% NaOCl+ fungicide respectively showed 100% shoot survival rate.



Fig. 1. Treatment T7 (3%NaOCl), T10 (5% NaOCl) showing fungal contamination after 14 days of culturing.

Results after 14 days (Table 3) showed that treatment T1 (0.1 % HgCl_2), T9 (3% NaOCl for 3 minutes), T14 (3% NaOCl + fungicide) and T23 (0.5% HgCl_2 +fungicide for 1 minute) showed 80 % survival rate.

However, results after 28 days of inoculation (Table

4) revealed that most of the treatments containing NaOCl as sterilizing agent showed 0% survival rate. According to Ahmed, *et al.*, (2014) considerable reduction in contamination of explant was obtained by treating the explants with sodium hypochlorite 5% for 10 minutes after rapid rinsing in 70% ethanol for 30 seconds.



Fig. 2. *Skimmia laureola* explants showing shoot emergence in treatment T4 (0.5 % mercuric chloride) as after 7 days of culturing.

In the present study 3% NaOCl along with fungicide for 1 minute also showed 20% contamination and 80% survived explants even after 28 days of culturing.

Death percentage

Data regarding death percentage of explants after 14 days of culturing (Table 3) showed that sterilization procedure T4, T5 (0.5 % HgCl_2 for 30 seconds and 1 minute, respectively) and T19, T22 having 0.1 % and

0.5 % HgCl_2 +fungicide for 30 seconds gave significantly ($P \leq 0.05$) lowest percentage of explants death i.e 0%. While treatment T7 (3 % NaOCl), treatment T10 (5 % NaOCl), T13 (3% NaOCl + fungicide) for 30 seconds showed 100% death of explants (Fig 3). Sterilization protocols T3 (0.1 % HgCl_2 for 3 minute), T12 (5% NaOCl), T15 (3% NaOCl + fungicide), T6 and T24(0.5 % HgCl_2 for 3 minutes) showed 80% death of explants.




Fig. 3. Fungal contamination on *Skimmia laureola* explants (a), causing explant death (b).

After 28 days of inoculation (Table 4) same trend was observed except treatment T3 (0.1 % HgCl₂ for 3 minute) which showed 100% explants death. Results similar to present study was also reported by Sen *et al.*, (2013) who found that at high concentration of HgCl₂ with fungicide controls contamination but cause corrugation to explants.

Percent discoloration

It was observed that the sterilization protocols NaOCl, NaOCl + fungicide at both concentration (3 % and

5%) for all time periods (30 seconds, 1minute and 3 minutes) gave significantly ($P \leq 0.05$) lowest discoloration percentage of explants (0%) (Table 4).

Similarly 0.5% HgCl₂ for two time period (30 seconds and 1 minute) and 3 % HgCl₂+ fungicide for 30 seconds also showed 0% discoloration. While significantly maximum discoloration i.e 80 % was observed in 0.1% HgCl₂ for 3 minutes (T3) (Fig. 4a), 0.5% HgCl₂ (T6) and 0.5% HgCl₂+fungicide treatments for 3 minute (Fig. 4b).




Fig. 4. Treatments T1 (a), and T24 (b) showing discoloration of explants after 14 days of culturing.

Our results are in accordance with the findings of Danso, *et al.*, (2011) who reported that exposure to HgCl₂ may have negative effects on survival rate of explants leads to browning and death of explants. The present study also showed deleterious effect of HgCl₂ at high concentration and long exposure.

Shoot length

The effect of sterilizing agents on growth of explants was recorded after 14 days of culturing (Table 3). The significantly ($P \leq 0.05$) maximum shoot length (2.24cm) was observed in treatment T4 (0.5% HgCl₂ for 30 seconds) (Fig. 5) followed by treatment T22 (0.5 % HgCl₂ + fungicide) 1.3 cm shoot length.



Fig. 5. *Skimmia laureola* explants showing shoot growth.

It was reported by Mahmoud and Nabeel, (2016) that at concentration (0.1%) mercuric chloride gave (90%) decontamination percentage and (60%) of viable nodal segments showing shoot development. No shoot growth was recorded in NaOCl treatments.

Results after 28 days of culturing (Table 4) showed an increase in shoot growth (3.26cm) in treatment T4 (0.5% HgCl₂ for 30 seconds) whereas treatment T5 (0.5% HgCl₂ 0.5% for 1 minute) and treatment T22 showed 1.74cm 1.50 cm respectively of shoot length. Our results are in line with the findings of Sen *et al.*, (2013) who found that at minimum concentration (0.1%) with the shortest duration of treatment (1 min), HgCl₂ was efficient as sterilizing agent giving 22.19% of *A. aspera* seed culture to be contamination free with a vigorous growth of plantlets of 5.67 cm in length was recorded.

Number of leaves

Data collected after 14 days (Table 3) showed that treatment T19 (0.1% HgCl₂ + fungicide for 30 seconds) showed maximum mean number of leaves i.e 3.2 followed by T4 and T22 in which 3 leaves per growing plantlets were observed. An increase in average leaf number was observed after 28 days of culturing (Table 4). This may be due to the fact that as the maximum number of plants remained survived in HgCl₂+fungicide treatment at both concentrations for 30 seconds (Table 3), these plants also showed growth in terms of shoot length and leaf number. However in literature direct effect of sterilizing agents on leaf number has not been reported.

Conclusion

Among the different sterilization protocols tested for the successful establishment of *in vitro* culture of *Skimmia laureola*, it has been concluded that mercuric chloride (HgCl₂) at 0.5 % concentration is most effective sterilizing agent when used alone or in combination with fungicide for controlling 100 % contamination with 100% survival rate for 30 seconds and 1 minute. However, 0.5 % HgCl₂ for 3 minutes showed 80 % discoloration and death of explants. It has been recommended that HgCl₂ alone or in combination with fungicide for short time of exposure

can be used as sterilant for sterilization of important medicinal plants and further work is needed for standardizing HgCl₂ concentrations and time of exposure for establishment of *in vitro* culture of endangered medicinal plants.

References

- Alam F, Ekhlas-Ud-din M, Ruhul Amin, Razzak AM, Manik MA, Khatun MM.** 2016. Studies on the effect of various sterilization procedure for *in vitro* seed germination and successful micropropagation of *Cucumis sativus*. International Journal of Pure Biosciences. **4(1)**, 75-81.
<http://dx.doi.org/10.18782/2320-7051.2226>
- Ali SI, Qaiser M.** 2007. Flora of Pakistan. Botany Deptt. Uni. of Karachi. **61**, 93-112.
- Ashfaq S, Ahmad M, Arshad M.** 2003. Ethnomedicinal observations of medicinally important plants of Tehsil Fateh Jang, District Attock. Pakistan Journal of Arid Agriculture **7(1)**, 25-33.
- Baart JLG, Baart-Bremer EL, Sagar MZ.** 2004. Names of plants in Kalam Kohistan (Pakistan) Work Papers of the Summer Institute of Linguistics, University of North Dakota Session[C]. 48.
- Barkatullah, Ibrar M, Muhammad N.** 2012. Phyto sociological and ethnobotanical attributes of *Skimmia laureola*. International Journal of Botany. **2**, 5-84.
- Bhattarai S, Chaudhary RP, Taylor RSL.** 2006. Ethnomedicinal plants used by the people of Manang District, central Nepal. Journal of Ethnobiology and Ethnomedicine **2**, 41. 3.
<https://doi.org/10.1186/1746-4269-2-41>
- Central Potato Research Institute (CPRI).** Shimla. 1992. Tissue Culture technique for potato health, conservation, micro propagation and improvement. CPRI, Shimla: 1-23.
- Danso KE, Azu E, Elegba W, Asumeng A, Amoatey HM, Klu, GYP.** 2011. Effective decontamination and subsequent plantlet regeneration of sugarcane (*Saccharum officinarum* L.) *in vitro*. International Journal of Integrative Biology **11**, 90-96.

- Daud NH, Jayaraman S, Mohamed R.** 2012. An improved surface sterilization technique for introducing leaf, nodal and seed explants of *Aquilaria malaccensis* from field sources into tissue culture. *AsPac Journal of Molecular Biology and Biotechnology*. **20(2)**, 55-58.
- Debnath M, Malik CP, Bisen PS.** 2006. Micropropagation: A Tool for the Production of High Quality Plant-based Medicines. *Current Pharmaceutical Biotechnology* **7**, 33-49. <https://doi.org/10.2174/138920106775789638>
- Dodds JH, Roberts LW.** 1985. Experiments in plant tissue culture (second edition). Cambridge University Press. 21-35.
- Dukan S, Belkin S, Touati D.** 1999. Reactive oxygen species are partially involved in the bactericidal action of hypochlorous acid. *Archives of Biochemistry and Biophysics*. **367**: 311–316. <https://doi.org/10.1006/abbi.1999.1265>
- Hamayun M.** 2007. Traditional uses of some medicinal plants of Swat Valley, Pakistan. *Indian Journal of Traditional Knowledge* **6**, 636–641.
- Hamayun M, Afzal S, Khan MA.** 2006. Ethnopharmacology, indigenous collection and preservation techniques of some frequently used medicinal plants of utror and gabral, district swat, Pakistan. *African journal of Traditional complementary and alternative medicines*. **3(2)**, 57-73.
- Hassan-ud-Din, Gzanfar SA.** 1980. Flora of Pakistan. family Rutaceae. Pakistan Agriculture Research Council, Islamabad. **132(1)**.
- Katky A, Handique PJ.** 2010. Micropropagation and screening of antioxidant potential of *Andrographis paniculata* (Burm. f) Nees. *Journal of Hill Agriculture* **1(1)**, 15-20.
- Martin KP.** 2004. Plant regeneration protocol of medicinally important *Andrographis paniculata* (Burm.f) Wallich ex Nees via somatic embryogenesis. *In Vitro Cellular and Developmental Biology-Plant*. **40**, 204-209. <https://doi.org/10.1079/IVP2003520>
- Mahmoud NS, Nabeel K, Al Ani.** 2016. Effect of Different Sterilization Methods on Contamination and Viability of Nodal Segments of *Cestrum nocturnum*. *International Journal of Research Studies in Biosciences*. **4**, 4-9. <http://dx.doi.org/10.20431/23490365.0401002>.
- Murashige T, Skoog F.** 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiology* **15**, 473-497. <https://doi.org/10.1111/j.13993054.1962.tb08052.x>
- Naika HR, Krishna V.** 2008. Plant regeneration from callus culture of *Clematis gouriana* Roxb. a rare medicinal plant. *Turkish Journal of Biology* **32**, 99-103.
- Okudera Y, Ito M.** 2009. Production of agar wood fragrant constituent in *Aquilaria calli* and cell suspension culture. *Plant Biotechnology*. **26**, 307-315. <https://doi.org/10.5511/plantbiotechnology.26.307>
- Pandey P, Rakesh M, Upadhyay, Ravi.** 2013 In-vitro propagation of an endangered medicinal plant *Psoralea corylifolia* Linn. *Asian Journal of Pharmaceutical and Clinical Research*. **6**, 115-118.
- Preethi D, Sridhar TM, Naidu CV.** 2011. Efficient protocol for indirect shoot regeneration from leaf explants of *Stevia rebaudiana* (Bert.)– an important calorie free bio sweetner. *Journal of Phytology* **3**, 56–60.
- Purkayastha J, Sugla T, Paul A, Solleti S, Sahoo L.** 2008. Rapid in vitro multiplication and plant regeneration from nodal explants of *Andrographis paniculata*: A valuable medicinal plant. *In Vitro Cellular and Developmental Biology-Plant*. **44(5)**, 442-447. <https://doi.org/10.1007/s11627-008-9156-8>
- Qin Y, Zeng F, Sun X, Yingli F, Cuiqin Y.** 2012. Propagation of *Cleome spenos*: a Jack through tissue culture. *Journal of Biotechnology and Food Science*. **1**, 1319-1327.

- Qureshi RA, Ghufran MA, Gilani S, Yousaf Z, Abbas G, Batool A.** 2009. Indigenous medicinal plants used by local women in Southern Himalayan regions of Pakistan. *Pakistan Journal of Botany*. **41**, 19.
- Schenk RV, Hilderbrandt AC.** 1972. Medium and techniques for induction and growth of monocotyledonous dicotyledonous plant cell cultures. *Canadian Journal of Botany* **50**, 199-204.
<https://doi.org/10.1139/b72-026>
- Sen MK, Jamal M, Abu HM, Nasrin S.** 2013. Sterilization factors affect seed germination and proliferation of *Achyranthes aspera* cultured in vitro. *Environmental and Experimental Biology*. **11**, 119–123.
- Sen MK, Hassan MM, Nasrin S, Jamal MAHM, Mamun-OrRashid ANM, Dash BK.** 2013. In vitro sterilization protocol for micropropagation of *Achyranthes aspera* L. node. *International Research Journal of Biotechnology* **4**, 89–93.
- Trombetta D, Castelli F, Sarpietro MG, Venuti V, Cristiani M, Daniele, Sajia, Mazzanti A, Bisignano.** 2005. Mechanism of antibacterial action of three monoterpenes. *Antimicrobial Agents and Chemotherapy* **49**, 2474–2478.
<https://doi.org/10.1128/AAC.49.6.2474-2478.2005>
- Webster S, Mitchell SA, Ahmad MH.** 2003. Anovel surface sterilization method for reducing fungal and bacterial contamination of field grown medicinal explants intended for in vitro culture. Proceedings of 17th SRC conference entitled ‘Science and Technology for Economic Development: Technology Driven Agriculture and Agro-Processing’ SRC Jamaica.
- Yadav K, Narender S, Sharuti V.** 2012. Plant tissue culture: a biotechnological tool for solving the problem of propagation of multipurpose endangered medicinal plants in India. *Journal of Agricultural Technology* **8**, 305-3184.
- Kiani M, Fatemeh SY, Mohammad HE, Pandarinath S, Phanikanth J, Abbagani S.** 2017. Studies on the Effect of Various Seed Surface Sterilization and Growing Media on the In-vitro Germination of Lemon Balm (*Melissa officinalis* L.). *Indian Journal of Science and Technology* **10(3)**, 1-6.
<https://doi.org/10.17485/ijst/2017/v10i3/102666>.