



RESEARCH PAPER

OPEN ACCESS

Seed germination in maize (*Zea mays* L.) under the influence of different drinks flavors as a source of plant nutrients

Ikram-ul Haq*, Noman Ali, Danish, Nazia Parveen Gill¹, Mahnoor Dua, Mariyam Shaikh

Institute of Biotechnology and Genetic Engineering (IBGE), University of Sindh, Jamshoro, Pakistan.

¹*Department of Statistics, University of Sindh, Jamshoro, Pakistan.*

Article published on March 30, 2018

Key words: Maize (*Zea mays* L.), Fruit juices, Seed germination, Reducing sugars, Antioxidants

Abstract

In present work, effect of different nutrients in the form of various drinks flavors available in local market on plants was assessed on especially seed germination of maize (*Zea mays* L.) var., MMRI-Yellow and its early growth. Seeds germinated on different nutrient conditions like as T₀ (dH₂O), T₁ (MAY-sparkling fruit juice), T₂ (1/4T₁), T₃ (Murree's Wheat Beer), T₄ (1/4T₃), T₅ (Apple Mixed Juice), and T₆ (1/4T₅) and maintained for 2-weeks. Among the morphological attributes maximum seed germination, seedlings biomass, lengths of plumules and radicals observed on T₀ and T₆ cultures and each are significantly lesser among seeds growing under alcoholic bears supplements (T₃ and T₄). The biochemical contents including chlorophyll (ab), total sugars and protein contents also showed same trend among the cultures ($p < 0.05$). Meanwhile, chlorophyll b, total carotenoids, reducing sugars, antioxidants activity and phenolics observed maximum in plumules and radicals of seedlings grew in T₁ and T₃ cultures ($p < 0.05$). These biocontents reduced among the dilutions to T₀ and T₆ cultures. These both cultures showed good seedling growth responses, whereas T₆ remained most growth supportive. It means that seed mobilization and its growth suppressed with supplementation of MAY-sparkling fruit juice (T₁) and Murree's Wheat Beer (T₃) even concentrated apple juice also (T₅). The T₁ and T₃ cultures showed impaired growth retardation during seed germination. Each of mixed fruit juice, non-alcoholic wine and alcoholic bears have performed significant role in seed germination and its initial subsequent growth.

*Corresponding Author: Dr Ikram-ul Haq ✉ rao.ikram@yahoo.com

Introduction

The maize (*Zea mays* L.) crop has got 2nd position among cereals after wheat in Pakistan (FAO, 2015). It is being used as human nutrition, livestock feed and its raw material in agro-based industries (Gwirtz and Garcia-Casal, 2014; Notenbaert *et al.*, 2013). A number of pressures exerted in the form of land limitation due to increasing population and plant pathogens are decreasing both of its quality and quantity. Consequently, approaches of conventional and modern biotechnology are emphasized. Conventional methods are economic and useful to get high yield from crops. On the basis of general characteristics of seeds that large sized seeds showed higher germination rate with its rapid emergence and further growth due to its large storage reservoirs in cotyledons (Alenca *et al.*, 2012; De Souza *et al.*, 1999; Seiwa and Kikuzawa, 1991).

The increasing rates of external applied environmental stresses are ultimately causing to decrease in crop yields including number of seeds or grain and its seed size. Large sized seeds have optimum raw material for its germination and initial growth of emerging shoots due to availability of substrate, enzymes and energy (ATP) for protein synthesis (Gallardo, 2002; Souza and Fagundes, 2014). *In-vitro* tissue cultures like as callus formation also dependent on seed size as well as its embryo's maturity (Akinyosoye *et al.*, 2015; Shah *et al.*, 2003). Meanwhile, a farmer is primarily believes that crop yields can be increased with applications of fertilizers, pesticides and good irrigation. For crop production, pesticide play an essential part to kill surface growing pathogen that makes possible to produce good yield to feed human as well as animals. Various varieties of different crops responses differently, when grown on the land with variant environmental biotic and abiotic stresses. The chemical stresses among abiotic stresses are known to exert negative impact on cell's metabolic processes, which results into reduction in vegetative growth of plant and its yield (Suzuki *et al.*, 2014; Verma *et al.*, 2016; Yumurtaci, 2015).

Meanwhile, little information about the effect of maize seed size on *in vitro* seed germination is available. Large sized seeds germinate nicely but

small sized seed shows low germination rate. It could be due to low number of essential components of seeds. The deficiency of essential components may be cause of slow down or reduction of seed germination. Supplements of nutrients could be helpful to enhance the rate of seed germination.

The plants are able to retrieve their nutrients from soil medium or other sediments including *in-vitro* cultures. The surplus supply of inorganic nutrients in the form of hydroponics could be useful for plant growth while being costly. Almost similar typed nutrients including inorganic and organic molecules are present in the fruit juices. These juices are being the cheapest source of various nutrients in balanced form and are available in the commercial super markets.

Instead to use the fresh available commercial juices, their wastes can be used for this purpose after proper sterilization. Various fruit industries have a considerable waste of their final products. Mostly wastes are not recycled just thrown away on earth or even drainage out in the water. Both are toxics for animals and plants on the soil as well as in the water. It is being a source of environmental pollution.

The proper management of such wastes in the form of bio-nutrients could be useful source of nutrition for other organisms. Like as different fruit juices have attractive composition of various salts, vitamins and sugars. Each component of the juice is also variable in types and concentrations from juice to juice of fruits even including wines and bears also.

For a specific crop or even its variety, the selection of a fruit juice is a major task. For this purpose, the aim of this study is to measure the effects of local drinks flavor (mixed fruit juice, non-alcoholic wine and alcoholic bear) on subject of interest of seed germination, early growth rate of newly emergence and further their metabolic rates. A maize var., MMRI-Yellow is subjected for detailed evaluation of its seed germination on the basis of their biochemical changes under the influence of variant concentrations of local commercial fruit drinks.

Materials and methods

Plant seed materials

Present experiment was conducted in glass petri-dishes. The bases of dishes were lined with sterile whatman-40 filter paper. Seeds of maize (*Zea mays* L.) var., MMRI-Yellow collected from local grain market. The seeds were rinsed in 90% ethanol for 1 minute than stirred in 0.1% mercuric chloride (HgCl₂). After 10 minutes, seeds were washed with sterilized distilled water for 5-8 times. The 5 seeds were arranged on sterile filter paper and subjected to fruit juice treatments.

Application of fruit juices treatments

Three local drinks namely MAY-sparkling fruit juice (non-alcoholic), Murree's Wheat Beers (5.3% alcohol) and Multivitamins mix-fruit juice were purchased from local supermarket.

Different concentrations of each drink were raised by mixing with sterile distilled water as shown in table 1. Exact 3ml of each treatment was poured over filter-paper. The cultures were incubated at 25°C±2, 25% humidity and 18/6 h day and light photoperiod (light intensity ~2000 lux) for 2-weeks.

Table 1. Various treatments and its composition used for seed germination in maize (*Zea maize* L.).

#s.	Treatments	Compositions (Nutrient contents)
a.	T ₀	Distilled water (dH ₂ O)
b.	T ₁	<i>May Sparkling Fruit Juice (240 ml)</i> : Sodium 10 mg, carbohydrate 26 mg, sugar 26 mg, vitamin C 60%, iron 8%, calcium 2%.
c.	T ₂	¼ T ₁
d.	T ₃	<i>Murree's wheat beer (240 ml)</i> : Calories 160, alcohol 5.3 %, fat 1 g, sodium 320 mg, carbohydrate 31 mg, dietary fiber 2 g, sugar 9 g, protein 4 g.
e.	T ₄	¼ T ₃
f.	T ₅	Mixed juice (240 ml): 45 calories, fat 0.1 g, sodium 4 mg, potassium 101 mg, carbohydrate 11.6 g, dietary fiber 0.1 g, sugar 8.5 g, protein 0.1 g, vitamin A 1 %, vitamin C 106%, calcium 1 %, iron 2 %, thiamin 1 %, riboflavin 1 %, vitamin B ₆ 2 %, niacin 1 %, magnesium 1 %, phosphorus 1 %, copper 2 %, pantothenic acid 1 %.
g.	T ₆	¼ T ₅

Maize seed germination

Seeds of each culture were kept under moisture conditions with their respective concentration of drinks. After 2-weeks of seed incubation in plant growth room, seeds were examined day by day and number of germinated seeds were recorded. On 15th day, seedlings from each culture were harvested and subjected for fresh biomass measurements after counting germinated seeds. Both plumules and radicals lengths were taken and seedlings were dried in the electric oven at 72°C. On 3rd day, seedling's dry biomass was also measured.

Biochemical analysis of seedlings

Seedlings of each culture were subjected for analysis of various biochemical contents. Fresh seedlings were chopped into smallest fine pieces and agitated in 80% acetone. Mixture was incubated in dark at room temperature for overnight. Absorbance were taken at 663 nm, 645nm, 453nm and 470nm and various chlorophyll contents and total carotenoids were calculated by following Arnon (chlorophyll) and

Sumata *et al* (carotenoids) suggested formulas (Arnon, 1949; Sumanta *et al.*, 2014). For the estimation of total proteins, exact 1 ml extract of fresh seedlings was mixed with 2.5ml alkaline copper reagent and after 0.25ml follin-ciocalteau reagent also poured slowly than Absorbance was measured at 750nm (Dawson and Heatlie, 1984). Total sugars were analyzed by following Lee and Montgomery's method (1961). The 0.5ml sample mixed with 2.50ml conc. H₂SO₄ than 50µl 80% phenol was added. After 15 minutes OD was read at 485nm. Also for reducing sugars, 1ml of sample mixed with 1ml 2, 6-dinitrosalicylic acid (DNS) and heated for 5 minutes in boiling water bath. When cool-down, the absorbance was read at 540nm (Miller, 1959).

Estimation of enzyme activities

Among the seedlings cultures, antioxidants activity was determined by mixing 10µl of sample with reaction mixtures and it was capped with aluminum foil. It was heated on boiling water bath for 90 minutes.

After cooling, the absorbance was read at 734nm (Pisoschi and Negulescu, 2012; Prior *et al.*, 2005; Proestos *et al.*, 2013). *Protease* activities was estimate in 2.0ml sample after mixing with 1.5ml sodium phosphate buffer (pH 7.6) by following methods reported by Alef and Nannipieri, (1995) and Vágnerová and Macura, (1974).

Decontaminated conditions

All steps of this experiment are performed under sterilized conditions on laminar air-flow cabinet. The glass-ware i.e. petri-dishes, volumetric flasks, filter-paper and other stuff used in this experiment cleaned with vim and then sterilized at 121°C, 15 lbs/cm² in electric-autoclave for 15 minutes. Autoclaved stuff was dried in electric oven (65°C, for 4-6 hours) and was opened to use in laminar air flow cabinet.

Statistical analysis of data

Data was collected from each seedling cultures and subjected for treatment significance with analysis of variance (ANOVA) on CoStat (3.03) *CoHort* software, Berkeley (USA). Mean differences between treatments also calculated with Duncan Multiple Range (DMR) test at 5% (Behrens, 1997; Henley, 1983).

Results and discussion

The maize (*Zea mays* L.) cereal is cropped widely due to its increasing demand to reach human consumption. With the passage of time its quality and quantity is decreasing with changing environmental stresses especially seed health (Kansiime and Mastenbroek, 2016; Liu *et al.*, 2016). The seeds with poor health cause inhibition for its germination and remain susceptible to the attacks of pathogens and insects (Donohue *et al.*, 2010; Shepherd and Chapman, 1998; Toh *et al.*, 2008).

Influences of other nutrient alternates could facilitate seed germination like as soil conditions either with sufficient nutrients or its physical texture (Rivera-Aguilar *et al.*, 2005). Soil fertility remains important for seed germination, its further development (Venable and Lawlor, 1980). Seed germination response depends on available quantity and sources of supplements of essential plant nutrient element (Pérez-García and González-Benito, 2006).

Seed germination responses may quite be associated with the soil nutritional level. In this experiment various nutrient cultures are established for the cultivation of MMRI-Yellow var., of maize. Composition of each seed germination culture raised with local fruit based drinks (Table 1). A significantly differential pattern in seed germination as well as further growth of seedling among the treated to control cultures is observed. Maximum seed germination observed in control (T₀) and mixed juice treated cultures (T₆) as 93.33% and 86.66% respectively. It is decreased in the cultures supplemented with MAY - sparkling fruit juice and Murree's wheat bear either supplied in concentrated or diluted forms ($p < 0.05$). Similar growth pattern after seed germination, seedling parts like as length of plumules and radicals even seedlings F. Wt. and its D. Wt. has been observed (Table 2).

Seed germination and its subsequent growth is impaired with various biochemical processes of plant metabolism, which consequently regulate their growth rate (Ashraf and Foolad, 2005; Kranner and Colville, 2011; Miransari and Smith, 2014).

The local drinks like as commercial fruit mixed juices, and beers even vines are based on different components of nutrients. In this way each juice could show different impact on plant or seed growth including its germination even it is best for plant callusing cultures (Haq *et al.*, 2011; Mgaya *et al.*, 2014; Nour *et al.*, 2012). It means that reduction or enhancement of plant biomass depends on availability of proper nutrition. While higher concentrations of juices may inhibit the germination or its further growth due to adverse effects of juices on degradation as well as mobilization of seed reserves (Djossa *et al.*, 2008; Evenari, 1949; Gardner *et al.*, 2000).

Morphological phenotypes of germinated seeds and its seedlings are expressions developed due to undergoing biochemical changes in the seed reserves. These are specific markers of seedlings growth and their development for which various enzymes of internal tissues are hydrolyzing seed reserve to convert complex cellular and organelle's molecules into simplest forms.

However, variations in rates of conversion of molecules into others depend on crops, species as well as external environmental conditions (Dürr *et al.*, 2015; Kakhki *et al.*, 2011; Kerrison *et al.*, 2015). Such as seed proteins are major component of seed reserve among the plant species, which are hydrolyzed with proteolytic enzymes at seed germination stage.

Proteins including other like as carbohydrates and lipids plays pivotal role being a source of nitrogen, carbon and energy for growing tissues. External nutritional sources could play a role in activation of hydrolytic enzymes, source of nutrition or even delay or inhibit the seed germination (Bragina *et al.*, 2003; Smykal *et al.*, 2014; Turner, 2010).

Table 2. Various morpho-biochemical attributes of maize (*Zea mays* L.) seedlings germinated under moisture conditions raised with local drinks (a 2-weeks culture).

S	Characteristics	T ₀	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	P-sig.
a	Germination rate (%)	^a 93.33±6.667	^c 53.33±6.667	^{bc} 13.33±6.667	^d 26.66±6.667	^d 26.66±6.667	^{bc} 66.66±6.667	^{ab} 86.66±6.667	***
b	Plumule length (cm)	^b 7.333±0.273	^d 4.233±0.203	^c 5.100±0.115	^b 2.833±0.088	^{ab} 3.833±0.088	^{ab} 5.367±0.145	^{ab} 8.167±0.145	***
c	Radical length (cm)	^a 8.400±0.173	^c 5.733±0.233	^b 6.900±0.173	^c 3.800±0.173	^d 4.667±0.145	^b 7.167±0.145	^a 8.833±0.088	***
d	Seedlings F.Wt (g)	^a 9.400±0.095	^c 5.004±0.099	^d 5.403±0.123	^f 4.814±0.039	^e 4.813±0.090	^c 8.921±0.090	^a 9.889±0.061	***
e	Seedlings D.Wt (g)	^a 1.440±0.026	^c 0.890±0.009	^b 1.022±0.033	^e 0.687±0.008	^d 0.782±0.009	^b 0.991±0.009	^a 1.501±0.008	***
f	Chlorophyll a (mg/g)	^a 2.139±0.381	^c 1.622±0.029	^b 1.778±0.020	^e 1.106±0.013	^d 1.210±0.059	^b 1.751±0.033	^a 2.227±0.006	***
g	Chlorophyll b (mg/g)	^a 0.994±0.006	^c 0.770±0.009	^b 0.801±0.007	^d 0.547±0.005	^c 0.749±0.007	^b 0.814±0.008	^a 1.009±0.010	***
h	Chlorophyll ab (mg/g)	^b 3.133±0.043	^d 2.393±0.023	^c 2.577±0.013	^f 1.653±0.011	^e 1.959±0.053	^c 2.565±0.026	^a 3.236±0.005	***
i	Carotenoids (mg/g)	^f 2.249±0.015	^b 4.261±0.058	^d 3.254±0.031	^a 4.401±0.006	^c 3.489±0.035	^c 3.508±0.018	^e 2.665±0.072	***
j	Total sugars (mg/ml)	^{ab} 2.884±0.009	^c 2.229±0.006	^{bc} 2.571±0.012	^d 1.782±0.329	^c 2.347±0.014	^c 2.442±0.011	^c 2.975±0.006	***
k	R. sugars (mg/ml)	^d 0.671±0.012	^b 0.829±0.006	^c 0.771±0.012	^a 0.915±0.006	^b 0.847±0.014	^d 0.682±0.005	^b 0.559±0.003	***
l	Total proteins (mg/ml)	^a 3.717±0.041	^d 2.471±0.012	^c 2.529±0.006	^f 2.047±0.014	^e 2.113±0.013	^d 2.435±0.003	^b 2.765±0.017	***
m	Phenolics (mg/ml)	^f 1.268±0.012	^c 1.529±0.006	^d 1.465±0.006	^a 1.728±0.006	^b 1.647±0.014	^c 1.382±0.005	^f 1.280±0.004	***
n	AOA (mmolTE/g F.Wt)	^e 0.682±0.005	^b 0.872±0.003	^c 0.838±0.006	^a 0.895±0.006	^b 0.860±0.004	^d 0.730±0.003	^f 0.665±0.003	***

T₀: Distilled water control; T₁: MAY-sparkling fruit juice; T₂: ¼T₁ + distilled water; T₃: Murree's Wheat Beer; T₄: ¼T₃ + distilled water; T₅: Apple Mixed Juice; T₆: ¼T₅ + + distilled water; AOA: Antioxidants activity; R. sugars: Reducing sugars; *p*-sig: *p* significance; F.Wt. Fresh weight; D.Wt.: Dry weight.

Among the seed reserves, carbohydrates (exist as free sugar and polysaccharides-starch) are also being an important components available as essential energy rich sources for seed germination. Starch is degraded by α -amylase into soluble sugars after see germination (Bernal-Lugo Smykal Leopold, 1992; Hagely *et al.*, 2013; Hedge and Hofreiter, 1962). A differential pattern of total sugars observed among the seedlings maximum in T₀ and T₆ cultures, while minimum in T₃ and T₁ cultures. Similar pattern is also seen for total proteins, chlorophyll a and even overall total chlorophyll (ab) contents. Reduction in rate of starch degradation may be due to inhibition of hydrolytic enzymes lowered the seedlings growth rate. After seeds germination still rate of growth of seedling remained lowered in the T₃ and T₁ cultures, while observed maximum in T₀ and T₆ cultures. It might be declarable that after seed germination, seedlings are growing further under stress of supplied nutrients in the form of fruit juices. Stressed conditions are provable in the form of increase in reducing sugars, chlorophyll b, phenolics and

antioxidants among the cultures (Table 2). Alleviation of these contents in seedling or plants is being their ability against applied stresses and remained at the same level until stress is passed over to normal conditions (Ali and Alqurainy, 2006; Michalak, 2006; Singh *et al.*, 2015). Anyhow initial nutrition have a key impact on seed germination, which is achievable with the optimization of initial nutrition condition for early seed growth of each crop.

Conclusions

Seed germination is an important step of early plant growth. Healthy seeds are germinated easily without delay than small sized or week seeds. After seedling stage, further growth potential depends on health of both typed seeds. Best vegetative as well as reproductive growth is achievable with supplementation of nutrients required for seed germination and its early growth. In this study, differential rates of seed germination and its further early seedling growth has observed among the maize var., MMRI-Yellow seeds of 2-weeks culture.

A similar and comparative seed germination and seedlings biomass attributes observed in control (distilled water) and mixed apple fruit juice (1/4 dilution). Other seedlings (on alcoholic beer, non-alcoholic wine) showed reduction in plant growth rates because of imbalanced nutritional stress, which increases in osmoprotective among the cultures. Proper nutritional conditions at seed germination stage could be helpful to empower the seedlings for its further growth stages.

Acknowledgements

We are in cordial thankful for support to provide chemicals and glassware to University of Sindh. We also like to pay thanks to people from IBGE for proving managerial technical as well as clerical help whenever we needed for this research work.

References

- Akinyosoye ST, Adetumbi JA, Amusa OD, Olowolafe MO, Olasoji JO.** 2015. Effect of seed size on *in vitro* seed germination, seedling growth, embryogenic callus induction and plantlet regeneration from embryo of maize (*Zea mays* L.) seed. Nigerian Journal of Genetics **2015(2)**, 1-7. <https://doi.org/10.1016/j.nigjg.2015.06.001>
- Alef K, Nannipieri P.** 1995. *Protease* activity. In Methods in applied soil microbiology and biochemistry. Academic Press, pp. 576. <https://doi.org/10.1016/B978-012513840-6/50022-7>
- Alenca NLM, Innecco R, Gomes-Filho E, Gallao MI, Alvarez-Pizarro JC, Prisco JT, de Oliveira AB.** 2012. Seed reserve composition and mobilization during germination and early seedling establishment of *Cereus jamacaru* DC, ssp. *jamacaru* (*Cactaceae*). Anais Da Academia Brasileira de Ciencias **84(3)**, 823-832. <https://doi.org/10.1590/S0001-37652012000300024>
- Ali AA, Alqurainy F.** 2006. Activities of antioxidants in plants under environmental stress. In: Motohashi N (ed) The Lutein-prevention and treatment for diseases. Transworld Research Network, India pp. 187-256.
- Arnon DI.** 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. Plant Physiology **24(1)**, 1-15. <https://doi.org/10.1104/pp.24.1.1>
- Ashraf M, Foolad MR.** 2005. Pre-sowing seed treatment-A shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions. Advances in Agronomy **88**, 223-271. [https://doi.org/10.1016/S0065-2113\(05\)88006-X](https://doi.org/10.1016/S0065-2113(05)88006-X)
- Behrens JT.** 1997. Principles and procedures of exploratory data analysis. Psychological Methods **2(2)**, 131-160. <https://doi.org/10.1037/1082-989X.2.2.131>
- Bernal-Lugo I, Leopold C.** 1992. Changes in soluble carbohydrates during seed storage. Plant Physiology **98(3)**, 1207-1210. <https://doi.org/10.1104/pp.98.3.120>
- Bragina TV, Rodionova NA, Grinieva GM.** 2003. Ethylene production and activation of hydrolytic enzymes during acclimation of maize seedlings to partial flooding. Russian Journal of Plant Physiology **50**, 794-798. <https://doi.org/10.1023/B:RUPP.0000003277.22914.6c>
- Dawson JM, Heatlie PL.** 1984. Lowry method of protein quantification: Evidence for photosensitivity. Analytical Biochemistry **140(2)**, 391-393. [https://doi.org/10.1016/0003-2697\(84\)90183-0](https://doi.org/10.1016/0003-2697(84)90183-0)
- De Souza AF, De Andrade ACS, Ramos FN, Loureiro MB.** 1999. Ecophysiology and morphology of seed germination of the neotropical lowland tree *Genipa americana* (Rubiaceae). Journal of Tropical Ecology **15(5)**, 667-680. <https://doi.org/10.1017/S026646749900108X>
- Djossa BA, Fahr J, Kalko EKV, Sinsin BA.** 2008. Fruit selection and effects of seed handling by flying foxes on germination rates of shea trees, a key resource in northern Benin, West Africa. Ecotropica **14**, 37-48. http://www.gtoe.de/PDF/Ecotropica_2008/Djossa_et_al_Ecotropica_14-1_2008.pdf

- Donohue K, de Casas R, Burghardt L, Kovach K, Willis CG.** 2010. Germination, postgermination adaptation, and species ecological ranges. *Annual Review of Ecology Evolution and Systematics* **41(1)**, 293-319. <https://doi.org/10.1146/annurev-ecolsys-102209-144715>
- Dürr C, Dickie JB, Yang XY, Pritchard HW.** 2015. Ranges of critical temperature and water potential values for the germination of species worldwide: Contribution to a seed trait database. *Agricultural and Forest Meteorology* **200**, 222-232. <https://doi.org/10.1016/j.agrformet.2014.09.024>
- Evenari M.** 1949. Germination inhibitors. *The Botanical Review* **15(3)**, 153-194. <https://doi.org/10.1007/BF02861721>
- FAO.** 2015. *FAO Statistical Pocketbook. Food and Agriculture Organization of the United Nations.* <https://doi.org/978-92-5-108802-9>
- Gallardo K.** 2002. Proteomics of *Arabidopsis* seed germination. A Comparative study of wild-type and gibberellin-deficient seeds. *Plant Physiology* **129(2)**, 823-837. <https://doi.org/10.1104/pp.002816>
- Gardner PT, White TAC, McPhail DB, Duthie GG.** 2000. The relative contributions of vitamin C, carotenoids and phenolics to the antioxidant potential of fruit juices. *Food Chemistry* **68(4)**, 471-474. [https://doi.org/10.1016/S0308-8146\(99\)00225-3](https://doi.org/10.1016/S0308-8146(99)00225-3)
- Gwartz JA, Garcia-Casal MN.** 2014. Processing maize flour and corn meal food products. *Annals of the New York Academy of Sciences* **1312(1)**, 66-75. <https://doi.org/10.1111/nyas.12299>
- Hagely KB, Palmquist D, Bilyeu KD.** 2013. Classification of distinct seed carbohydrate profiles in soybean. *Journal of Agricultural and Food Chemistry* **61(5)**, 1105-1111. <https://doi.org/10.1021/jf303985q>
- Haq I, Zara, Taseer ZA, Rajput MT, Dahot MU, Mustafa G, Mukesh, Latif A.** 2011. Effects of different fruit juices used as carbon source on cucumber seedling under *in-vitro* cultures. *African Journal of Biotechnology* **10(38)**, 7404-7408. <https://doi.org/10.5897/AJB10.2322>
- Hedge JE, Hofreiter BT.** 1962. Carbohydrates chemistry. Vol **17**, Whistler RL and Be Miller JN (Eds), Academic Press, New York, USA.
- Henley S.** 1983. Principles and procedure of statistics: In A biometrical approach. *Computers & Geosciences*, Steel RGD, Torrie JH, McGraw-Hill, New York, (2nd Ed), pp. 633. <https://doi.org/10.1016>
- Kakhki AM, Amoozegar MA, Khaledi EM.** 2011. Diversity of hydrolytic enzymes in haloarchaeal strains isolated from salt lake. *International Journal of Environmental Science and Technology* **8(4)**, 705-714. <https://doi.org/10.1007/BF03326255>
- Kansiime MK, Mastenbroek A.** 2016. Enhancing resilience of farmer seed system to climate-induced stresses: Insights from a case study in West Nile region, Uganda. *Journal of Rural Studies* **47**, 220-230. <https://doi.org/10.1016/j.jrurstud.2016.08.004>
- Kerrison PD, Stanley MS, Edwards MD, Black KD, Hughes AD.** 2015. The cultivation of European kelp for bioenergy: Site and species selection. *Biomass and Bioenergy* **80**, 229-242. <https://doi.org/10.1016/j.biombioe.2015.04.035>
- Kranner I, Colville L.** 2011. Metals and seeds: Biochemical and molecular implications and their significance for seed germination. *Environmental and Experimental Botany* **72(1)**, 93-105. <https://doi.org/10.1016/j.envexpbot.2010.05.005>
- Lee YC, Montgomery R.** 1961. The carbohydrate of ovalbumin. *Archives of Biochemistry and Biophysics* **95(2)**, 263-270. <https://doi.org/10.1016/0003-9861>
- Liu J, Liu Q, Yang H.** 2016. Assessing water scarcity by simultaneously considering environmental flow requirements, water quantity, and water quality. *Ecological Indicators* **60**, 434-441. <https://doi.org/10.1016/j.ecolind.2015.07.019>
- Mgaya KB, Remberg S, Chive B, Wicklund T.** 2014. Physio-chemical, mineral composition and antioxidant properties of roselle (*Hibiscus sabdariffa* L.) extract blended with tropical fruit juices. *African Journal of Food, Agriculture, Nutrition and Development* **14(3)**, 8963-8978.

- Michalak A.** 2006. Phenolic compounds and their antioxidant activity in plants growing under heavy metal stress. *Plant Cell*, **15(4)**, 523-530. <https://doi.org/10.1016/j.fitote.2011.01.018>
- Miller GL.** 1959. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry* **31(3)**, 426-428. <https://doi.org/10.1021>
- Miransari M, Smith DL.** 2014. Plant hormones and seed germination. *Environmental and Experimental Botany* **99**, 110-121. <https://doi.org/10.1016/j.envexpbot.2013.11.005>
- Notenbaert A, Herrero M, De Groot H, You L, Gonzalez-Estrada E, Blummel M.** 2013. Identifying recommendation domains for targeting dual-purpose maize-based interventions in crop-livestock systems in East Africa. *Land Use Policy* **30(1)**, 834-846. <https://doi.org/10.1016/j.landusepol.2012.06.016>
- Nour EH, Hamza MA, Fayez M, Monib M, Ruppel S, Hegazi NA.** 2012. The crude plant juices of desert plants as appropriate culture media for the cultivation of rhizospheric microorganisms. *Journal of Advanced Research* **3(1)**, 35-43. <https://doi.org/10.1016/j.jare.2011.03.002>
- Pérez-García F, González-Benito ME.** 2006. Seed germination of five *Helianthemum* species: Effect of temperature and presowing treatments. *Journal of Arid Environments* **65(4)**, 688-693. <https://doi.org/10.1016/j.jaridenv.2005.10.008>
- Pisoschi AM, Negulescu GP.** 2012. Methods for total antioxidant activity determination: A review. *Biochemistry and Analytical Biochemistry* **1(1)**, 1-10. <https://doi.org/10.4172/2161-1009.1000106>
- Prior RL, Wu X, Schaich K.** 2005. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry* **53(10)**, 4290-4302. <https://doi.org/10.1021/jf0502698>
- Proestos C, Lytoudi K, Mavromelanidou OK, Zoumpoulakis P, Sinanoglou VJ.** 2013. Antioxidant capacity of selected plant extracts and their essential oils. *Antioxidants* **2013(2)**, 11-22. <https://doi.org/10.3390/antiox2010011>
- Rivera-Aguilar V, Godínez-Alvarez H, Manuell-Cacheux I, Rodríguez-Zaragoza S.** 2005. Physical effects of biological soil crusts on seed germination of two desert plants under laboratory conditions. *Journal of Arid Environments* **63(1)**, 344-352. <https://doi.org/10.1016/j.jaridenv.2005.03.012>
- Seiwa K, Kikuzawa K.** 1991. Phenology of Tree Seedlings in Relation to Seed Size. *Canadian Journal of Botany-Revue Canadienne De Botanique* **69(3)**, 532-538. <https://doi.org/10.1139/b91-072>
- Shah MI, Jabeen M, Ilahi I.** 2003. *In vitro* callus induction, its proliferation and regeneration in seed explants of wheat (*Triticum aestivum* L.) var. LU-26S. *Pakistan Journal of Botany* **35(2)**, 209-217.
- Shepherd VE, Chapman CA.** 1998. Dung beetles as secondary seed dispersers: Impact on seed predation and germination. *Journal of Tropical Ecology* **14(2)**, 199-215. <https://doi.org/10.1017/S026646749800016>
- Singh M, Kumar J, Singh S, Singh VP, Prasad SM.** 2015. Roles of osmoprotectants in improving salinity and drought tolerance in plants: A review. *Reviews in Environmental Science and Biotechnology* **14(3)**, 407-426. <https://doi.org/10.1007/s11157-015-9372-8>
- Smykal P, Vernoud V, Blair MW, Soukup A, Thompson RD.** 2014. The role of the testa during development and in establishment of dormancy of the legume seed. *Frontiers in Plant Science* **5**, 351. <https://doi.org/10.3389/fpls.2014.00351>
- Souza ML, Fagundes M.** 2014. Seed size as key factor in germination and seedling development of *Copaifera langsdorffii* (Fabaceae). *American Journal of Plant Sciences* **5**, 2566-2573. <https://doi.org/10.4236/ajps>

- Sumanta N, Haque CI, Nishika J, Suprakash R.** 2014. Spectrophotometric analysis of chlorophylls and carotenoids from commonly grown fern species by using various extracting solvents. *Research Journal of Chemical Sciences* **4(9)**, 2231-2236. <https://doi.org/10.1055/s-0033-1340072>
- Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R.** 2014. Abiotic and biotic stress combinations. *New Phytologist* **203(1)**, 32-43. <https://doi.org/10.1111/nph.12797>
- Toh S, Imamura A, Watanabe A, Nakabayashi K, Okamoto M, Jikumaru Y, Kawakami N.** 2008. High temperature-induced abscisic acid biosynthesis and its role in the inhibition of gibberellin action in *Arabidopsis* seeds. *Plant Physiology* **146(3)**, 1368-1385. <https://doi.org/10.1104/pp.107.113738>
- Turner BL.** 2010. Variation in pH optima of hydrolytic enzyme activities in tropical rain forest soils. *Applied and Environmental Microbiology* **76(19)**, 6485-6493. <https://doi.org/10.1128/AEM.00560-10>
- Vágnerová K, Macura J.** 1974. Determination of *protease* activity of plant roots. *Folia Microbiologica* **19(4)**, 322-328. <https://doi.org/10.1007/BF0287322>
- Venable DL, Lawlor L.** 1980. Delayed germination and dispersal in desert annuals: Escape in space and time. *Oecologia* **46(2)**, 272-282. <https://doi.org/10.1007/BF00540137>
- Verma V, Ravindran P, Kumar PP.** 2016. Plant hormone-mediated regulation of stress responses. *BMC Plant Biology* **16(1)**, 1-10. <https://doi.org/10.1186/s12870-016-0771-y>
- Yumurtaci A.** 2015. Utilization of wild relatives of wheat, barley, maize and oat in developing abiotic and biotic stress tolerant new varieties. *Emirates Journal of Food and Agriculture* **27(1)**, 1-23. <https://doi.org/10.9755/ejfa.v27i1.17852>