



RESEARCH PAPER

OPEN ACCESS

Evaluation of the toxic and physiological activities of Precocene I (Anti-juvenile hormone agent) against *Spodoptera littoralis* Bois. (Lepidoptera: Noctuidae)

Khalid Hamadah, Karem Ghoneim*

Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt

Key words: Adult, Development, Growth, Larva, Mortality, Pupa.

<http://dx.doi.org/10.12692/ijb/13.1.86-105>

Article published on July 15, 2018

Abstract

The cotton leafworm, *Spodoptera littoralis* is a destructive pest of cotton and various crops all over the year in Egypt as well as in some countries. Objective of the present study was to assess the effects of Precocene I (PI) on survival, growth, development and metamorphosis of this pest. Five doses: 150, 120, 90, 30 and 15 µg/larva had been topically applied (once) onto 1-day old penultimate instar larvae or 1-day old last instar larvae. Various mortality percentages were recorded among the larvae, pupae and adults. LD₅₀ values were calculated in 70.48 and 234.96 µg/larva, after treatment of 5th and 6th larval instars, respectively. PI exerted a diverse action on the maximal body weight (max. wt) of treated larvae, depending on the dose. The larval growth was considerably inhibited. The larval duration was conspicuously shortened while the pupal duration was slightly or significantly prolonged, depending on the dose. Regardless the larval instar under treatment, the developmental rate of pupae was generally regressed. PI failed to induce the precocious pupation, regardless the applied dose. PI exerted a strong inhibitory action on the pupation, regardless the larval instar under treatment. The adult emergence was partially blocked. Some morphologically abnormal pupae were produced after treatment of 6th instar larvae with the higher two doses.

*Corresponding Author: Karem Ghoneim ✉ karemghoneim@gmail.com

Introduction

In Egypt, the cotton leafworm, *Spodoptera littoralis* Boisduval (Lepidoptera: Noctuidae) has been considered as destructive phytophagous pest of cotton and several vegetable and field crops all over the year (Hosny *et al.*, 1986; Shonouda and Osman, 2000; El-Khawas and Abd El-Gawad, 2002; Adham *et al.*, 2009). The insect attacks almost 112 plant species belonging to 44 families in tropical and temperate regions of the old world (Magd El-din and El-Gengaihi, 2000).

For controlling this pest, different types of conventional insecticides have been used over the past 40 years (Casida and Quistad, 1998). The intensive and indiscriminate use of insecticides against this pest results in the development of resistance to many registered pesticides making their control even more difficult (Ishaaya *et al.*, 1995; Smagghe *et al.*, 1999; Miles and Lysandrou, 2002; Aydin and Gurkan, 2006; Davies *et al.*, 2007; Mosallanejad and Smagghe, 2009).

As a result of improper and excessive uses, also, these insecticides usually exhibit several adverse impacts on the human health and beneficial animals as well as cause serious toxicological problems to the environment because these chemicals have a long half-life and retention in the environmental systems for long time (Van Der Gaag, 2000; Costa *et al.*, 2008; Relyea, 2009; Tiryaki and Temur, 2010). In addition, these chemicals have a tendency to accumulate in different trophic levels of the food net (Damalas and Eleftherohorinos, 2011; Chowański *et al.*, 2014).

Therefore, eco-friendly chemicals have received a great attention in the world during the recent years. These alternative compounds have lower toxicity to non-target organisms than conventional insecticides and they are effective at low concentrations (Attathom, 2002; Gade and Goldsworthy, 2003). Also, they are biodegradable into harmless compounds, and hence the problems of environmental pollution can be avoided (Tiryaki and Temur, 2010; Walkowiak *et al.*, 2015; Li *et al.*, 2017).

Owing to the dangerous attacks of *S. littoralis*, it is a subject to intensive research for searching new measures to control it and for improving the effects of known control methods (Hussain, 2012).

At present, the use of insect growth regulators (IGRs) is considered as possible alternative agents of conventional insecticides for controlling this pest (Raslan, 2002). IGRs are regarded as a 'third generation of insecticides' because they differ in their mode of action from other insecticides and have low toxicity to non-target organisms (Zhou *et al.*, 2003). Because of their desirable characteristics, such as less environmental pollution, high selectivity, and low impact on natural enemies and people, IGRs are used to control various insects (Wu, 2002; Cedric, 2005; Wang and Wang, 2007).

Juvenile hormone (JH) is necessary for the developmental stages in insects (Staal, 1986). Also, JHs play important roles in other physiological processes, such as behaviour, polymorphism, migration, metabolism, reproduction, diapause, and innate immunity (Riddiford, 1994; Gilbert *et al.*, 2000; Mitsuoka *et al.*, 2001; Tatar *et al.*, 2001a,b; Truman and Riddiford, 2007; Riddiford, 2008; Flatt *et al.*, 2008; Denlinger *et al.*, 2012; Amsalem *et al.*, 2014a).

The anti-juvenile hormone agents that affect either the mevalonate pathway in JH biosynthesis can be considered insect growth regulators (IGRs) affecting the corpora allata (CA) directly (Staal, 1986). Precocenes are originally plant-derived chromenes (2H-1-benzopyran) with insecticidal activities (Bowers, 1976; Proksch *et al.*, 1983; Isman *et al.*, 1986).

In the context of exploring the growth regulatory activities of compounds related to precocenes, rational synthetic methods for 2, 2-dimethyl chromenes were developed for the preparation of precocene analogues or precocenoids (Banerji and Goomer, 1984; Banerji and Kalena, 1989; Szczepanik *et al.*, 2005; Banerjee *et al.*, 2008).

Some entomologists paid a great attention to precocene analogues due to their twin advantage; avoiding the former surgical allatectomy and as an effective agent in devising 'fourth generation insecticides' in future (Muraleedharan *et al.*, 1986; Sariaslani *et al.*, 1987; Moya *et al.*, 1997; Szczepanik *et al.*, 2005; Singh and Kumar, 2011).

The hemimetabolous insects, such as grasshoppers, aphids and cockroaches, are more susceptible to precocenes than some larvae of holometabolous insects (Burt *et al.*, 1979; Chênevert *et al.*, 1980; Kiss *et al.*, 1988). However, some authors (Mathai and Nair, 1984; Kozhanova and Nemeč, 1991; Khafagi and Hegazi, 2001) reported that a number of holometabolous insects, such as *Spodoptera mauritia*, *Tenebrio molitor* and *Spodoptera littoralis* are sensitive to Precocenes.

It has been demonstrated that the design of anti-JH agents is an effective tool for insecticide discovery (Bede *et al.*, 2001). In addition, compounds with anti-JH activity are considered as new representatives of IGRs lacking some disadvantages of juvenoid-type chemicals (Bowers, 1982; Staal, 1982).

These compounds are potentially effective for controlling the major insect pests, since the larval stage is the dangerous among the insect developmental stages (El-Ibrashy, 1982). Objective of the present study was to assess the effects of Precocene I on the survival, growth, development metamorphosis and morphogenesis of *S. littoralis*.

Materials and methods

Experimental insect

A pupal sample of the Egyptian cotton leafworm, *Spodoptera littoralis* Boisduval (Lepidoptera: Noctuidae) was kindly obtained from the culture of susceptible strain maintained for several generations in Plant Protection Research Institute, Agricultural Research Center, Doqqi, Giza, Egypt. In Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, a culture was raised under laboratory controlled conditions (27±2°C, 65±5% R.H., photoperiod 14h L and 10h D).

Rearing procedure was carried out according to Ghoneim (1985) and improved by Bakr *et al.* (2010). Larvae were provided daily with fresh castor bean leaves *Ricinus communis*. The emerged adults were provided with cotton pieces soaked in 10% honey solution as a food source. Moths were allowed to lay eggs on *Oleander* branches. The egg patches were collected daily, and transferred into Petri dishes for another generation.

Precocene I administration

Precocene I (PI)(7-methoxy-2,2-dimethyl chromene) was kindly provided by Dr. Heba Hassan, Prof. at Plant Protection Research Institute, Agricultural Research Center, Doqqi, Giza, Egypt. Molecular formula: C₁₂H₁₄O₂ PI was diluted in acetone to prepare five doses: 150, 120, 90, 30 and 15 µg/larva. Each dose was topically applied (once) onto the thoracic sterna of 1-day old 5th (penultimate) and 1-day old (last) instar larvae by Hamilton microapplicator (NHN 737). Groups of 20 healthy larvae were used as replicates for each dose.

The newly moulted larvae did not be used, in the present study, because the first 24h, in each larval instar, is the critical period of corpora allata of *S. littoralis* (El-Ibrashy, 1971) after which the juvenile hormone is released in the haemolymph. Control larvae had been topically applied only with 1µl acetone. All treated and control larvae were kept individually under the previously mentioned laboratory controlled conditions.

All larvae were provided with fresh castor bean leaves every day, during the feeding period. Starting from the day after treatment all treated and control insects were observed daily to record all criteria of study.

Criteria of study

Mortality in larvae, pupae and adults were determined in %. LD₅₀ values were calculated by Microsoft office Excel, 2007, according to Finny (1971). Coefficient of growth: Coefficient of growth was calculated according to El-Ibrashy and Aref (1985) as follows: maximal body weight (mg)/ duration (in days) for each larva. Developmental duration: Dempster's equation (1957) was used for calculating

the developmental duration (in mean days \pm SD) and Richard's equation (1957) was used for calculating the developmental rate. Pupa rate: Pupa rate was expressed in % of the developed pupae. Adult emergence was determined in %. Precocious metamorphosis was determined in % of precocious pupation. Impaired morphogenesis was determined in % of deformed larvae, pupae and adults.

Statistical analysis of data

Data obtained were analyzed by the Student's *t*-distribution, and refined by Bessel correction (Moroney, 1956) for the test significance of difference between means.

Results

Lethal action of PI against *S. littoralis*

After topical application of PI (once) onto the 1-day old 5th (penultimate) instar larvae, insecticidal action

of PI was expressed in mortality among larvae, pupae and adults. According to data arranged in Table (1), PI exerted a toxic effect on the treated 5th instar larvae, in no certain trend.

The successfully moulted 6th instar larvae suffered the toxic effect only at higher two doses (14.3 and 11.1% mortality, at 150 and 120 μ g/larva, respectively, vs. 0% mortality of control congeners). Also, PI exhibited toxicity against pupae only at the higher two doses.

In a similar trend, the emerged adult females had been subjected to a toxic effect of PI only after larval treatment with the higher two doses. LD₅₀ value was calculated in 70.48 μ g/larva.

Table 1. Toxicity (%) of PI against *S. littoralis* topically treated as penultimate (5th) or last (6th) instar larvae.

Dose (μ g/larva)	Treatment of 1-day old 5 th instar larvae					Treatment of 1-day old 6 th instar larvae					
	Larval mortality		Pupal mortality	Adult mortality	Total mortality	LD ₅₀ (μ g/larva)	Larval mortality	Pupal mortality	Adult mortality	Total mortality	LD ₅₀ (μ g/larva)
	5 th instar	6 th instar									
150	53.3	14.3	16.67	20.00	73.33	70.48	20.00	8.33	9.09	33.33	234.96
120	40.0	11.1	12.50	28.57	60.00		13.33	53.85	16.67	66.67	
90	33.3	0.0	0.00	0.00	33.33		0.00	6.67	0.00	6.67	
30	60.0	0.0	0.00	0.00	60.00		0.00	0.00	0.00	0.00	
15	6.7	0.0	0.00	0.00	6.67		0.00	0.00	0.00	0.00	
Control	0.0	0.0	0.00	0.00	0.00		0.00	0.00	0.00	0.00	

The same table contains data of PI toxicity after topical application onto 1-day old 6th (last) instar larvae. Depending on these data, PI exhibited an insecticidal activity against larvae only at the higher two dose levels. Pupal mortality was observed after topical application of the higher three doses. With regard to the successfully emerged adult females, these moths were subjected to the toxic effect of PI only at the higher two doses. LD₅₀ value was calculated in 234.96 μ g/larva. As obviously seen, sensitivity of *S. littoralis* to PI toxicity was higher when the larvae were treated as 1-day old 5th instar.

Effect of PI on growth of *S. littoralis*

After topical application of PI doses onto the 1-day old 5th instar larvae, data of the maximal body weight

(max. wt), duration and coefficient of growth (CG) of the treated larvae and the successfully moulted 6th instar larvae were assorted in Table (2). In the light of these data, max.wt of treated 5th instar larvae significantly decreased at the higher two doses. In contrast, max. wt remarkably increased as a response to the action of PI, at the lower three dose levels. Apart from this diverse effect of PI on max.wt, it exerted a strong inhibitory action on larval growth, since the CG was considerably regressed, especially at the higher three dose levels. With no exception, PI enhanced the successfully moulted 6th instar larvae to attain increasing max. wt, in a dose-dependent course. In a similar trend, CG was pronouncedly induced in a dose-dependent manner (details table 2).

Table 2. Growth of *S. littoralis* after topical application of PI onto 1-day old penultimate instar larvae.

Dose ($\mu\text{g}/\text{larva}$)	Penultimate instar larvae			Last instar larvae			Precocious pupation (%)
	Maximal body weight (mean mg \pm SD)	Duration (mean days \pm SD)	Coefficient of growth (mean \pm SD)	Maximal body weight (mean mg \pm SD)	Duration (mean days \pm SD)	Coefficient of growth (mean \pm SD)	
150	102.8 \pm 17.5 b	1.0 \pm 0.0 b	102.8 \pm 17.5 c	622.5 \pm 32.7 b	4.0 \pm 0.7b	160.4 \pm 29.4 b	0
120	103.7 \pm 5.8 b	1.0 \pm 0.1 b	103.7 \pm 7.8 b	610.8 \pm 45.0c	4.1 \pm 0.6 b	148.9 \pm 11.11 c	0
90	138.3 \pm 8.7 b	1.4 \pm 0.1 a	98.8 \pm 6.8 c	605.2 \pm 55.8c	4.2 \pm 0.7a	148.0 \pm 39.3 b	0
30	151.20 \pm 17.6 b	1.5 \pm 0.2 a	108.0 \pm 21.4 a	572.8 \pm 67.0a	4.5 \pm 0.5a	127.9 \pm 13.4 a	0
15	162.0 \pm 10.8 c	1.5 \pm 0.3 a	108.0 \pm 19.5 a	560.7 \pm 77.9 a	4.5 \pm 0.5a	124.6 \pm 15.0 a	0
Control	119.0 \pm 38.9	1.9 \pm 0.3	110.4 \pm 19.8	513.4 \pm 1238	4.5 \pm 0.5	108.5 \pm 29.8	0

Mean \pm SD followed with the same letter a: insignificantly different ($P > 0.05$), b: significantly different ($P < 0.05$), c: highly significantly different ($P < 0.01$).

After topical application of PI onto the 1-day old last instar larvae, data of max. wt and CG were summarized in Table (3). On the basis of these data, a diverse action of PI on max.wt was recorded, since it significantly increased at the higher two doses but

drastically reduced at the lower three doses. Also, a diverse action of PI was exerted on the CG, since the compound promoted larvae to gain more growth at the higher two dose levels but slightly suppressed the CG at the lower three doses.

Table 3. Growth of *S. littoralis* after topical application of PI onto 1-day old last instar larvae.

Dose ($\mu\text{g}/\text{larva}$)	Maximal body weight (mean mg \pm SD)	Duration (mean days \pm SD)	Coefficient of growth (mean \pm SD)
150	572.7 \pm 151.5 b	5.63 \pm 0.8 a	111.7 \pm 49.9 b
120	682.0 \pm 149.1 b	6.40 \pm 1.5 a	114.8 \pm 57.7 b
90	468.1 \pm 18.8 b	5.71 \pm 0.8 a	085.7 \pm 35.8 a
30	472.2 \pm 78.1 b	5.33 \pm 0.9 b	092.0 \pm 25.3 a
15	493.8 \pm 47.4 a	5.18 \pm 0.7 b	089.3 \pm 13.5 a
Control	558.8 \pm 104.0	6.50 \pm 0.7	105.3 \pm 27.9

a, b: see footnote of Table (2).

Effects of PI on development, metamorphosis and morphogenesis of S. littoralis

Affected duration and developmental rate

After topical application of PI onto 5th instar larvae, data of affected larval duration were assorted in Table (2). In view of these data, duration of the treated larvae was conspicuously shortened, almost in a dose-dependent course. Also, duration of the successfully moulted 6th instar larvae was considerably shortened; especially at the higher three doses.

As obviously shown in Table (3), topical application of PI onto 6th instar larvae resulted in an abbreviation of larval duration, especially at the lower two doses. Data of distributed in Table (4) revealed a slight prolongation of pupal duration, after treatment of 5th instar larvae.

Also, a slight prolongation of pupal duration was easily observed in Table (5) after treatment of 6th instar larvae. Moreover, the pupal duration was remarkably prolonged at the lower two doses of PI.

As easily seen in tables 4 and 5, the developmental rate of pupae was generally regressed, in no certain trend, after treatment of 5th or 6th instar larvae with PI.

Impaired metamorphosis and morphogenesis

As obviously observed in Table (2), PI failed to induce the precocious pupation in *S. littoralis* after treatment of the penultimate instar larvae. With regard to the pupation rate, PI unexceptionally exerted a strong inhibitory action on pupation, regardless the larval instar under treatment (see tables 4 and 5).

Table 4. Development and metamorphosis of *S. littoralis* after topical application of PI onto 1-day old penultimate instar larvae.

Dose ($\mu\text{g}/\text{larva}$)	Pupation rate	Pupal Duration (mean days \pm SD)	Pupal Developmental Rate	Abnormal pupae (%)	Morphogenic efficiency (%)	Adult emergence (%)
150	45.0	7.65 \pm 2.9 a	13.00	0	0	50.0
120	48.5	7.65 \pm 3.1 a	13.00	0	0	58.3
90	62.5	7.85 \pm 2.8 a	12.80	0	0	66.6
30	63.5	7.50 \pm 3.3 a	13.33	0	0	100
15	69.5	7.50 \pm 3.1 a	13.33	0	0	100
Control	100	7.30 \pm 1.9	14.70	0	---	100

a: See footnote of Table (2).

After topical application of penultimate instar larvae, the inhibitory action of PI on pupation increased parallel to the dose level (Table 4). After treatment of last instar larvae, PI exhibited an inhibitory effect on the pupation, in a reverse trend with the dose level, i.e., the pupation rate was regressed consecutively with the decreasing dose level (Table 5).

In respect of the adult emergence, PI exerted a blocking action on this crucial metamorphosis event, after treatment of 5th instar larvae only with the higher three doses (Table 4). A similar arresting action of PI on adult emergence was recorded after treatment of 6th instar larvae only with the higher two doses (Table 5).

Table 5. Development and metamorphosis of *S. littoralis* after topical application of PI onto 1-day old last instar larvae.

Dose ($\mu\text{g}/\text{larva}$)	Pupation rate	Pupal Duration (mean days \pm SD)	Pupal Developmental Rate	Abnormal pupae (%)	Morphogenic efficiency (%)	Adult emergence (%)
150	86.9	7.53 \pm 1.1 a	13.28	50.0	60.0	85.7
120	86.7	7.50 \pm 3.5 a	13.33	06.7	07.1	95.5
90	83.8	7.39 \pm 1.4 a	13.53	00.0	00.0	100
30	83.3	8.00 \pm 0.8 c	11.95	00.0	00.0	100
15	80.5	8.44 \pm 0.7 b	11.85	00.0	00.0	100
Control	100	7.35 \pm 1.1	13.61	00.0	---	100

a, b, c: See footnote of Table (2).

After treatment of 5th instar larvae, PI failed to exhibit morphogenic efficiency on the present insect, since no malformed pupae or adults had been observed.

On the other hand, PI exhibited morphogenic efficiency on pupae, after treatment of 6th instar larvae only with the higher two doses, since morphologically abnormal pupae had been observed (50.0 and 6.7% abnormal pupae, at 150 and 120 $\mu\text{g}/\text{larva}$, respectively, vs. 0% abnormality in control pupae). As clearly shown in Fig. (1), some features of the impaired pupation program appeared in dwarf-sized pupae which failed to metamorphose into adults.

Discussion

Affected survival of S. littoralis by PI

Toxicity of several anti-juvenile hormone (anti-JH) compounds had been reported against different insect species. For examples, both PI and PII exhibited larvicidal activities against several mosquito species, such as *Aedes aegypti*, *Anopheles sacharovi* and *An. stephensi* (Saxena et al., 1994; Yasyukevich and Zvantsov, 1999). Precocenes exhibited larvicidal effects, in a dose-dependent course, on the Colorado potato beetle *Leptinotarsa decemlineata* (Farazmand and Chaika, 2008). A toxicological effect of PII was reported by Abdullah (2009) against larvae of the red palm weevil *Rynchophorus ferrugineus*.

Also, PII exhibited larvicidal and pupicidal effects on the grey flesh fly *Parasarcophaga dux* (Nassar *et al.*, 1999); larvicidal effect on the lepidopterous pest *Pericallia ricini* (Khan and Kumar, 2000); and larvicidal effect on the Asian tiger mosquito *Aedes albopictus* (Liu and Liu, 2014). After exposure of the newly moulted 2nd or 4th (penultimate) instar nymphs of the grasshopper *Euprepocnemis plorans* to some doses of PII, various mortality percentages were recorded among the treated nymphs of different instars and the emerged adults (Ghoneim and Basiouny, 2017). Apart from precocenes, other anti-JH compounds displayed different degrees of toxicity against some insects, such as synthesized EMD (ethyl (E)-3-methyl-2-dodecenoate) (Kuwano *et al.*, 1988) and some synthesized analogues of FMev (tetrahydro-4-fluoromethyl-4-hydroxy-2H-pyran-2-one) (Shuto *et al.*, 1988) against the mulberry silkworm *Bombyx mori*.



Fig.1. Pupal abnormalities of *S. littoralis* after treatment of 6th instar larvae with the higher two doses of PI (150 and 120 $\mu\text{g}/\text{larva}$). Normal pupa (at left) and dwarf pupa (at right). These dwarf-sized pupae failed to metamorphose into adults.

Results of the present study on the Egyptian cotton leafworm *Spodoptera littoralis* were in agreement with those previously reported results, since topical application (once) of PI onto 1-day old 5th (penultimate) or 6th (last) instar larvae affected the survival of the insect, as recorded in various mortality percentages among larvae, pupae and adult females.

However, the larval deaths may be attributed to the prevention of moulting larvae to swallow air volumes for splitting the old cuticle and expand the new one during ecdysis (Linton *et al.*, 1997). Also, these larval deaths may be due to the prevented feeding and continuous starvation of the insect (Ghoneim *et al.*, 2000). The pupal deaths can be directly related to the hormonal activity of the tested anti-JH compound or may be to other causes, such as bleeding, desiccation and suffocation, due to imperfect exuvation, failure of vital homeostatic mechanisms, *etc.* (Smaghe and Degheele, 1994). The adult mortalities can be explained by the retention and distribution of PI in the insect body as a result of direct and rapid transport *via* the haemolymph to other tissues, and/or by lower detoxification capacity of adults against the tested compound (Osman *et al.*, 1984).

The reported LD₅₀ (or LC₅₀) values of anti-JH compounds are variable in different insects. For examples, LD₅₀ of PII against the red cotton stainer *Dysdercus koenigii* has been found to be 85.46 and 82.37 mg/l^{-1} for 4th and 5th instar nymphs, respectively (Banerjee *et al.*, 2008). After treatment of 4th instar larvae of *A. albopictus* with PI and PII, LC₅₀ values were estimated in 41.63 $\mu\text{g}/\text{ml}$ and 43.55 $\mu\text{g}/\text{ml}$, respectively (Liu and Liu, 2014). LC₅₀ of PII against the booklice *Liposcelis bostrychophila* was calculated in 30.4 $\mu\text{g}/\text{cm}^2$ but LC₅₀ of PI was found as 64.0 $\mu\text{g}/\text{cm}^2$ (Lu *et al.*, 2014). LC₅₀ of PI against the cat flea *Ctenocephalides felis* was estimated in 10.97 ppm (Rust and Hemsarh, 2017). LC₅₀ values of the anti-JH agent Pitavastatin against the tobacco hornworm *Manduca sexta* and the viviparous cockroach *Diploptera punctata* were estimated in 5.23, and 395.2 μM , respectively (Li *et al.*, 2017). LD₅₀ values of PII against *E. plorans* were 0.388 and 17.022 $\mu\text{g}/\text{cm}^2$ after topical treatment of newly moulted 2nd and 4th (penultimate) instar nymphs, respectively (Ghoneim and Basiouny, 2017). In the current study, LD₅₀ values were estimated in 70.48 and 234.96 $\mu\text{g}/\text{larva}$, after treatment of 5th and 6th instar larvae, respectively. Thus, the 5th instar larvae were more sensitive to PI than 6th instar larvae.

As reported in the currently available literature, LD₅₀ (or LC₅₀) value depends on several factors, such as susceptibility of the insect and its treated stage or instar, lethal potency of the tested compound and its concentration levels, method and time of treatment, as well as the experimental conditions.

Growth inhibition in S. littoralis by PI

After treatment of 5th or 6th instar larvae of *S. littoralis* with PI doses, in the current investigation, the compound exerted a diverse action on the maximal body weight (max. wt) of treated larvae, depending on the dose. The coefficient of growth was considerably regressed, irrespective of the instar of treated larvae. These results were, to some extent, in accordance with some of the reported results of inhibited growth of various insects by different anti-JH compounds. Growth of the last instar larvae of the mealworm beetle *Tenebrio molitor* was remarkably inhibited after treatment with several chromene derivatives (Roberto *et al.*, 1998). PI and PII exhibited growth-inhibiting activities against the mosquito species *A. aegypti*, *An. sacharovi* and *An. stephensi* (Saxena *et al.*, 1994; Yasyukevich and Zvantsov, 1999). After feeding of *M. sexta* larvae on a treated diet with HMG-CoA reductase inhibitors, Fluvastatin, Lovastatin or Pitavastatin, the growth rate of treated larvae was significantly slow (Li *et al.*, 2017). After exposure of the newly moulted 2nd or 4th (penultimate) instar nymphs of *E. plorans* to different doses of PII, the nymphal growth of both 4th and 5th instars had been slightly inhibited after treatment of 2nd instar nymphs, but remarkably regressed after treatment of 4th instar nymphs (Ghoneim and Basiouny, 2017). To understand the inhibition in growth of *S. littoralis*, in the current study, PI might affect the tissues and cells undergoing mitosis (Nasiruddin and Mordue, 1994). Also, PI might exhibit an inhibitory effect on the protein contents in haemolymph and fat body (Lange *et al.*, 1983).

Developmental disturbance in S. littoralis by PI

The developmental rate of an insect stage is usually reversely related to the developmental duration, i.e. shorter duration indicates faster rate and vice versa.

The larval duration in several insect species (holometabolous or hemimetabolous) had been prolonged as a response to the action of different anti-JH compounds. For examples, Bowers and Aldrich (1980) recorded a prolongation of 5th nymphal instar in the milkweed bug *Oncopeltus fasciatus* after treatment with PI. Treatment of the 4th instar nymphs of the desert locust *Schistocerca gregaria* with PII resulted in prolongation of the duration of both 4th and 5th nymphal instars (Eid *et al.*, 1982). Treatment of 6th instar larvae of the lawn armyworm *Spodoptera mauritia* with PII resulted in prolongation of duration in last larval instar (Mathai and Nair, 1983, 1984). The nymphal period of the grasshopper *Aiolopus thalassinus* was prolonged after topical application of PIII onto 5th instar nymphs (Osman, 1988). Treatment of the tobacco cutworm *Spodoptera litura* larvae with PI, PII or ethoxyprococene (a synthetic analog of P II) resulted in prolongation of larval period (Srivastava and Kumar, 1997, 1999). After treatment of 4th instar nymphs of *D. koenigii* with PII, duration of the successfully moulted 5th instar nymphs was prolonged (Banerjee *et al.*, 2008). Apart from precocenes, prolongation of the larval period in the fall webworm *Hyphantria cunea* was recorded after treatment with FMev (Farag and Varjas, 1983). Similar results of prolonged larval duration were reported in *B. mori* by KK-22 (phenylimidazoles) (Kuвано and Eto, 1983; Asano *et al.*, 1984). After treatment of 4th instar larvae of *B. mori* with the synthesized 3-(2-methyl-1-phenyl-1-propenyl)pyridine, the larval period was prolonged (Yoshida *et al.*, 2000). Results of the present investigation disagreed with those previously reported results, since topical application of PI onto 5th instar larvae of *S. littoralis* resulted in conspicuously shortened larval duration, almost in a dose-dependent course. Also, the larval duration was remarkably shortened after treatment of 6th instar larvae with PI, especially at the lower two doses.

On the other hand, our results were in accordance with those reported results of shortened larval duration after treatment with some anti-JH

compounds, such as *P. dux* after treatment of the 3rd instar larvae with PII (Nassar *et al.*, 1999); the flesh fly *Sarcophaga ruficornis* after treatment of the last instar larvae with PI, PII or PIII (Srivastava and Kumar, 1996); the house fly *Musca domestica* after treatment of the larvae with PII (Gaur and Kumar, 2009); *B. mori* after treatment of the 3rd and 4th instars with the imidazole compound SSP-11 (Kiuchi *et al.*, 1985). Moreover, topical application of PI onto 5th or 6th instars larvae of *S. littoralis*, in the present study, resulted in a slight or significant prolongation of the developed pupae, depending on the dose level. In general, the developmental rate of these pupae was slightly regressed. As far as our literature survey could ascertain, no information was available for the effects of anti-JH compounds on the pupal duration or developmental rate.

To explicate the shortened larval duration of *S. littoralis*, in the current investigation, it may be due to the response of these treated larvae for avoiding the adverse action of PI, as a xenobiotic agent. On the other hand, the prolongation of pupal period in *S. littoralis*, in the present study, indicated a retarding action of PI on the development as expressed in regression of the developmental rate. This prolongation in the pupal duration may be attributed to the indirect interference of this compound with the neuroendocrine organs responsible for the synthesis and release of tropic hormones, like prothoracicotropic hormone (Subrahmanyam *et al.*, 1989). Also, the recorded prolongation of pupal stage may be attributed to a disturbing action of PI on the persistence of JH in the haemolymph where it is only in the absence of JH that ecdysone could be activated and lead to the formation of the next stage (Bowers, 1982; Kuwano *et al.*, 1988). In addition, PI may exhibit a delaying effect on the pupal transformation into adults (Linton *et al.*, 1997). In particular, the final step of chitin biosynthesis pathway was inhibited by PI and the precursor was not converted into chitin leading to a prolongation of developmental duration (Djehader *et al.*, 2014).

Impaired metamorphosis and morphogenesis of S. littoralis by PI

Precocious metamorphosis

In the current study, PI failed to induce the precocious pupation or other features of precocious metamorphosis of *S. littoralis*. This result disagreed with many reported results of precocious pupation or metamorphosis in several insect species as a response to different anti-JH compounds, such as the precocious pupation in *H. cunea* (Farag and Varjas, 1983) and *S. mauritia* (Balamani and Nair, 1989) after treatment with the anti-JH compound Fluoromevalonate (FMev), the flesh fly *Neobellieria bullata* after treatment with PII (Darvas *et al.*, 1990), *P. ricini* after treatment with PII (Khan and Kumar, 2000) and *B. mori* after treatment of 4th instar larvae with a synthesized 3-pyridine derivative (Yoshida *et al.*, 2000). Exposure of 2nd instar nymphs of the grasshopper *Euprepocnemis plorans* to PII led to precociously moulting into 4th instar, skipping off 3rd instar (only at the lowest dose). Also, exposure of 4th instar nymphs to PII, some treated nymphs precociously metamorphosed into adultoids, omitting the 5th instar (Ghoneim and Basiouny, 2017).

However, the failure of PI to induce precocious pupation or other features of precocious metamorphosis, in the present study on *S. littoralis*, was in agreement with those reported results of failure of some anti-JH compounds to induce precocious forms, such as PI against the sunn pest *Eurygaster integriceps* (Tarrant *et al.*, 1982), PII against *S. mauritia* (Mathai and Nair, 1983), the synthesized anti-JH compound EMD or its analogues against *B. mori* (Kuwano *et al.*, 1988), the synthesized compound (S)-(+)-FMev against *B. mori* (Shuto *et al.*, 1988) and PII against *S. gregaria* (Islam, 1995).

The failure of PI to induce precocious pupation of *S. littoralis*, in the current study, can be understood because it did not exhibit an anti-JH activity indicating the inability to affect the larval JH levels (Edwards *et al.*, 1983). Some authors (Burt *et al.*, 1978; Haunerland and Bowers, 1985; Staal, 1986) reported that the larvae of holometabolous insects-

with few exceptions- are less susceptible to the action of Precocenes than hemimetabolous insects. In contrast to our finding, some holometabolous insects, such as *S. mauritia* and *S. littoralis* have been reported to be sensitive to the anti-JH activity of Precocenes (Mathai and Nair, 1984; Khafagi and Hegazi, 2001).

Pupation process In the current investigation, PI exerted a strong inhibitory action on the pupation rate in *S. littoralis*, irrespective of the larval instar under treatment, 5th or 6th instar. No information is available in the current literature for the inhibitory effects of anti-JH compounds on the pupation rate in insects. However, the present result was consistent with those reported inhibitory effects of some juvenile hormone analogues (JHAs) or other insect growth regulators (IGRs) on the pupation rate in various insects, such as the diamondback moth *Plutella xylostella* by Hexaflumuron (Mahmoudvand *et al.*, 2012); *S. littoralis* by Novaluron (Ghoneim *et al.*, 2015); the lesser mulberry pyralid *Glyphodes pyloalis* by Lufenuron (Aliabadi *et al.*, 2016); the pink boll worm *Pectinophora gossypiella* (Ghoneim *et al.*, 2017a) and the olive leaf moth *Palpita unionalis* (Ghoneim *et al.*, 2017b) by Novaluron.

Deranged morphogenesis

Precocenes have been reported to be anti-juvenile hormone, accelerating the development of insects and inducing dwarfness associated with low survival rates (Ghosh *et al.*, 2012). In the present study on *S. littoralis*, no malformed pupae or adults had been observed after treatment of 5th instar larvae with PI. On the other hand, some morphologically abnormal pupae were produced after treatment of 6th instar larvae with the higher two doses. These dwarf-sized pupae failed to metamorphose into adults. This result was, to a great extent, in agreement with those reported results of adverse action of some anti-JH compounds on the morphogenesis of a number of insects. For examples, Treatment of 3rd instar larvae of *B. mori* with some terpenoid imidazole compounds led to the formation of miniature pupae after molting to 4th instar larvae (Kuwano *et al.*, 1983).

Topical application of PIII onto eggs or 5th instar nymphs of *A. thalassinus* resulted in some prothetelic morphogenic disturbances (Osman, 1988).

Production of abnormal puparia was recorded in *S. ruficornis* after administration of PI, PII or PIII to the last instar larvae (Srivastava and Kumar, 1996). Treatment of *S. litura* larvae with PI, PII or ethoxyprecocene (a synthetic analog of P II) resulted in the production of abnormally formed pupae (Srivastava and Kumar, 1999). Puparial malformations were observed in *P. dux* after topical application of PII onto the 3rd instar larvae (Nassar *et al.*, 1999). Treatment of *M. domestica* maggots with PIII led to the formation of abnormal puparia (Gaur and Kumar, 2009). Larval treatment of *E. integriceps* with PI led to the production of some morphological abnormalities (Amiri *et al.*, 2010).

For interpretation of the pupation rate regression and impairment of the pupation program in *S. littoralis*, as caused by PI in the present study, PI might exert a prohibiting action on the prothoracic gland (ecdysone-producing gland) and hence the ecdysone could not be synthesized and/or released. It is well known that the absence of ecdysone leads to failure of ecdysis. Whatever the mode of action, PI suppressed the chitin synthesis and prevented the normal deposition of new cuticle during apolysis leading to the production of pupal deformities (Retnakaran *et al.*, 1985). In addition, PI might block the release of morphogenic peptides, causing alteration in both ecdysteroid and juvenoid titers (Barnby and Klocke, 1990). A suggestion of Gaur and Kumar (2010) may be appreciated. PI might disrupt the ecdysteroid metabolism or might alternatively act directly to inhibit the release of ecdysis-triggering hormone.

Inhibited adult emergence

Scarce studies have examined the effects of anti-JH compounds on the adult emergence in insects. Inhibition of adult emergence in *S. ruficornis* was recorded after larval treatment with PII (Khan and Kumar, 2005). KK-42 (a terpenoid imidazole) was reported to inhibit the adult emergence of *B. mori* when applied to the newly formed pupae (Kadono-Okuda *et al.*, 1987).

In the present study on *S. littoralis*, the adult emergence was slightly or drastically blocked, especially at the higher three or two doses of PI, depending on the larval instar under treatment, 5th or 6th instar.

For interpretation of this result, it is important to point out that the adult emergence in insects is a crucial physiological process and regulated by the eclosion hormone. The disturbance of this hormone appears to partially or completely arrest the adults to emerge. The present result of blocked adult emergence can be interpreted by the disruptive effect of PI on the normal metabolism of insect hormones during the development of the immature stages leading to failure of adult emergence (Trigo *et al.*, 1988). In particular, PI might disturb the adult eclosion hormone release and/or inhibition of the neurosecretion (Al-Sharook *et al.*, 1991; Joseph rajkumar *et al.*, 1999). On the molecular basis, anti-JH compounds, like PI, might cause misexpression of certain genes, particularly the brood complex (*br-C*) transcription factor gene, leading to symptoms of deranged metamorphosis, like blocking of adult emergence (Wilson, 2004; Nandi and Chakravarty, 2011).

In general, it is still unclear whether the effects of precocenes on insects are restricted to the corpora allata (JH-producing organs) (Garcera *et al.*, 1991; Burks *et al.*, 1992), or are more general due to their toxicity (Ergen, 2001), or indirect as consequence of the decreased JH titers (Amsalem *et al.*, 2014b), since many studies revealed multiple targets for the effects of Precocenes in insects (Chen *et al.*, 2005 a,b; Amiri *et al.*, 2010).

Conclusion

According to the obtained results of the present study, it can be concluded that PI exhibited various degrees of toxicity against all developmental stages of *S. littoralis*, as well as some disruptive effects on growth and development but failed to induce anti-JH activity against this pest. Therefore, PI may be recommended to use in the integrated pest management program after further investigation on its activity under the field conditions.

Acknowledgement

The authors greatly appreciate Dr. Heba Hassan, Prof. at Plant Protection Research Institute, Agricultural Research Center, Doqqi, Giza, Egypt, who kindly provided a sample of susceptible strain of *S. littoralis* and PI.

References

Abdullah MAR. 2009. Identification of the biological active compounds of two natural extracts for the control of the red palm weevil, *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae). Egyptian Academic Journal of Biological Sciences **2(2)**, 35-44.

Adham FK, Rashad EM, Shoukry IF, Nasr EE. 2009. Host plants shifting affects the biology and biochemistry of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). Journal of Biological Sciences **2(1)**, 63-71.

Aliabadi FP, Sahragard A, Ghadamyari M. 2016. Lethal and sublethal effects of a chitin synthesis inhibitor, lufenuron, against *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae). Journal of Crop Protection **5(2)**, 203-214.
<http://dx.doi.org/10.18869/modares.jcp.5.2.203>

Al-Sharook Z, Balan K, Jiang Y, Rembold H. 1991. Insect growth inhibitors from two tropical Meliaceae: Effects of crude seed extracts on mosquito larvae. Journal of Applied Entomology **111**, 425-430.
<https://doi.org/10.1111/j.1439-0418.1991.tb00344.x>

Amiri A, Bandani AR, Ravan S. 2010. Effect of an anti-juvenile hormone agent (Precocene I) on sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae) development and reproduction. African Journal of Biotechnology **9(36)**, 5859-5868.

Amsalem E, Malka O, Grozinger C, Hefetz A. 2014a. Exploring the role of juvenile hormone and vitellogenin in reproduction and social behavior in bumble bees. BMC Evolutionary Biology **14**, 1-13.
<http://dx.doi.org/10.1186/1471-2148-14-45>

- Amsalem E, Teal P, Grozinger CM, Hefetz A.** 2014b. Precocene-I inhibits juvenile hormone biosynthesis, ovarian activation, aggression and alters sterility signal production in bumble bee (*Bombus terrestris*) workers. The Journal of Experimental Biology **217**, 3178-3185.
<http://dx.doi.org/10.1242/jeb.107250>
- Asano S, Kuwano E, Eto M.** 1984. Anti-juvenile hormone activity of 1-citronellyl-5-phenylimidazole in the 3rd instar silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae). Applied Entomology and Zoology **19(2)**, 212-220.
<https://doi.org/10.1303/aez.19.212>
- Attathom T.** 2002. Biotechnology for insect pest control. Proceedings of a Satellite Forum: "Sustainable Agricultural System in Asia". Nagoya, Japan, 73-84 p.
- Aydin MH, Gurkan MO.** 2006. The efficacy of spinosad on different strains of *S. littoralis* (Boisduval) (Lepidoptera: Noctuidae). Turkish Journal of Biology **30**, 5-9.
- Bakr RFA, El-barky NM, Abd Elaziz MF, Awad MH, Abd El-Halim HME.** 2010. Effect of Chitin synthesis inhibitors (flufenoxuron) on some biological and biochemical aspects of the cotton leafworm *Spodoptera littoralis* Bosid. (Lepidoptera: Noctuidae). Egyptian Academic Journal Biological Sciences **2(2)**, 43-56.
- Balamani E, Nair VSK.** 1989. Effects of anti-juvenile hormone agents and a juvenile hormone analog on neck-ligated post-feeding last instar larvae of *Spodoptera mauritia* Boisid. (Lepidoptera: Noctuidae). International Journal of Tropical Insect Science **10(5)**, 583-589.
<https://doi.org/10.1017/S1742758400021706>
- Banerjee S, Kalena GP, Banerji A, Singh AP.** 2008. New synthetic precocenoids as potential insect control agents. Journal of Environmental Biology **29(6)**, 951- 957.
- Banerji A, Goomer NC.** 1984. Synthesis of naturally occurring 6-oxygentaed-2, 2-dimethyl chromenes. Indian Journal of Chemistry **23B**, 885-886.
- Banerji A, Kalena GP.** 1989. Heterocycles from diacetyl phenols: Synthesis of benzodipyran, acetyl chromenes of pyrano-(1,2) benzisoxazoles. Heterocycle **28**, 711-718.
- Barnby MA, Klocke JA.** 1990. Effects of azadirachtin on levels of ecdysteroids and prothoracicotropic hormone-like activity in *Heliothis virescens* (Fabr.) larvae. Journal of Insect Physiology **36**, 125-131.
[https://doi.org/10.1016/0022-1910\(90\)90183-G](https://doi.org/10.1016/0022-1910(90)90183-G)
- Bede JC, Teal PE, Goodman WG, Tobe SS.** 2001. Biosynthetic pathway of insect juvenile hormone III in cell suspension cultures of the sedge *Cyperus iria*. Plant Physiology **127(2)**, 584-593.
<http://dx.doi.org/10.1104/pp.010264>.
- Bowers WS.** 1976. Discovery of insect anti-allatotropins. In: "The Juvenile Hormones" (Gilbert LI. Ed.). New York, NY: Plenum Press, 394-408 p.
- Bowers WS.** 1982. Toxicology of the precocenes. In: "Insecticide Mode of Action"(Coats JR. ed.). New York, Academic Press, 403-427 p.
- Bowers WS, Aldrich JR.** 1980. In vivo inactivation of denervated corpora allata by precocene II in the bug *Oncopeltus fasciatus*. Experientia **36**, 362-364.
<http://dx.doi.org/10.1007/BF01952328>
- Burks CS, Evans LD, Kim Y, Krafur ES.** 1992. Effects of Precocene and methoprene application in young-adult *Musca autumnalis*. Physiological Entomology **17**, 115-120.
<https://doi.org/10.1111/j.1365-3032.1992.tb01187.x>
- Burt ME, Kuhr RJ, Bowers WS.** 1978. Metabolism of precocene II in the cabbage looper and European corn-borer. Pesticide Biochemistry and Physiology **9**, 300-303.
[https://doi.org/10.1016/0048-3575\(78\)90010-X](https://doi.org/10.1016/0048-3575(78)90010-X)

- Casida JE, Quistad GB.** 1998. Golden age of insecticide research: past, present, or future. Annual Review of Entomology **43**, 1-16.
<http://dx.doi.org/10.1146/annurev.ento.43.1.1>
- Chen YR, Chen SC, Chang HW, Sun G, Kou R.** 2005a. Effect of exogenous juvenile hormone III and precocene II on agonistic behavior and the *Corpora allata in vitro* activity in the male lobster cockroach *Nauphoeta cinerea* (Dictyoptera: Blaberidae, Oxyhaloinae). Zoological Studies **44**, 409-416.
- Chen Z, Madden RD, Dillwith JW.** 2005b. Effect of precocene II on fatty acid metabolism in the pea aphid, *Acyrtosiphon pisum*, under cold stress. Journal of Insect Physiology **51(4)**, 411-416.
- Chenevert R, Perron JM, Paquin RL, Robitaille M, Wang YK.** 1980. Activity of precocene analogues on *Locusta migratoria* (Forsk.). Experientia **36**, 379-380.
- Chowański S, Kudlewska M, Marciniak P, Rosiński G.** 2014. Synthetic Insecticides- is there an Alternative? Polish Journal Environmental Studies **23(2)**, 291-302.
- Costa LG, Giordano G, Guizzetti M, Vitalone A.** 2008. Neurotoxicity of pesticides: A brief review. Frontiers in BioScience **13**, 1240-1249.
- Damalas CA, Eleftherohorinos IG.** 2011. Pesticide exposure, safety issues, and risk assessment indicators. International Journal of Environmental Research and Public Health **8(5)**, 1402-1419.
- Darvas B, Kuwano E, Eto M, Tag al-din MH, Timar T.** 1990. Effects of some anti-juvenile hormone agents (Precocenes-2, J-2710, KK-110) on post embryonic development of *Neobellieria bullata*. Agricultural and Biological Chemistry **54(11)**, 3045-3047.
<https://doi.org/10.1080/00021369.1990.10870450>
- Davies TGE, Field LM, Usherwood PNR, Williamson MS.** 2007. DDT, pyrethrins and insect sodium channels. International Union of Biochemistry and Molecular Biology Life **59**, 151-162.
DOI: 10.1080/15216540701352042
- Dempster C.** 1957. The population dynamic of *Moroccan locust, Dociostarus maroccanus* in Cyprus. Anti Locust Bull., 27 p.
- Denlinger DL, Yocum GD, Rinehart JP.** 2012. Hormonal control of diapause. In: "Insect Endocrinology" (Lawrence IG. ed.). Academic Press, San Diego, CA, USA.430-463 p.
- Djeghader NEH, Aïssaoui L, Amira K, Boudjelida H.** 2014. Impact of a chitin synthesis inhibitor, Novaluron, on the development and the reproductive performance of mosquito *Culex pipiens*. World Applied Sciences Journal **29(7)**, 954-960.
<http://dx.doi.org/10.5829/idosi.wasj.2014.29.07.82190>
- Edwards JP, Bergot BJ, Staal GB.** 1983. Effects of three compounds with anti-juvenile hormone activity and a juvenile hormone analogue on endogenous juvenile hormone levels in the tobacco horn worm, *Manduca sexta*. Journal of Insect Physiology **29(1)**, 83-89.
[https://doi.org/10.1016/0022-1910\(83\)90109-9](https://doi.org/10.1016/0022-1910(83)90109-9)
- Eid MA, Salem MS, El-Ibrashy MT, Abdel-Hamid M.** 1982. Effects of precocene II, Cycloheximide and C16-JH on the growth rate and adult morphometrics of *Schistocerca gregaria* Forsk. Bulletin of Entomological Society of Egypt (Economic Series) **13**, 95-105.
- El-Ibrashy MT.** 1971. The effect of allatectomy upon reproduction and development of the cotton leaf-worm *Spodoptera littoralis*. Journal of Insect Physiology **17**, 1783-1790.
[https://doi.org/10.1016/0022-1910\(71\)90075-8](https://doi.org/10.1016/0022-1910(71)90075-8)
- El-Ibrashy MT.** 1982. Juvenile hormone mimics in retrospect and antagonists in prospect. Journal of Applied Entomology **94**, 217-236.
<https://doi.org/10.1111/j.1439-0418.1982.tb02569.x>

- El-Ibrashy MT, Aref NB.** 1985. Effects of certain juvenoids on growth and morphogenesis in *Spodoptera littoralis* Boisduval. Journal of plant protection in the tropics **2(2)**, 105-116.
- El-Khawas MAM, Abd El-Gawad HAS.** 2002. The efficiency of two plant extracts (Fenugreek and Lupine) and commercial biofungicide (Biofly) on the cotton leafworm, *Spodoptera littoralis* (Boisd) (Lepidoptera: Noctuidae) larvae as a new approach of control. Journal of Egyptian German Society of Zoology **37**, 39-57.
- Farag AI, Varjas L.** 1983. Precocious metamorphosis and moulting deficiencies induced by an anti-JH compound FMev in the fall webworm, *Hyphantria cunea*. Entomologia Experimentalis et Applicata **34**, 65-70.
<https://doi.org/10.1111/j.1570-7458.1983.tb03291.x>
- Farazmand H, Chaika EY.** 2008. Effects of Precocene-I and II on the development of Colorado potato beetle, *Leptinotarsa decemlineata* (Col.: Chrysomelidae). Applied Entomology and Phytopathology **76(1)**, 27-43.
- Finney DJ.** 1971. Probit analysis. 3rd ed. Cambridge, England: Cambridge University Press, 318 p.
- Flatt T, Heyland A, Rus F, Porpiglia E, Sherlock C, Yamamoto R, Garbuzov A, Palli SR, Tatar M, Silverman N.** 2008. Hormonal regulation of the humoral innate immune response in *Drosophila melanogaster*. Journal of Experimental Biology **211**, 2712-2724.
<http://dx.doi.org/10.1242/jeb.014878>
- Gäde G, Goldsworthy GJ.** 2003. Insect peptide hormones: a selective review of their physiology and potential application for pest control. Pest Management Science **59(10)**, 1063-1075.
<http://dx.doi.org/10.1002/ps.755>
- Garcera MD, Ibanez P, Martinez R, Cunat P.** 1991. Variations induced by juvenile hormone and precocene II in the metabolic rate of 3 species of insects. Revista Espanola de Fisiologia. **47**, 31-36.
- Gaur R, Kumar K.** 2009. Effect of precocene on morphogenesis in housefly, *Musca domestica* Linn. Entomol **34**, 107-110.
- Gaur R, Kumar K.** 2010. Insect growth-regulating effects of *Withania somnifera* in a polyphagous pest, *Spodoptera litura*. Phytoparasitica **38(3)**, 237-241.
<http://dx.doi.org/10.1007/s12600-010-0092-x>
- Ghoneim KS.** 1985. Physiological studies on endocrine and reproductive systems of the cotton leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). Ph.D. Thesis, Fac. of Sci., Al-Azhar Univ., Cairo, Egypt.
- Ghoneim K, Basiouny A.** 2017. Insecticidal and antijuvenile hormone activities of Precocene II against the grasshopper *Euprepocnemis plorans* (Charp.) (Orthoptera: Acrididae). International Journal of Trend in Scientific Research and Development **1(6)**, 510-524.
- Ghoneim KS, Mohamed HA, Bream AS.** 2000. Efficacy of the neem seed extract, Neemazal, on growth and development of the Egyptian cotton leafworm, *Spodoptera littoralis* Boisduval. (Lepidoptera: Noctuidae). Journal of Egyptian German Society of Zoology **33**, 161-179.
- Ghoneim K, Tanani M, Hamadah Kh, Basiouny A, Waheeb H.** 2015. Bioefficacy of Novaluron, a chitin synthesis inhibitor, on survival and development of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). Journal of Advances in Zoology **1(1)**, 24-35.
- Ghoneim K, Hassan HA, Tanani MA, Bakr NA.** 2017a. Toxic and disruptive effects of Novaluron, a chitin synthesis inhibitor, on development of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). International Journal of Entomology Research **2(2)**, 36-47.
- Ghoneim K, Hamadah Kh, Mansour AN, Abo Elsoud AA.** 2017b. Toxicity and disruptive impacts of Novaluron, a chitin synthesis inhibitor, on development and metamorphosis of the olive leaf moth *Palpita unionalis* (Hübner) (Lepidoptera: Pyralidae). International Journal of Trend in Research and Development **4(3)**, 184-193.

- Ghosh A, Chowdhury N, Chandra G.** 2012. Plant extracts as potential mosquito larvicides. *The Indian Journal of Medical Research* **135**, 581-598.
- Gilbert LI, Granger NA, Roe RM.** 2000. The juvenile hormones: historical facts and speculations on future research directions. *Insect Biochemistry and Molecular Biology* **30**, 617-644.
[http://dx.doi.org/10.1016/S0965-1748\(00\)00034-5](http://dx.doi.org/10.1016/S0965-1748(00)00034-5)
- Gillott C.** 2005. Insects and humans. In: "Entomology" (Gillott C. Ed.). Third Edition. Springer, Dordrecht, 725-782 p.
- Hauerland NH, Bowers WS.** 1985. Comparative studies on pharmacokinetics and metabolism of the anti-juvenile hormone precocene II. *Archives of Insect Biochemistry and Physiology* **2**, 55-63.
<https://doi.org/10.1002/arch.940020106>
- Hosny MM, Topper CP, Moawad GG, El-Saadany GB.** 1986. Economic damage thresholds of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) on cotton in Egypt. *Crop Protection* **5**, 100-104.
[https://doi.org/10.1016/0261-2194\(86\)90088-8](https://doi.org/10.1016/0261-2194(86)90088-8)
- Hussain A.** 2012. The effect of non-host plant volatiles on the reproductive behaviour of the Egyptian cotton leafworm, *Spodoptera littoralis*. M.Sc. Thesis, Swedish Univ. Agric. Sci. 140 p.
- Ishaaya I, Yablonski S, Horowitz AR.** 1995. Comparative toxicity of two ecdysteroid agonists, RH-2485 and RH-5992, on susceptible and pyrethroid-resistant strains of the Egyptian cotton leafworm, *Spodoptera littoralis*. *Phytoparasitica* **23**, 139-145.
<http://dx.doi.org/10.1007/BF02980973>
- Islam MS.** 1995. Endocrine manipulation in crowd-reared desert locust *Schistocerca gregaria* (Forsk.) II. Effects of Precocene on mortality and reproductive parameters. *Bangladesh Journal of Zoology* **23**, 199-208.
- Isman MB, Yan JY, Proksch P.** 1986. Toxicity of natural chromene derivatives to a grasshopper. *Naturwissenschaften* **73**, 500-501.
<http://dx.doi.org/10.1007/BF00367198>
- Josephraj Kumar A, Subrahmanyam B, Srinivasan S.** 1999. Plumbagin and azadirachtin deplete haemolymph ecdysteroid levels and alter the activity profiles of two lysosomal enzymes in the fat body of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *European Journal of Entomology* **96**, 347-353.
- Kadono-Okuda K, Kuwano E, Eto M, Yamashita O.** 1987. Inhibitory action of an imidazole compound on ecdysone synthesis in prothoracic glands of the silkworm, *Bombyx mori*. *Development, Growth and Differentiation* **29**, 527-533.
<https://doi.org/10.1111/j.1440-169X.1987.00527.x>
- Khafagi WE, Hegazi EM.** 2001. Effects of juvenile hormones I & II and precocenes I & II on the immune response of *Spodoptera littoralis* larvae to supernumerary larvae of the solitary parasitoid, *Microplitis rufiventris* Kok. *Journal of Insect Physiology* **47**, 1249-1259.
[http://dx.doi.org/10.1016/S0022-1910\(01\)00110-X](http://dx.doi.org/10.1016/S0022-1910(01)00110-X)
- Khan IA, Kumar K.** 2000. Precocene II acts as insect growth regulator in polyphagous pest, *Pericallia ricini* F. (Lepidoptera: Arctiidae). The Proceedings of the National Academy of Sciences, India **70(B)**, 279-285.
- Khan IA, Kumar K.** 2005. Effect of precocene II on imaginal differentiation in flesh fly, *Sarcophaga ruficornis* F. (Diptera: Sarcophagidae). *Entomon* **30**, 187-191.
- Kiss I, Fodor A, Timar T, Hosztafi S, Sebok P, Torok T, Viragh E, Berenyi M.** 1988. Biological activity of precocene analogues on *Locusta migratoria*. *Experientia* **44**, 790-792.
- Kiuchi M, Kimura K, Akai H.** 1985. Induction of trimolters from a tetramolter strain of *Bombyx mori* by anti-juvenoid treatment. *Journal of Sericultural Science of Japan* **54(1)**, 77-81.
<https://doi.org/10.11416/kontyushigen1930.54.77>

- Kozhanova NI, Nemeč V.** 1991. Comparison of the effects of hydroprene and precocene II on oogenesis in the mealworm, *Tenebrio molitor* (Coleoptera, Tenebrionidae). Acta entomologica Bohemoslovaca. **88**, 281-285.
- Kuwano E, Eto M.** 1983. Terpenoid imidazoles: New anti-juvenile hormones and insecticidal compounds. Proc. 10th Int. Congr. Plant Prot., Brighton 1, 278.
- Kuwano E, Takeya T, Eto M.** 1983. Terpenoid imidazoles: new antijuvenile hormones. Agricultural and Biological Chemistry **47**, 921-923.
<https://doi.org/10.1271/bbb1961.47.921>
- Kuwano E, Tanaka Y, Kikuchi M, Eto M.** 1988. Effects of anti-juvenile hormones and related compounds on development in the larvae of *Bombyx mori*. Journal of Faculty of Agriculture, Kyushu Univ. **33(1, 2)**, 17-28.
- Lange AB, Phillips DR, Loughton BG.** 1983. The effects of precocene II on early adult development in male *Locusta*. J. Insect Physiology **29(1)**, 73-81.
[https://doi.org/10.1016/0022-1910\(83\)90108-7](https://doi.org/10.1016/0022-1910(83)90108-7)
- Li Y-M, Kai Z-P, Huang J, Tobe SS.** 2017. Lepidopteran HMG-CoA reductase is a potential selective target for pest control. Peer J., 5, e2881.
<https://doi.org/10.7717/peerj.2881>
- Linton YM, Nisbet AJ, Mordue (Luntz) AJ.** 1997. The effect of azadirachtin on the testes of the desert locust *Schistocerca gregaria* (Forsk.). Journal of Insect Physiology **43**, 1077-1084.
[https://doi.org/10.1016/S0022-1910\(97\)00060-7](https://doi.org/10.1016/S0022-1910(97)00060-7)
- Liu XC, Liu ZL.** 2014. Evaluation of larvicidal activity of the essential oil of *Ageratum conyzoides* L. aerial parts and its major constituents against *Aedes albopictus*. Journal of Entomology and Zoology Studies. **2(4)**, 345-350.
- Lu XN, Liu XC, Liu QZ, Liu ZL.** 2014. Isolation of insecticidal constituents from the essential oil of *Ageratum houstonianum* Mill. against *Liposcelis bostrychophila*. Journal of Chemistry, Article ID 645687, 6 p.
<http://dx.doi.org/10.1155/2014/645687>
- Magd El-Din S, El-Gengaihi SE.** 2000. Joint action of some botanical extracts against the Egyptian cotton leafworm *Spodoptera littoralis* Bosid. (Lepidoptera: Noctuidae). Egyptian Journal of Biological Pest Control **10(1)**, 51-56.
- Mahmoudvand M, Abbasipour H, Sheikhi Garjan A, Bandani AR.** 2012. Decrease in pupation and adult emergence of *Plutella xylostella* (L.) treated with hexaflumuron. Chilean Journal of Agricultural Research **72(2)**, 206-211.
<http://dx.doi.org/10.4067/S071858392012000200007>.
- Mathai S, Nair VSK.** 1983. Effects of precocene II on last instar larvae of *Spodoptera mauritia* (Lepidoptera: Noctuidae). Current Science **52**, 376-377.
- Mathai S, Nair VSK.** 1984. Morphogenetic effects of precocene II on the larval stages of *Spodoptera mauritia* Boisid. (Lepidoptera: Noctuidae). In: Proceedings of Fifth All India Symposium on Developmental Biology (Agarwal SK, Goel SC. Eds.) 59-66 p.
- Miles M, Lysandrou M.** 2002. Evidence for negative cross resistance to insecticides in field collected *Spodoptera littoralis* (Boisd.) from Lebanon in laboratory bioassays. Mededelingen (Rijksuniversiteit te Gent. Fakulteit van de Landbouwkundige en Toegepaste Biologische Wetenschappen), **67**, 665.
- Mitsuoka T, Takita M, Kanke E, Kawasaki H.** 2001. Ecdysteroid titer, responsiveness of prothoracic gland to prothoracicotropic hormone (PTTH), and PTTH release of the recessive trimolter strain of *Bombyx mori* in extra-ecdysed larvae by JHA and 20E application. Zoological Science, Japan **18(2)**, 235-240.
<https://doi.org/10.2108/zsj.18.235>
- Moroney MJ.** 1956. Facts from figures. (3rded.). Penguin Books Ltd., Harmondsworth, Middlesex, 228 p.

- Mosallanejad H, Smaghe G.** 2009. Biochemical mechanisms of methoxyfenozide resistance in the cotton leafworm *Spodoptera littoralis*. Pest Management Science **65**, 732-736.
<http://dx.doi.org/10.1002/ps.1753>
- Moya P, Castillo M, Primo-Yufera E, Couillaud F, Martinez-Manez R, Garcera MD, Miranda HA, Primo J, Martinez-Pardo R.** 1997. Brevioxime: a new juvenile hormone biosynthesis inhibitor isolated from *Penicillium brevicompactum*. Journal of Organic Chemistry **62**, 8544-8545.
<http://dx.doi.org/10.1021/jo970397y>
- Muraleedharan D, Varghese A, Abraham G, Mathews RA.** 1986. Effect of precocene-II on endocrines, feeding and digestion in the semilooper caterpillar, *Achoea Janata*. In: "Insect Neurochemistry and Neurophysiology", Publisher: Springer, 307-314 p.
- Nandi PS, Chakravarty K.** 2011. Juvenoids and anti-Juvenoids as third generation pesticide to control lepidopteran field crop pests. Indian Streams Research Journal **1(6)**, 15 p.
<http://dx.doi.org/10.9780/22307850>
- Nasiruddin M, Mordue (Luntz) AJ.** 1994. The protection of barley seedlings from attack by *Schistocerca gregaria* using azadirachtin and related analogues. Entomologia Experimentalis et Applicata **70**, 247-252 .
<https://doi.org/10.1111/j.1570-7458.1994.tb00753.x>
- Nassar MM, Hafez ST, Farrag AM, Abahussain MO.** 1999. Efficacy of BaySir-8514 and precocene II against the grey flesh fly *Parasarcophaga dux* (Thomson) (Diptera: Sarcophagidae). Journal of the Egyptian Society of Parasitology **29(3)**, 697-707.
- Osman KSA.** 1988. Effect of precocene-3 on the hormonal regulation of oogenesis and larval mortality in *Aiolopus thalassinus* (Fabr.) (Saltatoria: Acrididae). Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie **6**, 464-487.
- Osman EE, Rarwash I, El-Samadisi MM.** 1984. Effect of the anti-moulting agent "Dimilin" on the blood picture and cuticle formation in *Spodoptera littoralis* (Boisd.) larvae. Bulletin of Entomological Society of Egypt (Economic Series) **14**, 3-46.
- Proksch P, Proksch M, Towers GHN, Rodriguez E.** 1983. Phototoxic and insecticidal activities of chromenes and benzofurans from *Encelia*. Journal of Natural Products **46**, 331-334.
<http://dx.doi.org/10.1021/np50027a006>
- Raslan SAA.** 2002. Preliminary report on initial and residual mortality of the natural product, spinosad for controlling cotton leaf worm egg masses. In: 2002 Cotton Season at Sharkia Governorate, Egypt. 2nd International Conference, Plant Protection Research Institute, Cairo, Egypt, 21-24 December **1**, 635-637.
- Relyea RA.** 2009. A cocktail of contaminants: how mixtures of pesticides at low concentrations affect aquatic communities. Oecologia **159**, 363-376.
<http://dx.doi.org/10.1007/s00442-008-1213-9>
- Retnakaran A, Granett J, Ennis T.** 1985. Insect growth regulators. In: "Comprehensive Insect Physiology, Biochemistry and Pharmacology" (Kerkut, G.A., Gilbert, L.I., eds.), vol. 12, pp: 529-601. Pergamon, Oxford.
- Richard AG.** 1957. Cumulative effects of optimum and suboptimum temperatures on insect development. In: "Influence of Temperature on Biological Systems" (Johnson FH. ed.). Ronald Press Comp., New York, 145-162.
- Riddiford LM.** 1994. Cellular and molecular action of juvenile hormone. I. General considerations and premetamorphic actions. Advances in Insect Physiology **24**, 213-243.
[https://doi.org/10.1016/S0065-2806\(08\)60084-3](https://doi.org/10.1016/S0065-2806(08)60084-3)
- Riddiford LM.** 2008. Juvenile hormone action: A 2007 perspective. Journal of Insect Physiology **54**, 895-901.
<https://doi.org/10.1016/j.jinsphys.2008.01.014>

- Roberto CF, Sosa ME, Favier LS, Penna F, Guerreiro E, Giordano OS, Tonn CE.** 1998. Growth inhibitory activities of benzofuron and chromene derivatives towards *Tenebrio molitor*. *Journal of Natural Products* **61**, 1209-1211. <http://dx.doi.org/10.1021/np9800248>
- Rust MK, Hemsarath WLH.** 2017. Intrinsic activity of IGRs against larval cat fleas. *Journal of Medical Entomology* **54(2)**, 418-421.
- Sariaslani FS, McGee LR, Ovenall DW.** 1987. Microbial transformation of precocene II: Oxidative reactions by *Streptomyces griseus*. *Applied and Environmental Microbiology* **53(8)**, 1780-1784.
- Saxena RC, Sharma M, Kumar ML, Bansal SK, Shrivastava D.** 1994. Developmental effects of 6, 7-dimethoxy-2,2-dimethyl chromene on the preimaginal stages of *Anopheles stephensi*. *Proceedings of the Academy of Environmental Biology, India* **3**, 181-184.
- Shonouda ML, Osman SL.** 2000. New botanical derivatives, used in medicinal reparations, showing bioactive action on insect pests. 1-Toxicological effect on the development of *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae). *Journal of Egyptian German Society of Zoology* **31**, 227-234.
- Shuto A, Kuwano E, Eto M.** 1988. Synthesis of two optical isomers of the insect anti-juvenile hormone agent fluoromevalonolactone (FMev) and their biological activities. *Agricultural and Biological Chemistry* **52(4)**, 915-919. <https://doi.org/10.1271/bbb1961.52.915>
- Singh S, Kumar K.** 2011. Sensitivity of last larval stadium of *Chrysomya megacephala* (Fabricius) to anti-allatin ageratochromene Precocene II. *Indian journal of Experimental Biology* **49**, 600-608.
- Smagghe G, Degheele D.** 1994. The significance of pharmacokinetics and metabolism to the biological activity of RH-5992 (tebufenozide) in *Spodoptera exempta*, *Spodoptera exigua* and *Leptinotarsa decemlineata*. *Pesticide Biochemistry and Physiology* **49**, 224-234. <https://doi.org/10.1006/pest.1994.1050>
- Smagghe G, Carton B, Wesemael W, Ishaaya I, Tirry L.** 1999. Ecdysone agonists-mechanism of action and application on *Spodoptera* species. *Pesticide Science* **55**, 343-389. [https://doi.org/10.1002/\(SICI\)10969063\(199903\)55:3<386::AID-PS922>3.0.CO;2-5](https://doi.org/10.1002/(SICI)10969063(199903)55:3<386::AID-PS922>3.0.CO;2-5)
- Srivastava S, Kumar K.** 1996. Effect of precocenes on metamorphosis of flesh fly, *Sarcophaga ruficornis* F. *Indian Journal of Experimental Biology* **34**, 547-553.
- Srivastava S, Kumar K.** 1997. Precocene I and II induced metamorphosis in a noctuid moth, *Spodoptera litura* Fabr. *The Proceedings of the National Academy of Sciences, India* **67(B)**, 213-226.
- Srivastava S, Kumar K.** 1999. Juvenilizing effects of ethoxyprecocene in a lepidopteran insect. *Indian Journal of Experimental Biology* **37(4)**, 379-383. <http://hdl.handle.net/123456789/19032>
- Staal GB.** 1982. Insect control with growth regulation interfering with the endocrine system. *Entomologia Experimentalis et Applicata* **31**, 15-23. <https://doi.org/10.1111/j.1570-7458.1982.tb03115.x>
- Staal GB.** 1986. Anti-juvenile hormone agents. *Annual Review of Entomology* **31**, 391-429.
- Subrahmanzam B, Müller T, Rembold H.** 1989. Inhibition of turnover of neurosecretion by azadirachtin in *Locusta migratoria*. *Journal of Insect Physiology* **35**, 493-500. [https://doi.org/10.1016/0022-1910\(89\)90056-5](https://doi.org/10.1016/0022-1910(89)90056-5)
- Szczepanik M, Obara R, Szumny A, Gabryś B, Halarewicz-Pacan A, Nawrot J, Wawrzńczyk C.** 2005. Synthesis and insect antifeedant activity of precocene derivatives with lactone moiety. *Journal of Agricultural and Food Chemistry* **53**, 5905-5910. <http://dx.doi.org/10.1021/jf058034j>
- Tarrant C, Cupp EW, Bowers WS.** 1982. The effects of precocene II on reproduction and development of triatomine bugs (Reduviidae: Triatominae). *American Journal of Tropical Medicine and Hygiene*. **31(2)**, 416-420. <http://dx.doi.org/10.4269/ajtmh.1982.31.416>

- Tatar M, Kopelman A, Epstein D, Tu M-P, Yin C-M, Garofalo RS.** 2001a. A mutant *Drosophila* insulin receptor homolog that extends life-span and impairs neuroendocrine function. *Science* **292**, 107-110.
<https://doi.org/10.1126/science.1057987>
- Tatar M, Priest N, Chien S.** 2001b. Negligible senescence during reproductive diapause in *D. melanogaster*. *The American Naturalist* **158**, 248-258.
<http://dx.doi.org/10.1086/321320>
- Tiryaki D, Temur C.** 2010. The fate of pesticide in the environment. *Journal of Biodiversity and Environmental Sciences* **4(10)**, 29-32.
- Trigo JR, Campos S, Pereira AM.** 1988. Presença de alcalóides pirrolizidínicos em *Ageratum conyzoides* L. In: Simposio de Plantas Mediciniais do Brasil, Sao Paulo. (Resumos).p. 13.
- Truman JW, Riddiford LM.** 2007. The morphostatic actions of juvenile hormone. *Insect Biochemistry and Molecular Biology* **37**, 761-770.
<https://doi.org/10.1016/j.ibmb.2007.05.011>
- Van Der Gaag N.** 2000. Pick your poison. *New Internationalist* **323**, 9-11.
- Walkowiak K, Spochacz M, Rosinski G.** 2015. Peptidomimetics- A new class of bioinsecticides. *Postepy Biologii Komorki* **42(2)**, 235-254.
- Wang Y, Wang M.** 2007. The research of IGRs. *World Pesticides* **29**, 8-11.
- Wilson TG.** 2004. The molecular site of action of juvenile hormone and juvenile hormone insecticides during metamorphosis: how these compounds kill insects. *Journal of Insect Physiology* **50(2/3)**, 111-121.
<https://doi.org/10.1016/j.jinsphys.2003.12.004>
- Wu J.** 2002. The application of IGRs in agricultural pest control. *Pesticides* **41**, 6-8.
- Yasyukevich VV, Zvantsov AB.** 1999. Effects of antihormones on the susceptibility of *Anopheles sacharovi* mosquitoes to the malaria causative organism *Plasmodium gallinaceum*. *Meditinskaya Parazitologiya i Parazitarnye Bolezni* **68**, 46-48.
- Yoshida T, Shiotsuki T, Kuwano E.** 2000. Synthesis and precocious metamorphosis-inducing activity of 3-(1-alkenyl) pyridines. *Journal of Pesticide Science* **25**, 253-258.
<https://doi.org/10.1584/jpestics.25.253>
- Zhou Z, Deng G, Luo S.** 2003. Study and application of IGRs. *Guangxi. Agriculture Science* **1**, 34-36.