



RESEARCH PAPER

OPEN ACCESS

Serological evidence and associated risk factors of toxoplasmosis among pregnant women in District Mardan, Khyber Pakhtunkhwa, Pakistan

Mudassir Shah^{1,2}, Muhammad Zahid^{*1}, Sobia Attaullah¹

¹Department of Zoology, Islamia College Peshawar, KP, Pakistan

²Department of Zoology, Government Superior Science College, Peshawar, KP, Pakistan

Key words: *Toxoplasma gondii*, Seroprevalence, Antibodies, Pregnant women, ELISA.

<http://dx.doi.org/10.12692/ijb/13.2.179-186>

Article published on August 30, 2018

Abstract

Toxoplasmosis is one of the widely distributed zoonotic disease which is caused by an intracellular protozoan parasite *Toxoplasma gondii*. The objective of this study was to determine the seroprevalence and risk factors associated with infection in pregnant women of District Mardan, Khyber Pakhtunkhwa, Pakistan. Blood sample of 5mL was taken and the serum was tested for *T. gondii* IgG and IgM antibodies using Indirect Enzyme-Linked Immunosorbent Assay (ELISA) technique. Further information was collected by using preplanned questionnaire. A total of 1200 samples were tested for toxoplasmosis, out of which 39.5% were positive. Among seropositive pregnant women the seropositivity for IgG, IgM and both IgG and IgM antibodies was 43.03%, 40.5% and 16.45% respectively. The seropositivity was highest (58.18%) in age group 36-45 whereas lowest (33.92%) in age group 26-35 which was statistically significant. The findings of the present study suggest that *T. gondii* infection is widely spread in the study area which may constitute a significant risk for congenital defects, stillbirths and abortions during pregnancy in women. Screening for the infection among pregnant women can be done twice at least during pregnancy to protect both mother and child from toxoplasmosis. There should be proper awareness in the educational institutes as well as community awareness programs about the risk factors of toxoplasmosis.

* **Corresponding Author:** Muhammad Zahid ✉ drmzahid@icp.edu.pk

Introduction

Toxoplasmosis is a zoonotic infection which is caused by a protozoan parasite called *Toxoplasma gondii* (Kijlstra and Jongert, 2009). *T. gondii* was first discovered by Manceaux and Nicolle in 1908 in a rodent, *Ctenodactylus gundi* of North Africa. In the same year it was also found by Splendor in Brazil in a rabbit. Its name was based on its morphology (toxo: arc or bow, plasma: life) (Ferguson, 2009). The species name was patented from a rodent (*Ctenodactylus gundi*) from which parasite was isolated and genus name referring to crescent shape of the parasite (Black and Boothroyd, 2000).

T. gondii is found inside the cell and is worldwide in distribution (Shah *et al.*, 2013). All warm-blooded animals are infected by this parasite (Dubey, 2008). Tachyzoites, bradyzoites, sporozoites are three infectious stages of *T. gondii* (Sibley *et al.*, 2009). It completes its sexual phase of life cycle in wild animals and domestic cats which act as definitive hosts and its asexual phase of life cycle is completed in warm blooded animals and humans which act as definitive hosts. Indirect way of infection is through taking improperly cooked meat or vegetables already contaminated with *T. gondii* (Shah *et al.*, 2014). Man is directly infected by consumption of meat and vegetables, direct contact with contaminated cat litter boxes, blood transfusion (Singh, 2003), transplantation of organs (Dubey and Jones, 2008).

Different studies have shown that warm and humid environment is responsible for the high prevalence of toxoplasmosis (Studeníčová *et al.*, 2006). About 33% of the human's population have this parasite (Shah *et al.*, 2013). Infection rate varies geographically in pregnant women. The prevalence rate in Korea is 3.7% and in Saudi Arabia 24.1% (Aqeelyet *et al.*, 2014), in London 17.3% (Flatt and Shetty, 2012) in Thailand 28.3% (Nissapatorn and Sawangjaroen, 2011), 68.6% in Brazil (Srokaet *et al.*, 2010), 92.5% in Ghana (Ayi *et al.*, 2009). The overall prevalence rate of toxoplasmosis in human population in different countries also varies. The prevalence rate in Taiwan is 84%, USA is 10-70%, Iran 55%, Egypt 44%, Peru 38%, Jordan 37%, India 18% (Bari and Khan, 1990).

Infection may lead to pneumonia, pericarditis, and neurologic disorders in immunocompetent hosts (Lee *et al.*, 2012). Infections goes asymptomatic in case of most immune competent individuals and results in chronic persistence of dormant cysts in tissue or the symptoms are mild (Sarkar *et al.*, 2012) but in case of immune-compromised individuals such as in HIV patients, the persons are at risk of producing acute toxoplasmosis. In 10% of HIV patients it results in cervical lymphadenopathy or ocular disease. Mortality rate in USA and Europe due to toxoplasmosis in AIDS patients is about 10% and 30% respectively (Hill and Dubey, 2002).

If the pregnant women are infected by this parasite it results in toxoplasmic encephalitis, blindness, foetal abnormalities, abortion and even stillbirth (Cook *et al.*, 2000). *T. gondii* infection also results in abortion and infertility in women (Shah *et al.*, 2014). Clinical observations in human include enlargement of the lymph node which results in high fever, headache, muscle ache and sore throat (Hill and Dubey, 2002). The tachyzoite multiply rapidly. In humans it causes congenital defects and in case of domestic livestock it results in abortion (Chaudhary *et al.*, 2006). There is 40% death chance in fetus if mother is infected during pregnancy (Jones *et al.*, 2001). Congenital toxoplasmosis during the first trimester of pregnancy results in spontaneous abortion and still birth whereas acquired infection in late pregnancy passes asymptomatic in the new-born but may results in retinochoroiditis and even complete blindness (Shah *et al.*, 2013). Symptoms in cats are pancreatitis, pressing of head, teeth grinding, low coordination, voice abnormality, uneasy feelings, circling, weak reflex actions, paralysis (Holzworth, 1987)

Transmission of disease depends on environmental conditions, culture, and habits of eating (Kapperud *et al.*, 1996). Toxoplasmosis diagnosis is performed by detection of IgG and IgM antibodies against *T. gondii*. IgG avidity test is carried out for acute infection with low IgG avidity and for chronic infection with high IgG avidity (Liesenfeld *et al.*, 2001). Serologic tests are carried out to find out IgG antibodies for testing infection. Immunoglobulin M (IgM) level is measured

by Avidity test to find out the infection time in pregnant mother. For detection of IgG and IgM antibodies in serum Elisa test is performed (Hasan, 2011) Latex Agglutination Test kit (LAT) is commercially used for IgG antibodies (Tasawar *et al.*, 2012). Sabin-feldmen dye test (SFDT), complement fixation test, intradermal skin test, indirect hemagglutination assay (IHA) and Modified Agglutination Test (MAT) are other serological tests which are carried out to detect the infection (Goz *et al.*, 2007). The parasite can be observed directly in the tissues by staining or the diagnosis can also be made by biopsy. Polymerase Chain Reaction (PCR) is very important, sensitive and efficient method in diagnosis of toxoplasmosis (Aldebert *et al.*, 2011). The objective of the present study is to determine the seroprevalence of anti-*T. gondii* antibodies in pregnant women and to identify the risk factors associated with concern infection in pregnant women of District Mardan, Khyber Pakhtunkhwa, Pakistan.

Materials and methods

Study area

Samples were collected from different tehsiles (Takht Bhai, Katlang and Mardan) of District Mardan.

Sample collection and processing

A total of 5mL venous blood was collected from each individual through sterilized syringes.

Sample processing

The blood was transferred to the blood collecting tubes (EDTA). It was then centrifuged at 3000 rpm for 10 minutes, at 25°C for serum separation from RBCs, in eppendorf tubes.

Serological test for *T. gondii* antibodies

T. gondii antibodies were detected in the sera of individuals by using an indirect Enzyme-Linked Immunosorbent Assay (ELISA). The kit used was Accu Diag™ Toxo IgM and IgG ELISA Kits (The Diagnostic Automation/Cortez Diagnostics, Inc. *T. gondii* (Toxo) IgM and IgG Enzyme-Linked Immunosorbent Assay (ELISA). The Toxo IgM ELISA Kits had a 100% specificity and a 100% sensitivity, while the Toxo IgG ELISA Kits had a 100% specificity

and a 95.3% sensitivity. With each test assay calibrator and controls were run.

Cut-off calibrator value and Immune Status Ratio

For calculation of cut-off calibrator value and Immune Status Ratio (ISR) the optical densities were obtained (OD). The results were interpreted with respect to the ISR values. For IgM, a sample having $OD \leq 0.90$ was considered negative, for sample having $OD \geq 1.10$ was considered positive, and samples with $OD = 0.91-1.09$ was considered indeterminate. In case of IgG, a sample with $OD < 0.90$ was declared as negative, while samples with $OD > 1.10$ was considered positive, and OD within $0.91-1.09$ was considered intermediate (Wam *et al.*, 2016).

Statistical analysis

The data was analyzed by using SPSS software V20 (IBM corp) and was shown in tables. *P*-values less than 0.05 were considered statistically significant.

Results

A total of 1200 pregnant women were approached in which 474 (39.5%) women were seropositive in which the seroprevalence of anti-*T. gondii* antibodies were calculated, among the seropositive women 204 (43.03%) were seropositive for IgG antibodies, 192 (40.5%) were seropositive for IgM antibodies and 78 (16.45%) were positive for both IgG and IgM antibodies. Regarding the participants residential area, 411 of the study lived in rural area in which 165 (40.41%) were infected by toxoplasmosis (Table 1) while studying the anti-*T. gondii* antibodies for the suspected rural participants 81 (49.1%) were seropositive for IgG, 57 (34.5%) were seropositive for IgM and 27 (16.4%) were seropositive for both IgG and IgM antibodies. Similarly, a total of 789 pregnant women were examined in Urban area, in which 309 (39.16%) were suspected positive and were infected with toxoplasmosis. While studying the seropositivity of antibodies in Urban area 123 (39.8%) were seropositive for IgG, 135 (43.7%) were seropositive for IgM and 51 (16.5%) were seropositive for both IgG and IgM antibodies. Although there is no significant difference of the disease between rural and urban areas but there is high significant difference between the types of infection (Table 2).

Table 1. Overall seroprevalence of toxoplasmosis in pregnant women of District Mardan.

Characteristics	Total cases (N)	No. of -Ve (%)	No. of +Ve (%)	P-Value
Age				
16-25	699	435 (62.2)	264 (37.76)	0.000
26-35	336	222 (66.1)	114 (33.92)	
36-45	165	69 (41.8)	96 (58.18)	
Area				
Urban	789	480 (60.8)	309 (39.16)	0.741
Rural	411	246 (59.9)	165 (40.14)	
Education				
Uneducated	882	537 (60.9)	345 (39.1)	0.650
Educated	318	189 (59.4)	129 (40.5)	

Table 2. Age, area and education-wise prevalence of toxoplasmosis in pregnantwomen of District Mardan.

Characteristics	Total cases	No. of +Ve (%)	Type of infection			P-Value
			IgG (%)	IgM (%)	IgG+IgM (%)	
Age						
16-25	699	264 (37.76)	117(44.3)	90 (34.1)	57 (21.6)	0.001
26-35	336	114 (33.92)	48 (42.1)	51 (44.7)	15 (13.2)	
36-45	165	96 (58.18)	39 (40.6)	51 (53.1)	6 (6.2)	
Area						
Urban	789	309 (39.16)	123(39.8)	135(43.7)	51(16.5)	0.112
Rural	411	165 (40.14)	81(49.1)	57 (34.5)	27 (16.4)	
Education						
Uneducated	882	345 (39.1)	150(43.5)	135(39.1)	60 (17.4)	0.516
Educated	318	129 (40.5)	54(41.9)	57 (44.2)	18 (14.0)	

The age of the pregnant women which were examined for the toxoplasmosis were between 16 to 45 years. Among pregnant women, 699 examined were having age group 16-25, out of which 264 (37.76%) were infected with *T. gondii* infection in which the seropositivity of IgG anti-*T. gondii* antibodies was 117 (44.3%) and that of IgM antibodies prevalence was 90 (34.1%) and both IgG and IgM antibodies seroprevalence was 57 (21.6%). It shows that the seropositivity of IgG antibodies is more in pregnant women of age group 16-25. Among 1200 pregnant women, 336 were of age group 26-35. Out of which 114 (33.92%) were positive for toxoplasmosis, in which the seropositivity of IgG antibodies is 48 (42.1%) and 51 (44.7%) were seropositive for IgM antibodies and 15 (13.2%) were seropositive for both IgG and IgM antibodies. A total of 165 pregnant women of age group 36-45 were examined, in which 96 (58.18%) were infected with *T. gondii*. The analysis of the infected women of age group 36-45 shows that 39 (40.6%) were seropositive for IgG antibodies, while 51 (53.90%) were seropositive for IgM antibodies and 6 (6.2%) were detected

seropositive for both IgG and IgM antibodies. This illustrates that seropositivity of toxoplasmosis is higher in age group 36-45. The disease is highly significant in age groups (Table 2).

The data collected was also analyzed on education basis. Out of 1200 pregnant women 318 were educated, in which 129 (40.56%) were infected with *T. gondii* and 882 were uneducated in which 345 (39.11%) were detected seropositive for toxoplasmosis (Table 1).

Discussion

Toxoplasmosis is one of the life-threatening disease (Negash *et al.*, 2008). This study investigated toxoplasmosis by detecting antiToxoplasma antibodies in serum of pregnant women in rural and urban areas in District Mardan. The findings of the present study revealed that the prevalence of anti-*T. gondii* antibodies among pregnant women was 39.5% which was similar to the results of NWFP (38%) but was lower than Punjab (63%), Azad Kashmir (48%) (Bari and Khan, 1990), nijinikon, NW Cameroon 54.5% (Wam *et al.*, 2016).

The present study results were also very lower than the study carried out in Ethiopia (85.3%) (Abamecha and Awel, 2016), Yaounde, Douala 70% (Njunda *et al.*, 2011). These geographic differences in the seroprevalence may be explained by differences in rural and urban settings. This might explain differences in risk factors of toxoplasma seropositivity as well as climatic changes, geographical location and culture (Shah *et al.*, 2014).

The unexposed people are at risk of getting an acute infection which may be responsible for congenital toxoplasmosis in pregnant women or which may be life threatening in immune-suppressive patients if it reactivated. There are studies that show very low seroprevalence of the infection compared to the present study such as in South Africa 6.4% (Mwambe *et al.*, 2013), Zambia 5.87% (Frimpong *et al.*, 2017), Korea 3.7%, United Kingdom 9% (Aqeely *et al.*, 2014). In Southern Thailand the seroprevalence is 28.3% (Sroka *et al.*, 2010), 24.1% in Saudi Arabia (Aqeely *et al.*, 2014) and 30.9% in Tanzania (Nissapatorn and Sawangjaroen, 2011). Among 474, pregnant women, 43.03% were positive for IgG antibodies having chronic infection and 40.5% were positive for IgM having acute toxoplasmosis, whereas both IgG and IgM seropositivity was 16.45% which was lower than the prevalence in Northern parts of Pakistan where IgG antibodies were 46%, whereas was higher to 27.7% for IgM antibody (Bari and Khan, 1990). It was also lower to latent infection but was higher to acute infection of Cameroon whose results for IgG and IgM was 88.7%, 30.9% (Wam *et al.*, 2016). But the present results are also significantly lower to 71.8% and 67.6% respectively obtained from London (Flatt and Shetty, 2012). The latent infection prevalence is also lower to 82.3% in Ethiopia whereas the seropositivity for both IgG and IgM in current study is 16.45% which is higher than the work done in Ethiopia. Latent infection in the current study is higher than the 3.02% in Ethiopia (Abamecha and Awel, 2016). The latent infection prevalence among Pakistani pregnant women studied in Norway which is 17% (Bjerke *et al.*, 2011) is lower to the present study. Moderate prevalence of 30-50% has been found in countries of

central and southern Europe (Robert-Gangneux and Dardé, 2012) which shows similar prevalence range in present study. The overall prevalence of toxoplasmosis reported previously in Brazil was 68.37% and the chronic infection prevalence of 63.03% and acute infection of 5.33% (Da Silva *et al.*, 2015) show a higher overall and also chronic infection seroprevalence from present study but show a lower acute infection prevalence than present study. The socio-economic conditions and presence of pets and stray cats, geographic location and age has a great role in high prevalence rate of toxoplasmosis.

Conclusion

The present study findings suggest that *T. gondii* infection is widely spread in the study area which may constitute a significant risk for congenital defects, stillbirths and abortions during pregnancy in women. Screening for the infection among pregnant women can be done twice at least during pregnancy to protect both mother and child from toxoplasmosis. There should be proper awareness in the educational institutes as well as community awareness programs about the risk factors of toxoplasmosis. This is the pioneer study on toxoplasmosis in pregnant women in District Mardan. The present study also opens ways for further studies to be carried out in the concerned area. This will further create avenue for management and treatment of this disease in this population.

Acknowledgements

We are thankful to all the participants who sacrificed their reasonable time and donated their blood for the current study.

Conflict of interests

The authors declare that they have no conflict of interests.

References

Abamecha F, Awel H. 2016. Seroprevalence and risk factors of *Toxoplasma gondii* infection in pregnant women following antenatal care at Mizan Aman General Hospital, Bench Maji Zone (BMZ), Ethiopia. *BMC infectious diseases* **16**, 460. <https://doi.org/10.1186/s12879-016-1806-6>.

- Aldebert D, Hypolite M, Cavailles P, Touquet B, Flori P, Loeuillet C, Cesbron-Delauw MF.** 2011. Development of high-throughput methods to quantify cysts of *Toxoplasma gondii*. *Cytometry Part A* **79(11)**, 952-958.
<https://doi.org/10.1002/cyto.a.21138>.
- Aqeely H, El-Gayar EK, Perveen Khan D, Najmi A, Alvi A, Bani I, Mahfouz MS, Abdalla SE, Elhassan IM.** 2014. Seroepidemiology of *Toxoplasma gondii* amongst pregnant women in Jazan Province, Saudi Arabia. *Journal of tropical medicine* 2014, 1-6.
<http://dx.doi.org/10.1155/2014/913950>.
- Ayi I, Edu S, Apea-Kubi K, Boamah D, Bosompem K, Edoh D.** 2009. Sero-epidemiology of toxoplasmosis amongst pregnant women in the greater Accra region of Ghana. *Ghana medical journal* **43(3)**, 107-114.
<http://dx.doi.org/10.4314/gmj.v43i3.55325>.
- Bari A, Khan Q.** 1990. Toxoplasmosis among pregnant women in northern parts of Pakistan. *JPM* **40**, 288-289.
- Bjerke SEY, Vangen S, Holter E, Stray-Pedersen B.** 2011. Infectious immune status in an obstetric population of Pakistani immigrants in Norway. *Scandinavian journal of public health* **39(5)**, 464-470.
- Black MW, Boothroyd JC.** 2000. Lytic cycle of *Toxoplasma gondii*. *Microbiology and Molecular Biology Reviews* **64(3)**, 607-623.
DOI: 10.1128/MMBR.64.3.607-623.2000.
- Chaudhary Z, Ahmed R, Hussain S, Shakoori A.** 2006. Detection of *Toxoplasma gondii* infection in butchers and buffaloes by polymerase chain reaction and latex agglutination test. *Pakistan Journal of Zoology* **38(4)**, 333-336.
- Cook A, Holliman R, Gilbert R, Buffolano W, Zufferey J, Petersen E.** 2000. Sources of toxoplasma infection in pregnant women: European multicentre case-control study. *Commentary: Congenital toxoplasmosis-further thought for food.* *Bmj* **321(7254)**, 142-147.
DOI: <https://doi.org/10.1136/bmj.321.7254.142>.
- Da Silva MG, Vinaud MC, de Castro AM.** 2015. Prevalence of toxoplasmosis in pregnant women and vertical transmission of *Toxoplasma gondii* in patients from basic units of health from Gurupi, Tocantins, Brazil, from 2012 to 2014. *PLoS One* **10(11)**, e0141700.
<https://doi.org/10.1371/journal.pone.0141700>.
- Dubey J, Jones J.** 2008. *Toxoplasma gondii* infection in humans and animals in the United States. *International journal for parasitology* **38(11)**, 1257-1278.
<https://doi.org/10.1016/j.ijpara.2008.03.007>.
- Dubey JP.** 2008. The history of *Toxoplasma gondii*-the first 100 years. *Journal of Eukaryotic Microbiology* **55(6)**, 467-475.
<https://doi.org/10.1111/j.1550-7408.2008.00345.x>.
- Ferguson DJ.** 2009. *Toxoplasma gondii*: 1908-2008, homage to Nicolle, Manceaux and Splendore. *Memorias do Instituto Oswaldo Cruz Rio de Janeiro* **104(2)**, 133-148.
<http://dx.doi.org/10.1590/S0074-02762009000200003>.
- Flatt A, Shetty N.** 2012. Seroprevalence and risk factors for toxoplasmosis among antenatal women in London: a re-examination of risk in an ethnically diverse population. *The European Journal of Public Health* **23(4)**, 648-652.
<https://doi.org/10.1093/eurpub/cks075>.
- Frimpong C, Makasa M, Sitali L, Michelo C.** 2017. Seroprevalence and determinants of toxoplasmosis in pregnant women attending antenatal clinic at the university teaching hospital, Lusaka, Zambia. *BMC infectious diseases* **17(1)**, 10.
<https://doi.org/10.1186/s12879-016-2133-7>.
- Göz Y, Babür C, Aydin A, Kiliç S.** 2007. Seroprevalence of toxoplasmosis, brucellosis and listeriosis in horses in Hakkari, eastern region of Turkey. *Revue de Medecine Veterinaire* **158(11)**, 534-539.
- Hasan SF.** 2011. Seroprevalence of toxoplasmosis among comers to marriage in Kerbala governorate. *Kerbala Journal of Pharmaceutical Sciences* **2**, 97-102.

- Hill D, Dubey J.** 2002 Toxoplasma gondii: transmission, diagnosis and prevention. *Clinical microbiology and infection* **8(10)**, 634-640. <https://doi.org/10.1046/j.1469-0691.2002.00485.x>
- Holzworth J.** 1987. Diseases of the cat: Medicine and surgery. Philadelphia: Saunders.
- Jones JL, Kruszon-Moran D, Wilson M, McQuillan G, Navin T, McAuley JB.** 2001. Toxoplasma gondii infection in the United States: seroprevalence and risk factors. *American journal of epidemiology* **154(4)**, 357-365. <https://doi.org/10.1093/aje/154.4.357>.
- Kapperud G, Jenum PA, Stray-Pedersen B, Melby KK, Eskild A, Eng J.** 1996. Risk factors for Toxoplasma gondii infection in pregnancy: results of a prospective case-control study in Norway. *American journal of epidemiology* **144(4)**, 405-412. <https://doi.org/10.1093/oxfordjournals.aje.a008942>.
- Kijlstra A, Jongert E.** 2009. Toxoplasma-safe meat: close to reality? *Trends in parasitology* **25(1)**, 18-22. <https://doi.org/10.1016/j.pt.2008.09.008>.
- Lee S-E, Hong S-H, Lee S-H, Jeong Y-I, Lim SJ, Kwon OW.** 2012. Detection of ocular Toxoplasma gondii infection in chronic irregular recurrent uveitis by PCR. *The Korean journal of parasitology* **50(3)**, 229-231. DOI: 10.3347/kjp.2012.50.3.229.
- Liesenfeld O, Montoya JG, Kinney S, Press C, Remington JS.** 2001. Effect of testing for IgG avidity in the diagnosis of Toxoplasma gondii infection in pregnant women: experience in a US reference laboratory. *The Journal of infectious diseases* **183(8)**, 1248-1253. <https://doi.org/10.1086/319672>.
- Mwambe B, Mshana SE, Kidenya BR, Massinde AN, Mazigo HD, Michael D.** 2013. Sero-prevalence and factors associated with Toxoplasma gondii infection among pregnant women attending antenatal care in Mwanza, Tanzania. *Parasites & vectors* **6(1)**, 222. <https://doi.org/10.1186/1756-3305-6-222>.
- Negash T, Tilahun G, Medhin G.** 2008. Seroprevalence of Toxoplasma gondii in Nazareth town, Ethiopia. *East African Journal of Public Health* **5(3)**, 211-214.
- Nissapatorn V, Sawangjaroen N.** 2011. Parasitic infections in HIV infected individuals: diagnostic & therapeutic challenges. *The Indian journal of medical research* **134(6)**, 878-897. doi: 10.4103/0971-5916.
- Njunda AL, Assob JC, Nsagha DS, Kamga HL, Nde PF, Yugah VC.** 2011. Seroprevalence of Toxoplasma gondii infection among pregnant women in Cameroon. *Journal of public health in Africa* **2(2)**.
- Robert-Gangneux F, Dardé M-L.** 2012. Epidemiology of and diagnostic strategies for toxoplasmosis. *Clinical microbiology reviews* **25(2)**, 264-296. DOI: 10.1128/CMR.05013-11.
- Sarkar MD, Anuradha B, Sharma N, Roy RN.** 2012. Seropositivity of toxoplasmosis in antenatal women with bad obstetric history in a tertiary-care hospital of Andhra Pradesh, India. *Journal of health, population, and nutrition* **30(1)**, 87-92.
- Shah M, Aftab MZ, Sthanadar A, Ali PA.** 2014. Seroprevalence of Toxoplasma gondii Infection in Human Population of Mohmand Agency Khyber Pakhtunkhwa Pakistan. *Pakistan Journal of Zoology* **46(4)**.1169-1172.
- Shah M, Zahid M, Asmat P, Alam A, Sthanadar A.** 2013. Seroprevalence of Toxoplasma gondii in goats and sheep of district Mardan, Pakistan. *International Journal of Biosciences* **7(7)**, 90-97. doi: <http://dx.doi.org/10.12692/ijb/3.7.90-97>
- Shah M, Zahid M, Sthanadar AA, Asmat P, Kausar A, Jan AH.** 2013. Seroprevalence of Toxoplasma gondii infection in domestic animals of. *Journal of Coastal Life Medicine* **1(1)**, 70-73.
- Sibley LD, Khan A, Ajioka JW, Rosenthal BM.** 2009. Genetic diversity of Toxoplasma gondii in animals and humans. *Philosophical Transactions of the Royal Society B: Biological Sciences* **364(1530)**, 2749-2761. DOI: 10.1098/rstb.2009.0087.

Singh S. 2003. Mother-to-child transmission and diagnosis of *Toxoplasma gondii* infection during pregnancy. *Indian journal of medical microbiology* **21(2)**, 69-76.

Sroka J, Wójcik-Fatla A, Szymanska J, Dutkiewicz J, Zajac V, Zwolinski J. 2010. The occurrence of *Toxoplasma gondii* infection in people and animals from rural environment of Lublin region-estimate of potential role of water as a source of infection. *Annals of Agricultural and Environmental Medicine* **17(1)**,125-132.

Studeníčová C, Benčaiová G, Holková R. 2006. Seroprevalence of *Toxoplasma gondii* antibodies in a healthy population from Slovakia. *European journal of internal medicine* **17(7)**, 470-473.

DOI: <https://doi.org/10.1016/j.ejim.2006.07.007>.

Tasawar Z, Aziz F, Lashari MH, Shafi S, Ahmad M, Lal V, Hayat CS. 2012. Seroprevalence of Human toxoplasmosis in southern Punjab, Pakistan. *Pakistan Journal of Life and Social Sciences* **10(1)**, 48-52.

Wam EC, Sama LF, Ali IM, Ebile WA, Aghangu LA, Tume CB. 2016. Seroprevalence of *Toxoplasma gondii* IgG and IgM antibodies and associated risk factors in women of child-bearing age in Njinikom, NW Cameroon. *BMC research notes* **9**, 406. <https://doi.org/10.1186/s13104-016-2206-0>.