



RESEARCH PAPER

OPEN ACCESS

Estimation of heavy metals in river harrow at Pambala Khyber Pakhtunkhwa, Pakistan

Khalid Usman^{*1}, Rizwan Ullah Khan², Hameed Ur Rehman², Khalid Pervaiz³,
Inayat Ullah Malik⁴, Haleema Sadia⁵, Muhammad Ibrahim⁶, Muhammad Ali Subhani⁶,
Nisara Wahab⁶, Saiqa Bashir⁶, Nida Shah⁷

¹Department of Zoology, Hazara University, Mansehra, Khyber Pakhtunkhwa, Pakistan

²Department of Chemistry, Kohat University of Science and Technology, KUST, KP, Pakistan

³Fisheries Research & Training Institute, Government of the Punjab, Lahore, Pakistan

⁴Department of Biological Sciences, Subcampus Mianwali, University of Sargoda, Pakistan

⁵Department of Biotechnology, Balochistan University of Information Technology Engineering and Management Sciences, Quetta, Pakistan

⁶Department of Chemistry, University of Kotli, Kotli, Kashmir, Pakistan

⁷Department of Botany, Bacha Khan University, Charsada, KP, Pakistan

Key words: River, Harrow, Range, Pambala, KP, Concentration

<http://dx.doi.org/10.12692/ijb/13.1.382-386>

Article published on July 30, 2018

Abstract

A research study was conducted to explore the concentration of heavy metals in River Harrow at Pambala Khyber Pakhtunkhwa, Pakistan. For this purpose six heavy metals were selected. Sampling was carried out from Downstream, Mid Point and Upstream points of the river. Heavy metals examined in the present study were in the range of Zn 1.16-1.79ppm; Cu 1.04-1.23ppm; Cd 0.05-1.33ppm; Pb 0.05-1.24ppm; Cr 0.05-0.14ppm and Mn 0.05-0.09ppm respectively. Cd, Cu, Pb and Cr were found above the permissible limits. The present survey demonstrated that the water in this site is contaminated with heavy metals.

* **Corresponding Author:** Khalid Usman ✉ khalidusmankhattak1985@gmail.com

Introduction

Pollution of the environment by heavy metals is very prominent in areas of mining sites and reduces increasing distance away from these sites. Another contribution of anthropogenic metals of terrestrial origin is from industrial development and other activities such as agriculture, metallurgy and transport from mining activities, ground water is most vigorously polluted (Abderahman and Abu-Rukah, 2000; Buccolieri, 2006). The harmful effect of trace elements when consumed above the recommended limit can be toxic (acute, chronic or sub-chronic), and heavy metals can be neurotoxic, carcinogenic, mutagenic or teratogenic. The general symptoms of humans related to metal [e.g., Cd, Pb, As, Hg, Zn, Cu and aluminium (Al) poisoning include vomiting, convulsions, paralysis, ataxia, hemoglobinuria, gastrointestinal disorder, diarrhoea, stomatitis, tremor, depression and pneumonia (McCluggage, 1991). There has been an increasing interest in the utilization of fishes as bio-indicators of the integrity of aquatic environmental systems in recent years (Tawari and Ekaye, 2007). Heavy metals influence cellular organelles and various enzymes involved in metabolic process, detoxification, and damage repair (Wong and Shi, 2001). Quantity of health hazard metals was detected by Usman *et al.* (2017b) in natural waters of river Kabul, KP Province, Pakistan. The concentrations of the metals recorded were in the range as: Pb 0.06-4.41ppm; Zn 4.11-7.11ppm; Cd 0.42-1.46ppm; Cu 1.07-3.86ppm; Mn 0.06-2.11ppm and Cr 0.05-2.11ppm. Concentration of heavy metals was analyzed by Usman *et al.* (2017c) in the River Kabul Shah Alam tributary, Peshawar Khyber Pakhtunkhwa, Pakistan. The concentration of the heavy metals were Zn 1.2-2.0ppm; Cu 0.17-1.48ppm; Cd 0.2-0.69ppm; Pb 1.01-1.23ppm; Cr 0.04-2.01ppm and Mn 0.01-0.82ppm respectively. Measurement of heavy metals was conducted by Fawad *et al.* (2017) to know the rate of bioaccumulation of Chromium (Cr (III) in the gills, intestine, and skin and its acute toxicity to goldfish (*Carassius auratus*) fingerlings. The behavioral change occurs in the fish is that all the fingerlings of goldfish come to the corner of the aquarium and their appetite also decrease due to chemical effect.

Amount of heavy metals were analyzed by Usman *et al.* (2017a) in different sites of River Kabul on Rohu, *Labeo rohita* (Hamilton). The highest concentrations of Zn (6.00ppm) was found at Jehangera Upper site, Cu (3.05ppm) at Dalda Oil Mill Nowshera site, Cr (1.05ppm) at Jehangera Lower, Mn (2.00ppm) at Jehangera Lower, Pb (0.02ppm) at Dalda Oil Mill Nowshera site and Cd (3.0ppm) at the Jehangera Upper site. Evaluation of heavy metals were carried out by Afridi *et al.* (2017) in the common carp (*Cyprinus carpio*) collected from two different water bodies the Tarbela dam, District Haripur, and River Soan District Rawalpindi Pakistan. Concentration of Mn, Ni, Cd, Cu, Pb, Se, Zn were determined in five tissues the gills, skin, kidney, liver and muscle. The concentration of detected metals found in different tissues of same species varied for Mn: 0.43-4.96, Ni: 0.49 – 1.60, Cd: 0.06 – 0.08, Cu: 0.36 – 0.81, Pb: 0.50 – 0.74, Se: 6.17 – 17.05, Zn: 0.59 – 3.74 µg/g wet wt. The aim of the current study was to find out the estimation of Heavy metals in River Harrow at Pambala Khyber Pakhtunkhwa, Pakistan

Materials and methods

Study Area

Pambala sampling site of River Harrow is very beautiful because of its natural greenery and picnic spots. In this site of the river there is no domestic discharged which badly affect the fish population. The water of this sampling site is clear. Water flow in this site is very high. The width of the river Harrow at this site is large.



Fig. 1. Map of River Harrow at Pumbala site Khyber Pakhtunkhwa Pakistan.

Sampling of Water

Water samples were stored in clean and dry plastic bottles with screw caps and labeled. The freshly collected samples were analyzed for Heavy metals analysis at GC University Faisalabad lab by using atomic absorption.

Method for preparation of stock solution

The stock solution was prepared as 1000ppm = 1000mg/l. Then 100ppm solution was prepared from stock solution using serial dilution equation of $C_1V_1 = C_2V_2$

Determination of heavy metals in water

The water samples were first filtered with the help of filter paper and then taken in 250 ml of glass bottles and subjected to the atomic absorption spectrophotometer (Zn, Cu, Cd, Mn, Cr, Pb) at GC University Faisalabad lab.

Results and discussion

To explore the amount of heavy metals, a detail study was design in River Harrow at Pambala site Khyber Pakhtunkhwa, Pakistan. For this purpose six heavy metals were selected. Sampling was carried out from Downstream, Mid-Point and Upstream points of the river. Heavy metals examined in the present study were in the range of Zn 1.16-1.79ppm; Cu 1.04-1.23ppm; Cd 0.05-1.33ppm; Pb 0.05-1.24ppm; Cr 0.05-0.14ppm and Mn 0.05-0.09ppm respectively. Cd, Cd, Pb and Cr were found above the permissible limits. The present survey demonstrated that the water in this site in contaminated with heavy metals. A survey was carried out by Usman *et al.* (2017e) to find out the concentration of heavy metals in Jhangira Upper site of the River Kabul KP, Pakistan. The results obtained from the current study were in the range of Zn 1.11-1.97ppm; Cu 1.05-1.63ppm; Cd 0.11-0.89ppm; Pb 0.07-1.07ppm; Cr 0.01-0.11ppm and Mn 0.02-0.28ppm respectively. Analysis of heavy metals was determined by Usman *et al.* (2017f) to explore the amount of heavy metals in River Kabul at Khairabad water KP Pakistan. The heavy metals recorded were Zn 1.5-1.59ppm; Cu 1.15-1.94ppm; Cd 0.02-0.05ppm; Pb 0.15-0.73ppm; Cr 0.01-0.02ppm and Mn 0.07-0.21ppm respectively. Assessment of heavy metals was carried out by Usman *et al.* (2017f) to examine the concentration of health hazard toxic metals in River Kabul at Khazana Suger Mill Peshawar KP, Pakistan. Heavy metals concentration obtained from this study was Zn 1.13-201ppm; Cu 0.55-0.9ppm; Cd 0.02-1.22ppm; pb 1.231.84ppm; Cr 0.21-1.2ppm and Mn 0.02-0.05ppm respectively.

In a research study demonstrated by Usman *et al.* (2018) to estimate the amount of heavy metals in the water of River Kabul at Jehangira Lower KP, Pakistan. The highest concentration of the heavy metals was found Cu 0.2-1.66, Cd 0.06-0.96, Pd 0.02-1.1, Cr 0.01-0.06 while the lowest concentration was found Mn 0.11-0.23 and zinc 1.13-2.37 respectively. Usman *et al.* (2017d) work on River Kabul at Cantt area Nowshera to evaluate heavy metals. The metals which were recorded in the present study were Zn 1.13-1.85ppm; Cu 1.02-1.21ppm; Cd 0.03-1.32ppm; pb 0.04-1.23ppm; Cr 0.010.16ppm and Mn 0.00-0.00ppm respectively. Another study was carried out by Ullah *et al.* (2016) to estimate 96hr LC₅₀ value of Cadmium sulphate for the fish, *Labeo rohita*. The results showed that the median lethal concentration (LC₅₀) of Lead Nitrate for the fish, *Labeo rohita* is 24mg/l. The susceptibility of *Labeo rohita* to the lethal effect of Cadmium sulphate was dependent on duration as well as on concentration. The mortality of the fishes is directly proportional to the concentration.

Table 1. Concentration of heavy metals (ppm) in River Harrow at Pambala site KP, Pakistan.

S.No	Metals	U.S	M.P	D.S	Permissible limits
1	Zn	1.16	1.79	1.33	5.0mg/l
2	Cu	1.04	1.23	1.07	0.05mg/l
3	Cd	0.05	1.33	0.14	0.05mg/l
4	Pb	0.05	1.24	0.18	0.05mg/l
5	Cr	0.05	0.14	0.08	0.05mg/l
6	Mn	0.05	0.09	0.07	50-70mg/l




Fig. 2. Concentration of heavy metals (ppm) in River Harrow at Pambala site KP, Pakistan. U.S (Up stream); M.P (Mid point); D.S (Down stream).

Conclusion

From this study it can be concluded that heavy metals are the main health hazard pollutants which contaminate aquatic life.

These heavy metals accumulate in our food stuff and cause health problems by their toxicity. For this purpose a study was conducted to estimate the heavy metals.

Acknowledgement

Immense Thankful to Dr. Khalid Pervaiz and Dr. Inayat Ullah Malik. I am greatly thankful to Hameed Ur Rehman (Department of Chemistry). I am also thankful to my brother Dr. Wahid Raza (Department of Management Sciences ICUP) who helps me throughout in water sampling collection.

References

- Abderahman N, Abu-Rukah Y.** 2000. An assessment study of heavy metal distribution within soil in upper course of Zarqa River Basin/Jordan. *Environ. Geol* **49(8)**, 1116-1124.
- Afridi AJ, Zuberi A, Rehman HU, Khan A, Saeed K, Achakzai WM, Saddozai S, Usman K, Ateeq M, Akbar NU.** 2017. Effect of the aquatic environment of different water bodies on metal contents of common carp (*Cyprinus carpio*) collected from two different water bodies. *Journal of Entomology and Zoology Studies* **5(1)**, 388-399.
- Buccolieri A, Buccolieri G, Cardellicchio N, Dell Atti A, Di Leo A, Maci A.** 2006. Heavy metals in marine sediments of Taranto Gulf (Ionian Sea, southern Italy) *mar Chem* **99(1-4)**, 227-235.
- Fawad M, Yousafzai AM, Haseeb A, Rehman HU, Afridi AJ, Akhtar NA, Saeed K, Usman K.** 2017. Acute toxicity and bioaccumulation of chromium in gills, skin and intestine of goldfish (*Carassius auratus*). *Journal of Entomology and Zoology Studies* **5(1)**, 568-571.
- McCluggage D.** 1991. Heavy metal poisoning. Colorado, USA: NCS Magazine.
- Tawari FP, Ekaye SA.** 2007. Fish species as indicator of pollution in Ikpoba River, Benin City, Nigeria. *Review of Fish Biology and Fisheries* **17**, 21-30.
- Ullah A, Rehman HU, Saeed W, Quraish MF, Ullah Z, Haroon Rehman RU, Awais S, Raqeebullah, Ahmad W, Saeed K.** 2016. Determination of 96-hr LC50 value of cadmium for a fish, *Labeo rohita*. *Journal of Entomology and Zoology Studies* **4(5)**, 380-382.
- Usman K, Nisa Zu, Gul S, Gul S, Rehman HU, Asad M, Waqar M, Ullah, Ishaq HK.** 2017c. Contamination of Heavy metals in River Shah Alam Peshawar: (A tributary of River Kabul) Khyber Pakhtunkhwa Pakistan. *Journal of Entomology and Zoology Studies* **5(5)**, 510-512.
- Usman K, Rehman HU, Adeel S, Shah NA, Shah A, Pervaiz K, Saeed N, Mussabeha.** 2017a. Heavy Metals Accumulation in *Labeo rohita* (Hamilton) of River Kabul, KPK, Pakistan. *Biological Forum- An International Journal* **9(1)**, 01-07.
- Usman K, Rehman HU, Adeel S, Shah W, Pervaiz K, Ullah Z, Ullah R, Arsalan M.** 2017b. An Investigation on the toxicity of some trace metals in river Kabul, Khyber Pakhtunkhwa Province of Pakistan. *Biological Forum- An International Journal* **9(1)**, 95-99.
- Usman K, Rehman HU, Khudadad HS, Pervaiz K, Ali SZUA, Maqsood MJ, Jawad SM, Din AU, Ihtesham Y.** 2018. Investigation of heavy metals in River Kabul at Jehangira lower Khyber Pakhtunkhwa, Pakistan. *Journal of Entomology and Zoology Studie* **6(1)**, 100-102.
- Usman K, Rehman HU, Khudadad HS, Pervaiz K, Ali SZUA, Maqsood MJ, Jawad SM, Nazir R, Ihtesham Y.** 2017g. Evaluation of heavy metals in River Kabul at Khazana Sugar Mill Peshawar Khyber Pakhtunkhwa, Pakistan. *Journal of Entomology and Zoology Studies* **5(6)**, 2417-2419.
- Usman K, Rehman HU, Khudadad S, Pervaiz K, Ahmad N, Bilal M, Hussain St, Jawad SM, Khan M, Ali M.** 2017d. Estimation of heavy metals in River Kabul at Cantt area Nowshera Khyber Pakhtunkhwa, Pakistan. *Journal of Entomology and Zoology Studies* **5(6)**, 2275-2277.

Usman K, Rehman HU, Khudadad S, Pervaiz K, Ahmad N, Bilal M, Hussain ST, Jawad SM, Khan M, Akbar MU. 2017e. Heavy metals analysis in River Kabul at Jhangira Upper Khyber Pakhtunkhwa, Pakistan. *Journal of Entomology and Zoology Studies* **5(6)**, 2485-2487.

Usman K, Rehman HU, Khudadad S, Pervaiz K, Ali SZUA, Maqsood MJ, Jawad SM, Din AU, Ihtesham Y. 2017f. Measurement of heavy metals in River Kabul at Khairabad Khyber Pakhtunkhwa, Pakistan. *Journal of Entomology and Zoology Studies* **5(6)**, 2263-2265.

Wang S, Shi X. 2001. Molecular mechanisms of metal toxicity and carcinogenesis *Molecular and Cellular Biochemistry* **222**, 3-9.