



RESEARCH PAPER

OPEN ACCESS

Evaluation of soybean aphids against various doses of synthetic insecticides collected from different regions of China

Mubashar Hussain^{1*}, RuiBi¹, Juan Cui¹, Naveed Ahmad², Jamal Nasar³, Shu-Sen Shi¹

¹College of Agronomy, Jilin Agricultural University, Changchun 130118, China

²College of Life Sciences, Jilin Agricultural University, Changchun 130118, China

³College of Resources and Environmental Sciences, Jilin Agricultural University, Changchun, China

Key words: Soybean aphids, Insecticides, Chemical control, Population growth.

<http://dx.doi.org/10.12692/ijb/14.2.242-249>

Article published on February 27, 2019

Abstract

Aphids are a diverse herbivorous group which can infect a variety of plants direct or in direct ways. The effect of soybean aphids may lead to decline plant vigour and growth rates and ultimately reduced the yield. We evaluated effects of Five insecticides on soybean aphids collected from five different regions of China (*Changchun*, *Jiamusi*, *Zhalan*, *suihua* and *Qiqihar*) at specific doses under controlled conditions during soybean growing session 2018. Insecticides used were *Omethoate*, *Imidacloprid*, *Spirotetramat* *Malathion* and *Acetamiprid* at recommended doses of 10, 30, 50, 100 and 150. Our results indicated that, In case of *Jiamusi* aphids, *Malathion* dose 10 showed relatively greater effect (19.5a) than others. In case of *Changchun* aphids, *Spirotetramat* dose 10 showed (19.33a) while *Qiqihar* aphids at dose 10 in response to *Spirotetramat* showed (19a). In case of *suihua* aphids, *Malathion* using 10 multiple gives 17a and *Zhalan* aphids against *Omethoate* at dose 10 showed minimal result of (16.5a). In conclusion, our findings revealed the effectiveness of all listed insecticides against all soybean aphids, however, the toxicity was determined from maximum to minimum in this sequence, *Malathion* > *Spirotetramat* > *Omethoate*.

* Corresponding Author: Mubashar Hussain ✉ mubashar.ento@yahoo.com

Introduction

Soybean aphid also known as *Aphis glycines* is the most invasive pest found in Asia (Ragsdale *et al.*, 2011). Soybean aphids act on phloem sap and ultimately affect soybean seed size and yield (Beckendorf *et al.*, 2008). It is the basic primary insect which is recently managed by soybean growers in North America (Hurley and Mitchell, 2017). However, aphid populations can still occasionally build to economically damaging levels on various plants, which ask for treatment with immediate insecticides to safeguard the worldwide yields. *A. glycines* are tiny, holding light yellowish green morphology along with 2 black cornicles. Their combination of morphological characteristics as well as its pattern of colonization upon soybean, distinguishes it from various other types of aphid species. (Voegtlin *et al.*, 2004).

The major symptoms of soybean aphid infested plants exhibits a variety of morphological, chemical, biochemical, and environmental changes such as stunted growth, plant senescence, distorted and wrinkled foliage, irregular roots, stems and leaves reduced branch size and minimal seed count (He *et al.*, 1995; Lin *et al.*, 1992, 1994; Wang *et al.*, 1962, 1996; Wu *et al.*, 1999). In China, Among many soybean insect pests, *Aphis glycines* Matsumura (Hemiptera: Aphididae) is attributed as severe threat to overall production of soybean (Sun *et al.*, 2000; Wang *et al.*, 1962, 1996; Wu *et al.*, 1999; Yue *et al.*, 1989). During 1998, a 30% of heavy loss in the total yield along with 112.5 million kilograms reduction was caused due to soybean aphid infestation at Suihua and Heilongjiang regions of china (Sun *et al.*, 2000). Moreover, In Japan, Indonesia, Russia, The Philippines, Australia and recently Canada and United states have been reporting severe invasion of soybean aphids (APPPC, 1987; CAB International, 2001; Fletcher and Desborough, 2002; USDA, 2002; Venette, 2004).

Numerous tactics have been applied to tackle the control of soybean aphids worldwide in order to minimize the effect and population densities of

soybean aphids as per suggestion of integrated pest management (IPM) programs. . However, foliar insecticides specifically pyrethroids and organophosphates, proved to be the basic and primary way for the suppression (Hodgson *et al.*, 2012). Multiple insecticides against soybean aphid have been studied such as omethoate, deltamethrin, cyhalothrin, phosalone, fenvalerate, Phorate, imidacloprid (Chen and Yu, 1988; Heinrichs *et al.*, 1984; Qu *et al.*, 1987; Ragsdale *et al.*, 2007; Tabadkani *et al.*, 2017). The purpose of our study was to find the efficiency, activity and chemical control of various groups and doses of insecticides applied on soybean aphids taken from five different parts of china.

Materials and methods

Experimental location and layout

Experiment was conducted at College of Agronomy, Jilin Agricultural University, Changchun, China. Soybean variety (agriculture entomology 38) was grown in pots with controlled temperature $26\pm1^{\circ}\text{C}$ in green house. Water was given to the plants two times a day along with sulphur and potassium fertilizers. Aphids sample were collected from native soybean crops of five different areas of china including (Changchun, Jiamusi, Zhalan, suihua and Qiqihar). These were multiplied on the soybeans in controlled environment with $26\pm1^{\circ}\text{C}$, $60\pm5\%$ relative humidity [RH] and a photoperiod of 16:8h, after two generations fresh colonies were obtained and subjected to chemicals. Insecticides were bought from Guangzhou, China.

Chemical control

In case of Changchun aphids we prepared five selected insecticides (Omethoate, Imidacloprid, Spirotetramat Malathion and Acetamiprid) with five different doses (10, 30, 50, 100 and 150). Treatments of Changchun aphid with aforementioned groups of insecticides were carried out in a 0.5 ml solution when the number of soybean aphids reached to its optimum growth followed by 30-40 minutes of drying. Each group of insecticide with multiple doses were applied one by one with the help of 1 foot long

glass tubes. Each reaction included 20 aphids per tube. The reaction was performed in triplicates. Data was recorded 5-6 hours post treatment of insecticides and compared with control group (o) of each insecticide. Similar methodology was followed for *Jiamusi* aphids, *Zhalan* aphids, *suihua* aphids and *Qiqihar* aphids. The comparative data analyses for each group against selected insecticides were calculated according to complete Randomized design (CRD) by Statistix 8.1 software.

Aphids multiplication

Five different groups of soybean aphids were sampled from different regions of China. (See methods for details). These five selected groups were further multiplied by feeding soybean variety (agriculture entomology 38) until third generation was obtained. The results indicated that these aphids were actively invading our soybean variety hence confirming that feeding soybean variety (agriculture entomology 38) was not naturally resistant to these groups of aphids.

Preparation of multiple doses of insecticides

Multiple doses (5) for each group of insecticides were

prepared in the following concentrations.

Water solution = 1000ml water+ CH₃COOH3
1000µl+TritonX-100 (1000µl)

Insecticidal solution = water solution 100ml+1000µl
insecticide

Dose 10 = 10ml of water solution+1000µl of
insecticide solution

Dose 30 = 30ml water solution + 1000µl
insecticide solution

Dose 50 = 50ml water solution + 1000µl
insecticide solution

Dose 100 = 10ml water solution + 100µl
insecticide solution

Dose 150 = 15ml water solution +100µl
insecticide solution

Results

Response of Changchun aphids

The chemical control of Changchun aphids on five groups of insecticides was analyzed after the extensive treatment. The results revealed that *Spirotetramat* shows relatively higher effect (19.33a) at dose 10 compare to other groups.

Table 1. The effect of *Changchun* aphids against five different groups of insecticides.

| Insecticide name | Dose o | Dose 10 | Dose 30 | Dose 50 | Dose 100 | Dose 150 | LSD | CV% |
|----------------------|--------|------------|------------|---------|----------|-------------|-----|-------|
| <i>Omethoate</i> | 2.0e | 17.6 a | 15.3 b | 14.3 b | 12.3 c | 9.6d | 2.2 | 6.58 |
| <i>Imidacloprid</i> | 2.0d | 18.3a | 19.0a | 17.6ab | 16.0bc | 15.6c | 2.2 | 6.7 |
| <i>Spirotetramat</i> | 2.0c | 19.33a | 17.33a | 16.6ab | 14.3b | 2.3c | 2.2 | 13.0 |
| <i>Malathion</i> | 2.0d | 16.5a | 12.5ab | 13ab | 10bc | 5.5cd | 2.4 | 25.87 |
| <i>Acetamiprid</i> | 2.0e | 11.6a | 8.6b | 8.3b | 6.6c | 4.3d | 2.2 | 11.26 |

In case of *Imidacloprid* exhibits nearly similar effect (19a) with previous one however the dose dependency was found different 30. Furthermore, *Omethoate response against Changchun aphids was recorded* (17.6a) when the dose was kept 10. Lastly the results of *Malathion on dose 10* were recorded maximum (16.5a) and *Acetamiprid on dose10 showed* (11.6a) response (Table 1). All of the given groups of insecticides and their multiple doses were found non significantly variable from each other at 5% probability in comparison to control.

Response of Jiamusi aphids

The chemical control experiment in case of *Jiamsui* aphids on five various groups of insecticides was recorded after treatment. Here *Malathion* showed maximum effect (19.5a) at dose 10 comparatively. The second group that exhibits maximum response was found *Omethoate* (17.5a) when encountered on dose 10. Afterwards, *Spirotetramat response against Changchun aphids was recorded* (17a) at dose 10. The data of *Imidacloprid* and *Acetamiprid on dose 10* exhibits (15.5a) and (10.6a) respectively (Table 2).

Table 2. The effect of *Jiamsui* aphids against five different groups of insecticides.

| Insecticide name | Dose 0 | Dose 10 | Dose 30 | Dose 50 | Dose 100 | Dose 150 | LSD | CV% |
|----------------------|--------|---------|---------|---------|----------|----------|-----|-------|
| <i>Omethoate</i> | 1.5d | 17.5a | 15ab | 11.5b | 6.5c | 4cd | 2.4 | 16.37 |
| <i>Imidacloprid</i> | 1.5d | 15.5a | 14ab | 12.5bc | 12.5bc | 10.5c | 2.4 | 9.39 |
| <i>Spirotetramat</i> | 1.5d | 17a | 15ab | 13bc | 15ab | 11.5c | 2.4 | 8.88 |
| <i>Malathion</i> | 1.5d | 19.5a | 16.5ab | 15b | 9c | 5.5cd | 2.4 | 16.3 |
| <i>Acetamiprid</i> | 1.6e | 10.6a | 9.3b | 6.3c | 5.3c | 3.3d | 2.2 | 11.57 |

All of the given groups of insecticides and their multiple doses were found non significantly variable from each other at 5% probability in comparison to control.

Response of *Zhalantun* aphids

Experimental control of *Zhalantun* aphids upon five different types of insecticides were tested post-treatment under controlled environment.

Data showed that *Omethoate* gives best response (16.5a) at dose 10 in comparison to others.

Table 3. The effect of *Zhalantun* aphids against five different groups of insecticides.

| Insecticide name | Dose 0 | Dose 10 | Dose 30 | Dose 50 | Dose 100 | Dose 150 | LSD | CV% |
|----------------------|--------|---------|---------|---------|----------|----------|-----|-------|
| <i>Omethoate</i> | 2.0e | 16.5a | 13b | 12bc | 9.0d | 9.5cd | 2.4 | 11.85 |
| <i>Imidacloprid</i> | 2.0c | 12a | 7.5b | 8.5ab | 7b | 5bc | 2.4 | 25.4 |
| <i>Spirotetramat</i> | 2.0e | 15.5a | 13ab | 12bc | 9cd | 8d | 2.4 | 13.3 |
| <i>Malathion</i> | 2.0e | 11.5a | 9.5b | 8.5b | 5.5c | 4c | 2.4 | 8.45 |
| <i>Acetamiprid</i> | 2.0a | 1.5ab | 1b | 0c | 0c | 0c | 2.4 | 38.5 |

The second group that exhibits maximum response was found *Spirotetramat* (15.5a) at the same dose as first one. Then, *Imidacloprid* response was recorded (12a) 10 dose. The results of *Malathion* and *Acetamiprid* on dose 10 exhibits (11.5a) and (1.5a) respectively (Table 3). All of the given groups of insecticides and their multiple doses were found none significantly variable from each other at 5% probability in comparison to control.

Response of *Suihua* aphids

Suihua aphids and their chemical control with five multiple types of insecticides were analyzed according to the previous method under controlled conditions.

The variable datasets were recorded with maximum response in case of *Malathion* (17a) at dose 10 in comparison to others. After *Malathion*, another group of insecticide named *Spirotetramat* exhibits (15a) at the same dose of 10. Furthermore, *Imidacloprid* and *Omethoate* showed a similar pattern of response of (14a) when the dose was kept as 10.

Table 4. The effect of *Suihua* aphids against five different groups of insecticides.

| Insecticide name | Dose 0 | Dose 10 | Dose 30 | Dose 50 | Dose 100 | Dose 150 | LSD | CV% |
|----------------------|--------|---------|---------|---------|----------|----------|-----|------|
| <i>Omethoate</i> | 1.5d | 14a | 10b | 9bc | 8bc | 6c | 2.4 | 16.4 |
| <i>Imidacloprid</i> | 1.5e | 14a | 13ab | 11.5bc | 10.5c | 8.5d | 2.4 | 8.3 |
| <i>Spirotetramat</i> | 1.5c | 15a | 12ab | 6.5bc | 5bc | 2.5c | 2.4 | 43.7 |
| <i>Malathion</i> | 1.5e | 17a | 15.5ab | 13.5bc | 11.5c | 8d | 2.4 | 10.3 |
| <i>Acetamiprid</i> | 1.5ab | 6a | 2.5ab | 2.5ab | 2ab | 0b | 2.4 | 88.6 |

The data of *Acetamiprid* at dose 10 exhibits (6a) which was found the least recorded statistics in *Suihua* aphids (Table 4). All of the given groups of insecticides and their multiple doses were found non significantly variable from each other at 5% probability in comparison to control.

Response of *Qiqihaer* aphids

The effect of *Qiqihaer* aphids was tested with five multiple types of insecticides were analyzed. Data was recorded for *Spirotetramat* to be the best among all selected insecticides which was (19a) at 10 dose followed by *Malathion* and *Imidacloprid* exhibiting

(14a) when the dose was kept as 10. Minimum response of qiqihaer aphids were recorded for *Omethoate* and *Acetamiprid* giving (10.5a) and (8a). Both of these insecticides showed this response on

dose 10 respectively. All of the given groups of insecticides and their multiple doses were found non significantly variable from each other at 5% probability in comparison to control (Table 5).

Table 5. The effect of *Qiqihaer* aphids against five different groups of insecticides.

| Insecticide name | Dose 0 | Dose 10 | Dose 30 | Dose 50 | Dose 100 | Dose 150 | LSD | CV% |
|----------------------|--------|---------|---------|---------|----------|----------|-----|-------|
| <i>Omethoate</i> | 2.5cd | 10.5a | 7b | 5bc | 3.5cd | 1d | 2.4 | 28.16 |
| <i>Imidacloprid</i> | 2.5d | 14a | 10b | 8.5bc | 6cd | 5cd | 2.4 | 20.6 |
| <i>Spirotetramat</i> | 2.3f | 19a | 17.3b | 12.3d | 13.6c | 10.7e | 2.2 | 4.2 |
| <i>Malathion</i> | 2.5e | 14a | 12.5a | 9.5b | 6.5c | 4.5d | 2.4 | 7.8 |
| <i>Acetamiprid</i> | 2.5b | 8a | 4ab | 3b | 2b | 1.5b | 2.4 | 48.1 |

Discussion

Soybean aphid is a serious insect of soybean crop. Before the outbreaks of soybean aphids a few amount of insecticides were used but these days insecticides are very common tool to control populations of soybean aphids and it was used as primary tool for soybean aphids control in United states in first decade. There are many kinds of insecticides available in market (Fernandez Cornejo and Jans 1999), (DiFonzo 2009, Hodgson *et al.*, 2010), (Beckendorf *et*

al., 2008).

(Faheem *et al.*, 2016) applied Malathion on *S. graminum* and observed that Malathion is highly effective against *S. graminum*, while in case of our study Malathion dose 10 shows 19.5a to Jiamusi aphids, 17a to suihua aphids, 16.5a to Changchun aphids, 14a to Qiqihaer aphids and 11.5a to Zhalantun aphids.

Table 6. Maximum damage shown by specific dose of insecticides against specific soybean aphids.

| Insecticide with Dose | Aphids | Damage |
|---------------------------------|------------------|--------|
| 1. <i>Malathion</i> dose 10 | Jiamusi aphids | 19.5a |
| 2. <i>Spirotetramat</i> dose 10 | Changchun aphids | 19.33a |
| 3. <i>Spirotetramat</i> dose 10 | Qiqihar aphids | 19a |
| 4. <i>Malathion</i> dose 10 | Suihua aphids | 17a |
| 5. <i>Omethoate</i> dose 10 | Zhalan aphids | 16.5a |

(Magalhaes *et al.*, 2008) observed a high temporal variation in field efficacy for imidacloprid against soybean aphid and they proved that imidacloprid showed good lethal and sublethal results against soybean aphid reproduction and survival. (Shafique *et al.*, 2016) and (Royer *et al.*, 2011) used Imidacloprid to control *S. graminum* and results proved that Imidacloprid gave maximum decrease in populations of *S. graminum* and (Faheem *et al.*, 2016) showed that imidacloprid showed only 41% reduction in *S. graminum* and in our study Imidacloprid showed 18.3a in Changchun aphids, 15.5a in Jiamsui aphids, 14a in

Qiqihar aphids and Suihua aphids and 12a in Zhalantun aphids.

(Y. Gong *et al.*, 2016) applied spirotetramat to cotton aphids for five days and revealed that it gives a slow effect on killing cotton aphids, their results showed that its lethal and sublethal doses decreases the fecundity of cotton aphids, LC75 dose of spirotetramat decreased 90% of cotton aphid offspring. While in our study spirotetramat 19.33a in Changchun aphids, 19a in Qiqihaer aphids, 17a in Jiamsui aphids, 15.5a in Zhalantun aphids and 15a in

Suihua aphids.

Previous studies indicate that acetamiprid causes reduction of populations against pomegranate aphid, *A. punicae* and its activity is not disturbed with passage of time like imidacloprid activity is decreased with time, it is also effective against *C. carnea* and *S. corolla* (A. Abd-Ella *et al.*, 2015). While our study shows acetamiprid shows higher effect 11.6a in Changchun aphids while minimum effect 1.5a in zhalantun aphids.

(Bishop *et al.*, 1998) revealed that Omethoate influences the population, structure and new emerging generation of lucerne flea, omethoate has effective, immediate and long lasting effect on populations of lucerne flea. Our studies reveals omethoate activity against changchun aphids is 17.6a while jiamsui aphids 17.5a, 16.5a in Zhalantun aphids, Suihua aphids 14a and 10.5a in Qiqihaer aphids.

Our experiment reveals that different insecticides shows different effects in different kinds of aphids while the activity of insecticides seen is Malathion followed by Spirotetramat followed by Omethoate.

Conclusion

We concluded that Jiamusi aphids were more killed by Malathion dose 10 as compared to others while against Changchun aphids and Qiqihqar aphids, Spirotetramat dose 10 performed better than others. Meanwhile, Suihua aphids died more by Malathion dose 10 and in case of Zhalan aphids, Omethoate at dose 10 gave better results than others. So, we recommend Malathion dose 10 for Jiamusi aphids and Suihua aphids, Spirotetramat dose 10 for Changchun aphids and Qiqihqar aphids and Omethoate dose 10 for Zhalan aphids (Table 6).

Acknowledgments

This research was mainly supported by the China Agriculture Research System (CARS-04).

Conflict of interest

The authors declare that they have no conflict of

interest.

References

- Abd-Ella A.** 2015. Effect of several insecticides on pomegranate aphid, *Aphis punicae* (Passerini) (Homoptera: Aphididae) and its predators used field conditions. *EPPO Bulletin* **45**(1).
- APPPC. Asia and Pacific Plant Protection Commission** 1987. Insect pests of economic significance affecting major crops of the countries in Asia and the Pacific region. Tech. Doc. **135**. Bangkok, Thailand: Regional Food and Agriculture Organization. Office for Asia and the Pacific.
- Baute T.** 2001. Soybean aphids found in Ontario. *Field Crops (Crop Technology)*, **6**(11).
- Beckendorf EA, Catangui MA, Riedell WE,** 2008. Soybean aphid feeding injury and soybean yield, yield components, and seed composition. *Agronomy Journal* **100**, 237–246.
- Bishop A, McKenzie H, Barchia I, Spohr L,** 1998. Efficacy of insecticides against the lucerne flea, *Sminthurus viridis*(L.) (Collembola: Sminthuridae), and other arthropods in lucerne. *Australian Journal of Entomology* **37**(1).
- CAB International.** 2001. Crop protection compendium. CD-ROM.
- Chen QH, Yu SY.** 1988. Aphids and control. Shanghai Science and Technology Press, Shanghai, China.
- DiFonzo CD.** 2009. Tiny terrors: the soybean aphid. *American Entomologist*. **55**.
- Faheem M, Sajjad A, Shafique RM, Rehman A, Aslam MN.** 2016. Field evaluation of different insecticides against wheat aphids and their natural enemies in Pakistan. *Asian journal of Agriculture Biology* **4**(4).

- Fernandez-Cornejo J, Jans S.** 1999. Pest management in U.S. Agriculture. Resource Economics Division, U.S. Department of Agriculture Economics Research Service, Agriculture Handbook No. 717.
- Fletcher MJ, Desborough P.** 2002. The soybean aphid, *Aphis glycines*, present in Australia.
- Guo JQ, Zhang MH.** 1989. Study on the important vectors of soybean mosaic virus and their transmission efficiency. *Journl of Soybean Sciences*. **8**, 55D63.
- He FG, Liu XD, Yan FY, Wang YQ.** 1995. Soybean resistance to the soybean aphid. *Liaoning. Journal of Agricultural Sciences* **4**, 30 D34.
- Heinrichs EA, Fabellar LT, Basilio RP, Wen TC, Medrano F.** 1984. Susceptibility of rice plant hoppers, *Nilaparvata lugens* and *Sogatella furcifera* to insecticides as influenced by level of resistance in the host plant. *Journal of Environmental Entomology*. **13**, 455–458.
- Hodgson EW, Van Nostrand G, O'Neal ME.** 2010. 2010 yellow book: report of insecticide evaluation for soybean aphid. Department of Entomology, Iowa State University, Publication 287–10.
- Hodgson EW, McCornack BP, Tilmon K, Knodel JJ.** 2012. Management recommendations for soybean aphid (Hemiptera: Aphididae) in the United States. *Journal of Integrated Pest Management* **3(1)**.
- Hurley T, Mitchell P.** 2017. Value of neonicotinoid seed treatments to US soybean farmers. *Journal of Pest Management Science* **73**, 102–112.
- Li WM, Pu ZQ.** 1991. Population dynamics of aphids and epidemics of soybean mosaic virus in summer sown soybean in Nanjing. *Journal of Acta Phytotaxonomica Sinica* **18**, 123D126.
- Lin CL, Xun ZS, Li LT, Zhang HK, Zhang GX, Wang YP.** 1994. Population dynamics and control stage of the soybean aphid in Jining prefecture, China Shandong. *Journal of Agricultural Sciences* **4**, 44.
- Lin CL, Xun ZS, Li LT, Wang YP, Zhang GX.** 1992. Control threshold of the soybean aphid in the field. *Journal of Soybean Science* **11**, 318 D321.
- Luo RW, Shang YF, Yang CL, Zhao JH, Li CS.** 1991. Study on the epidemiology and prediction of soybean mosaic virus. *Journal of Acta Phytotaxonomica Sinica* **18**, 267D271.
- Magalhaes LC, Hunt TE, Siegfried BD.** 2008. Development of methods to evaluate susceptibility of soybean aphid to imidacloprid and thiamethoxam at lethal and sublethal concentrations. *Entomologia experimentalis et applicata* **128**, 330 D336.
- Qu YX, Ma ZQ, Shan DA, Gao XH, Wang QS.** 1987. Effects of insecticides on population of destructive insects and their natural enemies in the soybean field. *Journal of Plant Protection* **13**, 4D6.
- Ragsdale DW, McCornack BP, Venette RC, Potter BD, MacRae IV, Hodgson EW, O'Neal ME, Johnson KD, O'Neil RJ, DiFonzo CD, Hunt TE, Glogoza PA, Cullen EM.** 2007. Economic threshold for soybean aphid (Hemiptera:Aphididae). *Journal of Economic Entomology* **100**, 1258–1267.
- Royer TA, Elliott NC, Giles KL, Kindler SD.** 2011. Field efficacy of wintertime insecticide applications against greenbugs, *Schizaphis graminum* (Rondani) (Hemiptera: Aphididae) on winter wheat (*Triticum aestivum* L.). *Journal of crop protection*. **30**.
- Shafique MA, Ahmed KS, Haider N, Khan RP, Majeed MZ.** 2016. Field evaluation of different insecticides against wheat aphid (*Schizaphis graminum* Rondani) and comparative yield assessment for different wheat cultivars. *Acad.*

Journal of Entomology and Zoology **9**(1).

Sun B, Liang SB, Zhao WX. 2000. Outbreak of the soybean aphid in Suihua prefecture in 1998 and its control methods. *Journal of Soybean Bully* **1**, 5.

Tabadkani SM, Khoobdel M, Tavakoli HR. 2017. Host resistance enhances susceptibility of *Callosobruchus maculatus* (Coleoptera: Chrysomelidae) to herbal extract of *Echinophora platyloba*. *Journal of Entomological Research* **47**, 28–34.

Takahashi S, Inaizumi M, Kawakami K. 1993. Life cycle of the soybean aphid *Aphis glycines* Matsumura, in Japan. *Journal of Applied Entomology and Zoology* **37**, 207–212.

U.S. Department of Agriculture [USDA]. 2002. Soybean Aphid Watch.

Voegtlin DJ, Halbert SE, Qiao G. 2004. A guide to separating *Aphis glycines* Matsumura and morphologically similar species that share its hosts. *Annals of the Entomological Society of America* **97**, 227–232.

Wang CL, Xiang NY, Zhang GS, Zhu HF. 1962. Studies on the soybean aphid, *Aphis glycines*

Matsumura. *Journal of Acta Entomologica Sinica* **11**, 31–44.

Wang SY, Bao XZ, Sun YJ, Chen RL, Zhai BP. 1996. Effect of population dynamics of the soybean aphid (*Aphis glycines*) on soybean growth and yield. *Journal of Soybean Science* **15**, 243–247.

Wu XB, Ni WJ, Liu PJ. 1999. Occurrence and control of the soybean aphid, *Aphis glycines* Matsumura. *Chinese Journal of Biological Control* **6**, 20.

Youhui Gong, Xueyan Shi, Nicolas Desneux, Xiwu Gao. 2016. Effects of spirotetramat treatments on fecundity and carboxylesterase expression of *Aphis gossypii* Glover. *Ecotoxicology* **25**(4).

Yue DR, Guo SG, Shan YL. 1989. Resistance of wild soybean *Glycine soja* to *Aphis glycines*. I. Screening for resistant varieties. *Journal of Jilin Agricultural Sciences* **3**, 15–19.

Zhang HJ. 1982. Relationship between population fluctuations of the soybean aphid *alatae* and epidemic level of soybean virus. *Chinese Journal of Oil Crop Sciences* **2**, 59–61.