



RESEARCH PAPER

OPEN ACCESS

Foliar applied iron and zinc improves the growth, physiological and yield related traits of wheat under drought

Rashid Iqbal^{1,*}, Muhammad Aown Sammar Raza^{1,*}, Muhammad Farrukh Saleem², Imran Haider Khan³, Muhammad Saqlain Zaheer^{1,4}, Salman Ahmad¹, Imran Haider¹, Muhammad Usman Aslam¹, Monika Toleikiene⁵

¹Department of Agronomy, University College of Agriculture and Environmental Sciences, The Islamia University of Bahawalpur, Pakistan

²University of Agriculture Faisalabad, Department of Agronomy, Faisalabad, Pakistan

³Department of Agronomy, Nanjing Agricultural University, China

⁴School of Agriculture, Food and Wine, The University of Adelaide, Australia

⁵Lithuanian Center for Agriculture and Forestry (LAMMC), Lithuania

Key words: Drought stress, Iron Spray, Zinc spray, Wheat, photosynthetic rate & water related traits.

<http://dx.doi.org/10.12692/ijb/14.3.361-372>

Article published on March 27, 2019

Abstract

Water is the basic unit for sustaining healthy plant life. Limited or unavailability of water leads to ultimate plant death. Drought severely affects the yield of wheat, especially when it occurs at the critical growth stages like tillering and anthesis. Therefore, attempts have been and being made to develop measures to combat the drought effects in wheat. This study investigated the sole and/or integrated use of Iron (Fe) and Zinc (Zn) spray on growth and yield of Wheat under drought stress. Drought was created at the tillering and anthesis stages by withholding the irrigation. At both these stages, different doses of Zn and Fe were applied alone and/or in combination. Results revealed that drought stress significantly ($P < 0.05$) reduced the 1000-grain weight (30 vs. 42g), number of grains per spike (27 vs. 62), spike length (11 vs. 17cm), number of spikelets per spike (22 vs. 9) physiological (leaf chlorophyll content, stomatal conductance & photosynthetic rate) and water related traits. Combine application of Fe and Zn to wheat under drought significantly increased the 1000 grain weight (40g), seed yield (5350 kg/ha), relative water content, leaf osmotic potential, leaf water potential, protein content (16%) and spike length (16cm). The Zn was more effective in mitigating the ill impacts of drought as compared to the Fe alone. While, combine spray of both Zn and Fe, gave best results as compared to the sole application of either nutrient. Results of this study lead us to conclude that both Zn and Fe have the ability to compensate the detrimental effects caused by drought stress and their combine use can be more useful as compared to their sole usage.

* Corresponding Author: Muhammad Aown Sammar Raza ✉ aown.sammar@iub.edu.pk

Introduction

Wheat (*Triticum aestivum* L.) is considered as one of the most important cereal crop and staple food in many countries of the world including Pakistan. It is used to make roti, nans, bread, cookies, pasta and other products. It is the cheapest source of human calories and protein content in the average diet (Lev-Yadun *et al.*, 2000). Among the all cereal crops, wheat is the most favorite food for human being and as feed for animals also (Vaughan and Judd, 2003).

Wheat grains contain 9.4-14 g/100g protein, 1.8-2.3 g/100g fiber, 1.2-2.5 g fat and have 69.1-75.4 g/100g available carbohydrates (Ken, 2004). Wheat provides about 55 percent of the carbohydrates and 20% of the food calories as consumed globally (Breiman and Graur, 1995). Wheat is grown under different variety of soil and climatic conditions in Pakistan. From the total acreage, about one third falls under rain fed areas where annual amount and intensity of rainfall is very low (Khanzada *et al.*, 2001). Drought stress makes threats to the agricultural production in the majority of the areas of world (Abolhasani and Saeidi, 2004), as 20,000,000 km² of the lands throughout the world lies in semi-arid regions (Alyari and Shekari, 2000).

Drought causes a drastic effect on plant productions each year in the different regions of the world. There are many reasons for the low availability of water to the plants not only due to high salinity conditions and high temperature of the surroundings but also due to the deviation in the normal rainfall pattern of an area (Baybordi, 2004; Soriano *et al.*, 2004). Drought stress is characterized by reduction of water content, turgor loss, very low leaf water potential, leaf stomata opening and closing, reduction in cell enlargement and decline in growth occurs. In the harsh conditions of water deficiency photosynthesis process in plants may stop which leads to the interruption of metabolism and at last the death of plant (Jaleel *et al.*, 2008). Shortage of water in soil also causes very slow movement of micro-nutrients like as Iron and Zinc in soil and plants. This disturbs the growth and many physiological processes of plants (Chimenti *et al.*, 2002).

The chlorophyll production and quantity of abscisic acid are affected by the utilization of Zn and Fe (Jiang and Huang, 2002). The increase in the quantity of chlorophyll ultimately enhances yield during the light reaction of photosynthesis (Conversion of carbon dioxide and water into glucose). Starch accumulation in the grain can be suppressed due to the limited availability of water during the process of assimilates translocation at the time of grain filling (Yang *et al.*, 2004). In the same way, grain weight and spike length are also adversely affected due to water deficit condition (Wang *et al.*, 2005). Less grain weight under drought conditions is the main reason of reduced seed yield of wheat (Baybordi, 2004).

Many management practices are used to improve the drought tolerance in wheat like stubble or residual mulches (Ahmad *et al.*, 2015), use of osmolytes (Raza *et al.*, 2012a) or solutes like spray of potassium (Raza *et al.*, 2012b), gibberelic acid and amino acid application and use of micronutrients (Iron, Zinc, Copper, etc). Studies on micronutrients have proved that they can be used efficiently in crop plants for mitigating the impact of different kinds of stresses like water stress. Iron and Zinc are among those micronutrients. Both the micro and macro nutrients are importance for plant nutrition and low quantity of any nutrient can badly effect the production of crop (Murphy *et al.*, 2008). Unbalanced application of phosphorus and nitrogen produce the low quality of grain and these fertilizers have also reduced the micronutrients uptake by plants (Maralian, 2007-2008). So, there is a need to check the effects of micronutrients application on plants. Micronutrients can also increase dry matter production, total straw yield, and grain yield of wheat (Asad and Rafique, 2002).

Many researchers have reviewed Zinc deficiency in the soils of Pakistan (Hadi *et al.*, 1997). Amongst the micronutrients, Zinc scarcity takes place both in humans and crop plants (Hotz and Brown, 2004). Latest reports indicated that almost 5000, 000 children die during 5 years of age per annum due to insufficiency of low Zn and Fe in their diet (Black *et*

al, 2008). Zinc sulphate fertilizer is the best one among the fertilizers needed by plants due to its performance on stomatal adjusting according to the changing environment and uptake of nutrients and ions by plants. So, there is need to pay attention during application of such kind of fertilizers either water is sufficient or inadequate (Karam *et al.*, 2007; Babaeian *et al.*, 2010).

Unbalanced use of synthetic fertilizers and prolonged drought are the main factors for low amount of Iron and Zinc in the soil solution. Seed yield and protein percentage in the plants increases with the application of Iron (Yahyavi *et al.*, 2004). Seed yield improved due to the utilization of Iron, which affects photosynthesis in plants, and ultimately there will enhance of effectiveness of dry matter substances and reallocation of photosynthates into the economic portion of wheat. The growers in the arid areas face the problem of Iron deficiency due to high pH of soils. Pakistani soils are high in pH which results reverse effects of Iron and Zinc with the other nutrients and consequently in the presence of other plant nutrients Zn and Fe availability to plants is less. Under drought stress conditions, we believe that the yield losses can be minimized by the sufficient supply of Zn and Fe. The objective of present work was to study the possible role of Zn and Fe applied on wheat foliage at different growth stages under drought, in mitigation of stress in terms of physiological components and nutrients uptake.

Materials and methods

To explore the objective as outlined above, a field experiment on clay-loam soil was conducted at agronomic research area, University College of Agriculture and Environmental Sciences, The Islamia University of Bahawalpur to investigate the effect of foliar application of iron and zinc on wheat growth, physiology and yield under drought stress. Soil analysis carried out prior to sowing revealed EC = 2.04 dsm⁻¹, pH = 8.6, O.C(%) = 0.79, soil P₂O₅ = 5 ppm, K₂O = 115 ppm, N = 0.109, field capacity = 21% W/W, wilting point = 10% W/W and the volume weight of the soil was 1.22 g.cm³.

The experiment was laid out in randomized block design with four replications. Treatments included were T₀=Control, T₁=Drought, T₂=Zn (150 g ha⁻¹), T₃=Fe (150 g ha⁻¹), T₄=Zn+Fe (150+150g ha⁻¹), T₅=Zn+Fe (200+150 g ha⁻¹) and T₆= Zn+Fe (150+200 g ha⁻¹).The foliar spray of Fe and Zn was done at tillering and anthesis after giving drought at both stages. Zinc was applied as zinc sulphate and Iron as Iron oxide both were applied through foliar application and at a rate of 150g/ha. Additionally, the wheat crop received P and N at a rate of 112 kg ha⁻¹ o and 150 kg ha⁻¹, respectively. Total amount of P was mixed in soil while soil preparation, whereas N was applied in two splits: at the time of soil preparation and at heading stage of wheat crop. Wheat seeds were sown in each plot with size of 5m x 3m with hand drill having 22.5 cm row to row distance. Leaf chlorophyll contents were measured by chlorophyll meter. Stomatal resistance/conductance measurements was made with an automatic porometer MK-3 (Delta-T Devices, Burwell Cambridge, England) Hertford, Herts, England) while Photosynthetic rate was recorded by using the an infrared gas analyzer (Li-COR-LI 6250). Relative water content (RWC) was calculated by means of the following formulas, RWC (%) = FW- DW/ TW- DW X 100.

For recoding of leaf water potential (-MPa), an entirely stretched youngest leaf (fourth from the top) was cut out from every plant and calculations were made with a water potential apparatus (Chas W. Cook Div., England). To compute the leaf osmotic potential (-MPa), a fraction of the leaf used for water potential purpose will be frozen for two weeks. After that, frozen leaf thawed and crushed using metal rod to take out the frozen sap. After centrifugation (8000 x g) for 4 minutes, the sap will be used for osmotic potential determination in a Vapor Pressure Osmometer (Wescor 5520, Logan, USA).

At maturity when moisture content in the seed was about 14%, area of one square meter in each plot was randomly selected using a quadrant. Thereafter, the plants in this area were manually harvested to estimate, grain yield biological yield, no. of spikes per

surface unit, number of grains per spike, number of spikelets per spike, length of the spike, height of the plant and stem, and thousands kernels.

The grains from each treatment were oven-dried at 70 °C and ground to pass a 1mm sieve. Dried ground form of wheat grain (0.1 g) was taken in digestion tubes and 5ml of concentrated sulphuric acid (H₂SO₄) was added to each digestion tube and incubated overnight at 25°C. Then 1 ml of hydrogen peroxide (H₂O₂) (34%) was poured into the sides of the digestion tube, ported the tubes in a digestion block and heated at 350°C, until fumes were produced. They were continued to heat for half an hour. The digestion tubes were removed from the block and cooled down. Then 1 ml of H₂O₂ was added and the tubes were kept back into the digestion block. The above step was repeated until the cooled digested material was colorless. Thereafter about 0.1 g was

used for the determination of total N content by Kjeldahal's method and later converted into protein content by multiplying N content with a factor of 5.70 (ISO, 2009). The K⁺ in the digests was determined with the help of flame photometer and phosphorus was determined by spectrophotometer at 420 nm using standard curve method (Lindner, 1944). Collected data regarding all the parameters was analyzed by using Fisher's analysis of variance technique and LSD test at 5% probability was used to compare the differences among treatments (Steel *et al.*, 1997).

Results

Growth & Physiological Related Parameters

Results revealed that no. of plants m⁻², plant height and number of fertile tillers m⁻² were not significantly affected under drought stress by foliar application of Iron (Fe) and Zinc (Zn) (Table 1).

Table 1. Variance analysis and means compares of drought stress and effects of foliar application of Iron and Zinc on wheat (*Triticum aestivum* L.) growth components.

| Treatments | Plant Population (m ⁻²) | No. of fertile tillers(m ⁻²) | Plant height (cm) | Spike length (cm) | Number of spikelets per spike |
|--|-------------------------------------|--|-------------------|-------------------|-------------------------------|
| T ₀ =Control | 210 a | 421a | 98 a | 16.83 a | 21.67 a |
| T ₁ =Drought | 198 b | 396 b | 86 b | 10.50 g | 9.33 g |
| T ₂ =Zn(150 g ha ⁻¹) as ZnSO ₄ +Drought | 200 b | 403 b | 93ab | 12.50 e | 13.33 e |
| T ₃ =Fe(150g ha ⁻¹) as Fe ₂ O ₃ + Drought | 206 ab | 415 ab | 93 ab | 11.58 f | 11.33 f |
| T ₄ =Zn+Fe (150+150g ha ⁻¹) +Drought | 205 ab | 413 ab | 91 ab | 13.67 d | 15.33 d |
| T ₅ =Zn+Fe (200+150g ha ⁻¹) +Drought | 204 ab | 413 ab | 95 a | 16.00 b | 19.67 b |
| T ₆ =Zn+Fe (150+200g ha ⁻¹) +Drought | 200 ab | 418 ab | 93 ab | 15.00 c | 17.33 c |

Significant results at 0.05 % probability.

Spike length was significantly affected by foliar applied micronutrient treatments (Table 1). Highest spike length was obtained in control treatment (T₀) and minimum in drought stressed treatment (T₁).

Treatment in which zinc (T₂) and iron (T₃) were applied solely produced more lengthy spikes than T₁. Combine application of micronutrient (T₄, T₅ and T₆) produced more lengthy spikes as compared to their sole application (T₂ and T₃). Among the combine micronutrient treatments, T₅ produced lengthiest spike.

Foliar applied micronutrient showed significant effects on number of spikelets per spike (Table 1). Maximum number of spikelets per spike (21.67) was obtained in control treatment (T₀) and minimum in drought stressed treatment (T₁). Treatment in which zinc (T₂) and iron (T₃) were applied solely produced more number of spikelets per spike than T₁. Combine application of micronutrient (T₄, T₅ and T₆) produced more number of spikelets per spike as compared to their sole application (T₂ and T₃). Among the combine micronutrient treatments, T₅ produced maximum number of spikelets per spike (19.67).

Table 2. Variance analysis and means compares of drought stress and effects of foliar application of Iron and Zinc on wheat (*Triticum aestivum* L.) physiological components.

| Treatments | Leaf chlorophyll contents (%) | Stomatal conductance (mmolm ⁻² s ⁻¹) | Photosynthetic rate (μmolm ⁻² s ⁻¹) | Leaf relative water contents (%) | Leaf osmotic potential (MPa) | Leaf water potential (MPa) |
|--|-------------------------------|---|--|----------------------------------|------------------------------|----------------------------|
| T ₀ =Control | 30.00 a | 360a | 15.50 a | 84.50 a | 1.80 a | 1.12 a |
| T ₁ =Drought | 14.00 g | 220 g | 7.03 g | 40.00 g | 1.11 g | 0.22 g |
| T ₂ =Zn(150 g ha ⁻¹) as ZnSO ₄ +Drought | 20.00 e | 290 e | 12.03e | 65.00 e | 1.42 e | 0.65 e |
| T ₃ =Fe(150g ha ⁻¹) as Fe ₂ O ₃ + Drought | 17.50f | 275 f | 11.80 f | 60.00 f | 1.35 f | 0.58 f |
| T ₄ =Zn+Fe (150+150g ha ⁻¹) +Drought | 22.13 d | 320 d | 12.80 d | 70.80 d | 1.62 d | 0.85 d |
| T ₅ =Zn+Fe (200+150g ha ⁻¹) +Drought | 27.50 b | 338 b | 14.20 b | 78.50 b | 1.72 b | 0.98 b |
| T ₆ =Zn+Fe (150+200g ha ⁻¹) +Drought | 23.50 c | 330 c | 13.40 c | 75.50 c | 1.68 c | 0.92 c |

Significant results at 0.05 % probability.

A significant effect of foliar applied micronutrient was observed on leaf chlorophyll contents (Table 2). More leaf chlorophyll content was obtained in control treatment (T₀) and minimum in drought stressed treatment (T₁). Solely applied zinc (T₂) and iron (T₃) produced more leaf chlorophyll content (%) than T₁. Combine application of micronutrient (T₄, T₅ and T₆) gave more leaf chlorophyll content than their sole application (T₂ and T₃). Among the combine micronutrient treatments, T₅ produced maximum leaf chlorophyll content.

Stomatal conductance (mmolm⁻²s⁻¹) was significantly affected by foliar applied micronutrient (Table 2). The higher stomatal conductance (mmolm⁻²s⁻¹) was obtained in control treatment (T₀) and minimum in T₁. Sole application of zinc (T₂) and iron (T₃) resulted in more stomatal conductance (mmolm⁻²s⁻¹) than T₁. Combine application of micronutrient (T₄, T₅ and T₆) produced more stomatal conductance (mmolm⁻²s⁻¹) as compared to their sole application (T₂ and T₃). Among the combine micronutrient treatments, T₅ produced highest stomatal conductance (mmolm⁻²s⁻¹). Table.2 indicates that foliar applied micronutrient had statistically significant effect on photosynthetic rate of wheat. The maximum photosynthetic rate (μmolm⁻²s⁻¹) was observed in control treatment (T₀) and minimum in T₁. Sole application of zinc (T₂) and iron (T₃) resulted in higher photosynthetic rate (μmolm⁻²s⁻¹) than T₁. Combine application of

micronutrient (T₄, T₅ and T₆) showed more photosynthetic rate (μmolm⁻²s⁻¹) as compared to their sole application (T₂ and T₃). Among the combine micronutrient treatments, T₅ produced highest photosynthetic rate (μmolm⁻²s⁻¹).

A significant effect of foliar applied micronutrient was observed on all water related traits of wheat i.e. leaf relative water content (%), leaf osmotic and leaf water potential (MPa) (Table 2). Higher water related parameters were obtained in control treatment (T₀) and minimum in drought stressed treatment (T₁). Solely applied zinc (T₂) and iron (T₃) showed more water related traits than T₁. Combine application of micronutrient (T₄, T₅ and T₆) gave more values of relative water content (RWC), leaf osmotic potential and leaf water potential than their sole application (T₂ and T₃). Amongst the combine micronutrient treatments, T₅ showed higher rates of all water related traits of wheat.

Yield Related Parameters

Results showed that contents of phosphorus and potassium in wheat grain were not significantly affected under drought stress by foliar application of Iron (Fe) and Zinc (Zn) (Table 3).

A significant effect of foliar applied micronutrient was observed on number of grains per spike (Table 3). The highest number of grains per spike was obtained in

control treatment (T_0) and minimum in drought stressed treatment (T_1). Solely applied zinc (T_2) and iron (T_3) produced more number of grains per spike than T_1 . Combine application of micronutrient (T_4 , T_5 and T_6) gave more number of grains per spike than their sole application (T_2 and T_3). Among the combine micronutrient treatments, T_5 produced maximum number of grains per spike. 1000-grain weight was significantly affected by foliar applied micronutrient (Table 3). The maximum 1000-grain weight was

obtained in control treatment (T_0) and minimum in T_1 . Sole application of zinc (T_2) and iron (T_3) resulted in more 1000-grain weight than T_1 . Combine application of micronutrient (T_4 , T_5 and T_6) produced maximum 1000-grain weight as compared to their sole application (T_2 and T_3). Among the combine micronutrient treatments, T_5 produced highest 1000-grain weight. Foliar applied micronutrient also showed significant effects on grain yield of wheat (Table 3).

Table 3. Variance analysis and means compares of drought stress and effects of foliar application of Iron and Zinc on wheat (*Triticum aestivum* L.) yield and yield components.

| Treatments | No. of grains per spike | 1000 grain weight (g) | Grain Yield (kg/ha) | Phosphorus content (%) | Potassium content (%) | Protein content (%) |
|--|-------------------------|-----------------------|---------------------|------------------------|-----------------------|---------------------|
| T_0 = Control | 62.33 a | 42.33 a | 5500.00 a | 0.38 a | 2.3 d | 11.55 g |
| T_1 = Drought | 27.66 g | 30.33 g | 3916.67 g | 0.38 a | 2.7 ab | 12.57 f |
| T_2 = Zn(150g ha ⁻¹) as ZnSO ₄ +Drought | 39.66 e | 33.66 e | 4350.00 e | 0.39 a | 2.2 d | 13.81 d |
| T_3 =Fe(150g ha ⁻¹) as Fe ₂ O ₃ +Drought | 33.66 f | 32.00 f | 4016.67 f | 0.41 a | 2.3 cd | 13.14 e |
| T_4 =Zn+Fe(150+150g ha ⁻¹) +Drought | 44.66 d | 35.66 d | 4716.67 d | 0.42 a | 2.4 bcd | 14.32 c |
| T_5 =Zn+Fe(200+150g ha ⁻¹) +Drought | 57.33 b | 40.33 b | 5350.00 b | 0.43 a | 2.7 abc | 16.05 a |
| T_6 =Zn+Fe(150+200g ha ⁻¹) +Drought | 51.00 c | 37.66 c | 5016.67 c | 0.39 a | 2.9 a | 14.91 b |

Significant results at 0.05 % probability.

The maximum grain yield (5500.00 kg/ha) was obtained in T_0 and minimum (3916.67 kg/ha) in T_1 . Treatment in which zinc (T_2) and iron (T_3) were applied solely produced more grain yield than T_1 . Combine application of micronutrient (T_4 , T_5 and T_6) yielded more grain than their sole application (T_2 and T_3). Among the combine micronutrient treatments, maximum grain yield (5350.00 kg/ha) was obtained from T_5 .

Grain protein content was significantly affected by foliar applied micronutrient (Table 3). Maximum grain protein content was obtained in T_5 and minimum in T_0 .

Treatment in which zinc (T_2) and iron (T_3) were applied solely produced more grain protein content than T_1 . Combine application of micronutrient (T_4 , T_5 and T_6) produced more grain protein content as compared to their sole application. Among the combine micronutrient treatments, T_5 produced highest grain protein content.

Discussion

Water deficit stress affects the growth, physiological, yield and quality parameters of wheat. Application of micronutrients (Fe and Zn) either alone or in combination was reported to be an effective technique to minimize the adverse effects of drought. Spike length of wheat adds enormous share in ultimate economic yield because it has considerable influence on number of spikelets, grain size and number of grains per spike. Drought affected wheat plants produced smaller spikes as compare to the proper irrigated plants (Taheri *et al.*, 2011). It is due to less turgor pressure, leads to less cell division and less growth (Jaleel *et al.*, 2009). Abbas *et al.* (2009) found that foliar applied Zinc (Zn) has a positive effect on spike length of wheat under water deficit conditions due to the role of Zn is in auxin production which is involve in cell division and cell elongation. Fe also has a significant effect on spike length of wheat, as Fe is helpful in chlorophyll formation which results in more photosynthesis and more growth. Combine application of Zn and Fe can result in the formation of

longer spikes (Kholdebarin and Islamzadeh, 2002; Monjezi *et al.*, 2013; Jam *et al.*, 2012). It might be due to synergistic effect of both nutrients (Gaffar *et al.*, 2011).

Chlorophyll concentration in plant leaves display its capacity to make photosynthates (Zhang *et al.*, 2007). Process of photosynthesis is badly affected by concentration of chlorophyll. This concentration is mostly lower under drought stress conditions due to the reduction in activity of chlorophyllase enzyme which is mainly involved in the formation of chlorophyll (Ashraf *et al.*, 2013). Simultaneous application of both iron and zinc produced more concentration of chlorophyll in wheat crop as compared to drought stress. Similar results were also proposed by Khakwani *et al.* (2013) in fully irrigated crop (more chlorophyll concentration) as compared to water deficit conditions. Iron is directly involved in the formation of chlorophyll.

Stomatal conductance is an imperative measurement of plant responses under drought or low availability of water (Baloch *et al.*, 2012). Synergistic effect of both iron and zinc is helpful for crop plants in water stress conditions (Gaffar *et al.*, 2011). These nutrients are involved in the controlling of stomatal aperture by producing auxin in this way a resistance may be created under drought stress conditions. Process of photosynthesis computes the overall growth and yield of crop plants. Under severe water conditions and low relative water content of leaves there is very slow rate of photosynthesis (Alkhaldi *et al.*, 2012). Stomatal conductance is directly related to the formation of glucose in photosynthesis.

Relative water content of plant leaves is the indication of presence of water in tissues of plant leaves. Computation of drought is also carried out through it (Sanchez-Blanco *et al.*, 2002). Control treatment showed higher values of relative water content due to more leaf area and cell volume in relation to drought (Stikic *et al.*, 2003). Combine use of zinc and iron showed better results than their sole application and drought. Low RWC is also attributed with the root

systems of various cultivars and their capability to absorb the soil moisture in water deficit conditions (Khakwani *et al.*, 2013). Leaf water and osmotic potentials have positive influences on the overall growth and physiology of crop plants. Leaf water potential shows the potential energy of water in leaf tissues. Simultaneously applied micronutrients improved the water related traits of wheat due to their synergistic effects as zinc is directly involved in the production of auxins in plants and iron is the main functional element in the formation of chlorophyll. Changing water potential of various varieties is might be due to the alteration in osmotic pressure of crop plants (Siddique *et al.*, 2000).

Potential of wheat production is estimated by the number of spikelets per spike. Number of spikelets per spike of wheat add huge share in final economic yield. Less number of spikelets per spike under water deficit conditions is mainly due to less spikelets primordial formation during tillering stage or may be due to death of floret at the terminal and basal ends of the spike (Maqbool *et al.*, 2015). Abbas *et al.* (2009) noted significant effect of different levels of Zn on number of spikelets per spike in wheat under water deficit conditions. Zn is more efficient on the formation of spikelets and/or the increase in the number of spikelets per spike due to role of Zn in the production of growth promoting hormone auxin and its role in many metabolic activities (Hafeez *et al.*, 2013). Foliar applied Fe also increased the number of spikelet per spike in drought affected wheat as compared to untreated plants due to more photosynthesis (Abbas *et al.*, 2009). However combined application of Fe and Zn produced more number of spikelets per spike, might be due to synergistic effects of both nutrients (Zeidan *et al.*, 2010).

Number of grains per spike is the main factor which particularly contributes to the economic yield of wheat. The number of grains per spike is the major factor which affects the grain yield. Under water deficit stress number of grains decreased mainly due to the dehydration of pollen grain (Khanzada *et al.*,

2001). Micronutrients had a significant effect on the number of grains per spike (Jam *et al.*, 2012). Foliar application of micronutrients enhanced the number of wheat grains per spike (Khan *et al.*, 2010). Zeidan *et al.* (2010) reported that micronutrients Fe and Zn alone application in different treatments increase the number of wheat grains per spike than the control treatment through positively affecting the fertility of flowers (Monjezi *et al.*, 2013).

1000-grain weight indicates the degree of the formation of grain in wheat crop. Treatment in which proper irrigation was applied produced more 1000-grain weight than drought affected plants. More 1000-grain weight with proper irrigations is due to more translocation of assimilates towards grain because of availability of sufficient amount of water in root zone (Maqbool *et al.*, 2015). Alone application of either Fe or Zn give higher 1000-grain weight than without application of these nutrients (Zeidan *et al.*, 2010). The increase in 1000-grain weight of wheat by the simultaneous application of Fe and Zn has been reported by Ziaieian and Malakoti (2002) and El-Majid *et al.* (2000). The increase in grain weight was resulted from the positive effects of these micronutrients on post-anthesis photosynthesis and the allocation of dry matter to grains which finally led to the increase in 1000-grain weight. Similar results were reported by Monjezi *et al.* (2013) and Jam *et al.* (2012).

Seed yield is the result of various yield contributing parameters like spike length, number of spikelets per spike, number of grains per spike, and grain weight. Drought stress decreases the grain yield of wheat. Less grain yield under water deficit stress is due to less translocation of photosynthates from the leaves and as drought stress faster the maturation process. More seed yield of wheat with foliar application of Fe is mainly due to its role in enzymatic reactions and photosynthesis, which ultimately result in higher seed yield (Habib, 2009). More seed yield with application of Zn is due to its role in production of protein, carbohydrates, cell metabolism, protection of cell organelles and other processes linked to make the

plants well-matched with water shortage conditions (Yari *et al.*, 2005). These results are in lined with the findings of Yilmaz *et al.* (1997), El-Majid *et al.* (2000) and Zeidan *et al.* (2010). Synergistic effect of both Zn and Fe results more grain yield as compared to their sole application (Gaffar *et al.*, 2011).

Drought stress at both tillering and anthesis stages shortened the grain filling stage of wheat, which in turn, increased the grain protein as compared to control treatment due to the increased accumulation of hydrocarbons (Monjezi *et al.*, 2013). Drought affected plants treated with Zn has more grain protein contents due to the positive effect of Zn on important plant processes, such as nitrogen metabolism and nitrogen uptake by the plants (Potarzycki and Grzebisz, 2009). Drought affected plants treated with Fe also has more grain protein contents than control treatment. Seadh *et al.* (2009) reported that foliar applied Fe @ 500 ppm on wheat enhanced grain protein content over untreated wheat plants. Micronutrients (Iron and Zinc) either alone or in a combination had significant effect on grain protein content of wheat (Zeidan *et al.*, 2010). Similar results were found by Jiang *et al.* (2002) and Monjezi *et al.* (2013). It is obvious that the foliar application of Iron and Zinc increased the grain protein contents but at the same time reduced the concentration of phosphorus and potassium contents of wheat grain due to low uptake of these nutrients through roots (Zeidan *et al.*, 2010).

Conclusions

Foliar applied Iron and Zinc significantly improved the growth, physiology, yield and grain protein contents of wheat (*Triticum aestivum* L.) under drought stress conditions. Combined application of Zn + Fe (200+150 g ha⁻¹) under drought at tillering and anthesis performed better in improving growth, physiological, yield related parameters and quality of wheat.

Acknowledgements

To the University College of Agriculture and Environmental Sciences, the Islamia University of

Bahawalpur for provision of research facilities is gratefully acknowledged.

References

Abbas G, Khan MQ, Jamil M, Tahir M, Hussain F. 2009. Nutrient uptake, growth and yield of wheat (*Triticum aestivum* L.) as affected by zinc application rates. International Journal of Agriculture and Biology **11**, 389-396.

Abolhasani K, Saeidi G. 2004. Relationships between agronomic characteristic of safflower under water stress and control conditions. Iranian Journal of Field Crop Research **1**, 127-138.

Ahmad S, Raza MAS, Saleem MF, Zahra SS, Khan IH, Ali M, Shahid AM, Iqbal R, Zaheer, MS. 2015. Mulching strategies for weeds control and water conservation in cotton. Journal of Agriculture and Biological Science **8**, 299-306.

Alyari H, Shekari F. 2000. Oil seeds. Amidi Press. Tabriz. 182.

Alkhalidi A, Aldarir AN, Janat M, Wahbi MA, Arslan A. 2012. Effect of regulated deficit irrigation and partial root-zone drying on some quantitative indicators and the efficiency of adding nitrogen fertilizer to (*Zea mays* L.) by using n15 Isotope. American-Eurasian Journal of Agriculture and Environmental Sciences **12**, 1223-1235.

Asad A, Rafique R. 2002. Identification of micronutrients deficiency of wheat in the Peshawar Valley, Pakistan. Communication in Soil-Science and Plant Analysis **33**, 349-364.

Ashraf M, Shahbaz M, Ali Q. 2013. Drought-induced modulation in growth and mineral nutrients in rape (*Brassica napus* L.). Pakistan Journal of Botany **45**, 93-98.

Babaeian M, Heidari M, Ghanbari A. 2010. Effect of water stress and foliar micronutrient application on physiological characteristics and

nutrient uptake in sunflower (*Helianthus annuus* L.). Iranian Journal of Crop Science **12**, 311-391.

Baybordi A. 2004 - Effect of Fe, Mn, Zn and Cu on the quality of wheat under salinity stress. Journal of Water and Soil Science **17**, 140-150.

Baydar H, Erbas S. 2005. Influence of seed development and seed position on oil, fatty acids and total tocopherol contents in sunflower (*Helianthus annuus* L.). Turkish Journal of Agriculture **29**, 179-186.

Baloch MJ, Dunwell J, Khan NU, Khakwani AA, Dennett M, Jatoi WA. 2012. Profiling dehydrin gene sequence and physiological parameters in drought tolerant and susceptible spring wheat cultivars. Pakistan Journal of Botany **44**, 801-806.

Black RE, Lindsay HA, Bhutta ZA, Caulfield LE, DeOnnis M, Ezzati M, Mathers C, Rivera J. 2008. Maternal and child under nutrition: global and regional exposures and health consequences. Lancet **371**, 243-260.

Breiman A, Graur D. 1995. Wheat evolution. Israel Journal of Plant Sciences **43**, 85-98.

Ehdaie B, Alloush GA, Waines JG. 2008. Genotypic variation in linear rate of grain growth and contribution of stem reserves to grain yield in wheat. Field Crops Research **106**, 34-43.

El-Majid AA, Knany RE, El-Fotoh HGA. 2000. Effect of foliar application of some micronutrients on wheat yield and quality. Journal of Agricultural Sciences **130**, 301-313.

Gaffar A, Ehsanullah, Nadeem A, Sultan HK. 2011. Influence of zinc and iron on yield and quality of sugarcane planted under various trench spacing. Pakistan Journal of Agricultural Sciences **48**, 25-33.

Govt. of Pakistan. 2014. Economic Survey of Pakistan. Ministry of Food, Agriculture and Livestock,

Finance Division, Economic Advisors wing, Islamabad, Pakistan.

Habib M. 2009. Effect of foliar application of Zn and Fe on wheat yield and quality. *African Journal of Biotechnology* **8**, 6795-6798.

Hadi A, Taran SA, Azeem A, Saeed Z, Aslam, M. 1997. Wheat response to zinc application in rice growing area of Balochistan. *Pakistan Journal of Soil Science* **13**, 25-27.

Hafeez B, Khanif YM, Saleem M. 2013. Role of Zinc in plant nutrition, A-review, *American Journal of Experimental Agriculture* **3**, 374-391.

ISO. 2009. Food products – Determination of the total nitrogen content by combustion according to the Dumas principle and calculation of the crude protein content- Part 2: Cereals, pulses and milled cereal products. ISO/TS 16634-2: 1-25.

Jaleel CA, Manivannan PA, Wahid A, Farooq M, Al-Juburi HJ, Somasundaram R, Panneerselvam R. 2009. Drought stress in plants: a review on morphological characteristics and pigments composition. *International Journal of Agriculture and Biology* **11**, 100-105.

Jam E, Sajed K, Ebadi A, Farjaminejad R, Ghasempour F. 2012. Effect of Fe and Zn micronutrients spray on grain yield of autumn wheat in Ardabil Region, Iran. *Plant Ecophysiology* **3**, 101-107.

Jiang Y, Huang B. 2002. Protein alterations in tall fescue in response to drought stress and abscisic acid. *Journal of Crop Sciences* **42**, 202-207.

Karam F, Lahoud R, Masaad R. 2007. Evaporation, seed yield and water use efficiency of drip irrigated sun flower under full and deficit irrigation conditions. *Agricultural Water Management* **90**, 213- 235.

Ken P. 2004. Forms of micronutrient fertilizers. Sask. Agriculture & Food, Rigas Karamanos (Westco), Agric. and Agri-Food Canada: 1-10.

Khan MB, Farooq M, Shahnawaz MH, Shabir G. 2010. Foliar application of micronutrients improves the wheat yield and net economic return. *International Journal of Agriculture and Biology* **12**, 953-956.

Khanzada B, Ashraf MY, Ala SA, Alam SM, Shirazi MU, Ansari MU. 2001. Water relations in different Guar (*Cyamopsis tetra gonoloba L.*) genotypes under water stress. *Pakistan Journal of Botany* **33**, 279-287.

Khakwani AA, Dennett MD, Khan NU, Munir M, Baloch MJ, Latif A, Gul S. 2013. Stomatal and chlorophyll limitations of wheat cultivars subjected to water stress at booting and anthesis stages. *Pakistan Journal of Botany* **45**, 1925-1932.

Kholdebarin B, Islamzadeh T. 2002. Mineral feeding of higher plants. 1, Shiraz University Press, p, 500.

Lindner RC. 1944. Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant Physiology*.1, 76.

Lev-Yadun S, Gopher A, Abbo S. 2000. The Cradle of Agriculture Science **288**, 1602-1603.

Maqbool MM, Ali A, Haq T, Majeed MN, Lee DJ. 2015. Response of spring wheat (*Triticum aestivum L.*) to induced water stress at critical growth stages. *Sarhad Journal of Agriculture* **31**, 53-58.

Monjezi F, Vazin F, Hassan M. 2013. Effect of iron and zinc spray on wheat in drought stress, *Cercetari Agronomice in Moldova* 1, 153.

Murphy K, Hoagland L, Reeves P, Jones S. 2008. Effect of cultivar and soil characteristics on nutritional value in organic and conventional wheat. 16th IFOAM Organic World Conference in

- Cooperation with the International Federation of Organic Agriculture Movements (IFOAM) and the Consorzio Modena Bio., Modena, Italy, 18-20 June. 614-617.
- Potarzycki J, Grzebisz W.** 2009. Effect of zinc foliar application on grain yield of maize and its yielding components. *Plant, Soil and Environment* **55**, 519-527.
- Raza MAS, Saleem MF, Ashraf MY, Ali A, Asghar HN.** 2012a. Glycine betaine applied under drought improved the physiological efficiency of wheat (*Triticum aestivum* L.). *Plant, Soil and Environment* **31**, 67-71.
- Raza MAS, Saleem MF, Anjum SA, Khaliqand T, Wahid MA.** 2012b. Foliar application of potassium under water deficit conditions improved the growth and yield of wheat (*Triticum aestivum* L.) *Journal of Animal and Plant Science* **22**, 431-437.
- Sanchez-Blanco MJ, Rodriguez P, Morales MA, Ortuno MF, Torrecillas A.** 2002. Comparative growth and water relations of *Cistus albidus* and *Cistus monspeliensis* plants during water deficit conditions and recovery. *Plant Science* **162**, 107-113.
- Seadh SE, Abady EL, ElGhamry MI, Farouk AM.** 2009. Influence of micronutrients foliar application and nitrogen fertilization on wheat yield and quality of grain and seed. *Journal of Biological Sciences* **9**, 851-858.
- Siddique MRB, Hamid A, Islam M.S.** 2000. Drought stress effects on water relations of wheat. *Botanical Bulletin Academia Sinica* **41**, 35-39.
- Stikic R, Popovic S, Srdic M, Savic D, Jovanovic Z, Prokic LJ, Zdravkovic J.** 2003. Partial root drying (PRD): a new technique for growing plants that saves water and improves the quality of fruit. *Bulgarian Journal of Plant Physiology* **29**, 164-171.
- Soriano MA, Orgaz F, Villalobos FJ, Fereres E.** 2004. Efficiency of water use of early plantings of sunflower. *European Journal of Agronomy* **21**, 465-476.
- Taheri S, Saba J, Shekari F, Abdullah TL.** 2013. Effects of drought stress condition on the yield of spring wheat (*Triticum aestivum* L.) lines. *African Journal of Biotechnology* **10**, 18339-18348.
- Vaughan JG, Judd PA.** 2003. *The Oxford Book of Health Foods*. Oxford University Press. ISBN 0-19-850459-4, p 35.
- Wang S, Gong H, Zhu X, Chen K, Zhang C.** 2005. Silicon alleviate oxidative damage of wheat plants in pots under drought. *Journal of Plant Science* **169**, 313-321.
- Yahyavi Tabriz SH, Sadrabadi HR.** 2004. Effect of irrigation on yield and yield component of three spring canola cultivars under environmental of Tabriz. *Iranian Journal of Field Crops Research* **1**, 305-313.
- Yang JC, Zhang JH, Wang ZQ, Xu GW, Zhu QS.** 2004. Activities of key enzymes in sucrose-to-starch conversion in wheat grains subjected to water deficit during grain filling. *Journal of Plant Physiology* **135**, 1621-1629.
- Yari L, Modares MA, Soroushade A.** 2005. The effect of foliar application of Mn and Zn on qualitative characters in five spring safflower cultivars. *Journal of Water and Soil Science* **18**, 143-151.
- Yilmaz A, Ekiz H, Torun B, Gultekin B, Karanlik S, Bagci SA, Cakmak I.** 1997. Effect of different zinc application methods on grain yield and zinc concentration in wheat grown on zinc-deficient calcareous soils in Central Anatolia. *Journal of Plant Nutrition* **20**, 461-471.
- Zeidan MS, Mohamed FM, Hamouda HA.** 2010. Effect of Foliar Fertilization of Fe, Mn and Zn on

Wheat Yield and Quality in Low Sandy Soils Fertility.
World Journal of Agriculture Science **6**, 696-699.

Ziaei AH, Malakoti MJ. 2002. Effects of Fe, Mn, Zn and Cu fertilization on the yield and grain quality of wheat in the calcareous soils of Iran. Journal of Plant Nutrition **92**, 840-841.

Zhang J, Sun J, Duan A, Wang J, Shen X, Liu X. 2007. Effects of different planting patterns on water use and yield performance of winter wheat in the Huang-Huai-Hai plain of China. Agricultural Water Management **92**, 41-47.