



Hydrolysate of extruded soy proteins concentrate attenuates post myocardial infarction remodeling and improves heart health

Fatima Sultana^{1*}, Nazir Ahmad¹, Farhan Saeed¹, Fatima Jalal², Zeeshan Hafeez³

¹*Institute of Home and Food Sciences, Faculty of Life Sciences, Government College University, Faisalabad-Pakistan*

²*Department of Zoology, Faculty of Life Sciences, Government College University, Faisalabad-Pakistan*

³*Composés Alimentaires: Biofonctionnalités et Risques Neurotoxiques, Faculty of Sciences et Technologies, University of Lorraine, Nancy-France*

Key words: Soybean, HESPC, Intake, Myocardial Infarction.

<http://dx.doi.org/10.12692/ijb/15.1.176-185>

Article published on July 06, 2019

Abstract

Soybean is a good source of dietary proteins associated with numerous nutritional benefits and attenuates metabolic disorders like myocardial infarction (MI). Present study was aimed to explore the role of hydrolysate of extruded soy proteins concentrate (HESPC) intake in mitigation of MI. Isoprenaline induced MI rat groups (3-6) were fed on diet containing 5.07, 10.14, 15.21, 20.30 g proteins from Hydrolysates of extruded soy proteins concentrates (HESPC), respectively. Group 1 (non-induced MI) and group 2 (Isoprenaline induced MI) were fed on casein diet as control. Feed intake, body and heart weight, lactate dehydrogenase (LDH), creatine kinase-MB (CK), troponin, creatinine and urea, alanine aminotransferase (ALT), aspartate aminotransferase (AST), lipid profile, plasma proteins and homocysteine, cardiac and antioxidant enzymes and histopathology analysis were performed. The diet containing HESPC intake significantly changed body and heart weight. Lowest concentrations of LDH, CK-MB and troponin and highest concentration of superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx) were observed in rats group fed on 20.30 g proteins. Blood cells count was significantly decreased on intake of HESPC in all groups. Plasma lipids, proteins, homocysteine, hepatic enzymes showed decreasing trend in rat groups with increasing percentage of HESPC in diet. Histopathological results showed healing of injured heart tissues was significant on HESPC intake. HESPC diet intake can improve biochemical parameters and cardiac tissues health.

* **Corresponding Author:** Fatima Sultana ✉ drnazirahmad@gcuf.edu.pk

Introduction

Soybean is glycine rich proteins and utilizes for human consumption in the form of flour, infant formulas, concentrated, protein isolates and textured fibres. The soy enriched foods include tofu, cheese, tempeh, and drinks are used as proteins substitutes and vegetarian food product (Decroos *et al.*, 2005; Chauhan and Chauhan, 2007). These proteins are used as milk replacer in many products, but the effect of soy protein consumption reduces as compared to milk proteins due to anti-nutritional, antigenic activities and amount of incorporation (Hagen *et al.*, 2009). Based on investigations, epidemiological data and clinical trials, Food and Drug Administration (FDA) has approved health claim related to the soybean for being helpful in reduction of total and LDL cholesterol (Hasler *et al.*, 2004). MI is one of the health complications that is not only increasing in western world but also rising rapidly in developing countries (Sabeena *et al.*, 2004). MI is categorized as an imbalance of coronary blood demand and supply that leads to myocardial ischemic injury and ultimately damages the cardiomyocytes (Kloner, 2015). Previous studies suggest health boosting potential of soybean proteins against cholesterol, low density lipoprotein and triglycerides, along with increased resistance against lipid oxidation (Pihlanto-Leppälä *et al.*, 2000; Zhan and Ho *et al.*, 2005).

Potential health claims have led to increase soybean protein based foods as fastest growing category in food industry resulting in demand for manufacturing soy protein ingredients with improved processing characteristics. Heat processing of soybean reduced flatulence and anti-nutritional factors which increases the use of soy proteins in many products. Novel technologies help in reduction of anti-nutritional, anti-genic factors and improve the value addition of these proteins in product development and production of bioactive peptides which are useful for health of cardiac tissues (Pihlanto-Leppälä *et al.*, 2000). Bioactive peptides are inactive within the parent protein; however, thermal extrusion processing helps in production of bioactive peptides that can act as physiological modulators of

metabolism on their release (Milán-Carrillo *et al.*, 2006) In this context, the present study was aimed to assess the nutritional consumption of HESPC incorporated as casein replacer in rat diet and its biological evaluation against post myocardial infarction remodeling and heart health.

Materials and methods

Procurement of material

Soybean seeds of 95-I variety were procured from Ayub Agricultural Research Institute, Faisalabad, Pakistan and transferred to respective laboratory of Government College University, Faisalabad.

Preparations of defatted and extruded soybean proteins concentrate

Proteins hydrolysates have bioactive peptides that play important therapeutic role in human health. While thermal extrusion also helps in minimizing anti-nutritional factors and produce anticardiovascular and anti-inflammatory bioactive peptides. The seeds were cleaned, grounded and defatted with n-hexane, desolventized and pulverized to obtain the defatted soy flour. The proteins were concentrated according to described method (Wang *et al.*, 2007). Enzymatic hydrolysis of was performed to produce bioactive peptide from protein concentrate using Protamex® (EC 3.4.24.28), Neutrase® 1.5 MG (EC 82 3.4.24.28) and Flavourzyme® 500 MG (EC 3.4.11.1) purchased from Novozymes (Denmark). The ratio of enzymes: substrate was 1:100 using pH and temperature control devices. 1.0 g of enzymes was mixed in a 10 L reactor. The mixture was incubated for 8 hrs at optimal stirring and temperature and then placed into boiling water bath for 10 min to inactivate the enzyme and vacuum dried. Twin screw extruder was used for thermal extrusion (FMHE-36, Hunan Fumake Engineering Technology Co., 90 Ltd., Hunan, China) to produced extruded powder. The optimized conditions of barrel exit temperature (125°C), screw speed (100 rpm), feed flow rate (60 kg/h), feed moisture content (20%) were used in extrusion process.

Composition analysis of soybean seeds and HESPC

The proximate analysis of soybean seeds and HESPC as moisture (AACC Method No. 46-30), crude fat (AACC Method No. 30-25), crude protein (AACC Method 46-10), total ash (AACC Method No. 08-01) and Crude fibre (AACC Method no. 32-10) were conducted according to mentioned procedures of The American Association of Cereal Chemists (AACC, 2000).

Animal handling and MI induction

Rats were purchased from National Institute of Health (NIH), Islamabad and housed in the animal room of Government College University, Faisalabad, in accordance with institutional and National Policies and Guidelines for Care and Use of Laboratory Animals compliance with declaration of Helsinki (Ref.no.gcu/erc/1886). They were housed in polypropylene cages (47 cm×34 cm×20 cm) lined with husk, changed after every 24 hrs under a 12:12 hrs light dark cycle at 22°C and had free access to tap water and food. The rats were fed on a standard casein diet (Table 1). The rats were divided into 6 groups, each of 12 rats. 5 groups (2-6) were subcutaneously induced by isoprenaline hydrochloride injection (Sigma Aldrich) of 100 mg/kg body weight, once daily for two successive days (Zaafan *et al.*, 2013) and MI induction was confirmed by MI enzymes biomarkers of randomly selected from pool of 60 rats.

Feed formulation and study design

Experimental rats were divided into 6 groups, 12 in each group. Four MI induced groups (3-6) were fed on functional diets containing 5.07, 10.14, 15.21, 20.30 g proteins of HESPC while group 1 (non-induced and group 2 (MI induced) were fed on casein diet for 4 weeks as presented in Table 1.

Feed intake and body weight gain

Feed intake of experimental groups was calculated on daily basis by eliminating spilled diet from the total diet consumed during the entire experiment (Wolf and Weisbrode, 2003). Gain in body weight and heart weight of each group were determined accordingly (Seena *et al.*, 2006).

Biochemical analysis

The blood was drawn from tail of animals on first day of MI induction and then every week for confirmation of MI induction through troponin level. Animals were anesthetized with chloroform and sacrificed at the end of study for collection of blood plasma and hearts for further analysis. Blood samples were centrifuged at 4000 rpm for 10 min to obtain the serum which was used for determining various biochemical parameters. MI induced was evaluated by measuring markers of heart injury in serum, and antioxidant enzymes in heart tissue, observing pathological changes of tissue. RBCs, WBCs and platelet count were assessed using a Sysmex Cell Counter (Sysmex, Kobe, Japan) (Burr *et al.*, 1992). Serum creatinine was measured spectrophotometrically by using commercially available kit (Creatinine Jaffe Ecoline® diagnostic kit Merck). Blood glucose level was determined using diagnosis reagent kit (DiaSys Diagnostic Systems GmbH, Germany) (Husni *et al.*, 2016).

The serum total cholesterol (TC) and total triacylglycerol (TG) were determined by using DiaSys Diagnostic Systems USA reagent kit method. High density lipoprotein cholesterol (HDL-C) was determined by reagent kit method (Randox, Randox Laboratories LTD, UK). Calculation was done according to given equation for LDL-cholesterol [LDL-cholesterol = (Total cholesterol) – Triglycerides – (HDL- cholesterol)]. Very low density lipoprotein (VLDL) was calculated using following equation: VLDL-C = TG/5. Total plasma protein (Bradford, 1976) and homocysteine in blood plasma were determined through high performance liquid chromatography (Durand *et al.*, 1996).

Cardiac biomarker enzymes LDH, CK-MB and troponin were assayed using commercial kits according to previously described methods (Okinaka *et al.*, 1961; King, 1965). Serum ALT and AST were measured using commercially available kits (Reference # BT294QY; Randox Laboratories, United Kingdom). Heart homogenate was prepared according to method of (Noori *et al.*, 2009) using

Hematoxylin and eosin staining. Heart tissue CAT, SOD and GPx were measured through commercially available kits (Reference # KAO882, Abnova Corporation, Taiwan using method of Bancroft and Gamble. Similarly, histopathology analysis of heart tissues were also performed according to cited method (Bancroft and Gamble, 2002). The section mounted, and histological images were captured using 40x and 100x magnification of light microscope.

Statistical analysis

All experiments were performed in triplicate and results are presented as means \pm SD. The statistical analyses were performed using SPSS (version 20.0, IBM Corporation, NY, USA) and MS excel. Biochemical data were analysed using one-way ANOVA followed by t-test. Least Significant Difference test was used to determined significance difference at p value of < 0.05 (Steel *et al.*, 1997).

Results and discussion

Feed intake, body weight gain and heart weight

The means values of feed intake, body weight gain and heart weight are presented in Table 2. The results show insignificant increase in feed intake in all groups except group 2.

The highest feed intake (16.9 ± 1.06 g/day) was observed in group 6 and lowest feed intake (16.03 ± 1.26 g/day) was observed in group 2. HESPC intake resulted in significant increase ($p < 0.05$) in body weight. Highest body weight gain was observed in group 6 feeding on 20.3 g HESPC proteins diet.

Heart weight change was significant in all groups feeding on HESPC diets as compared to control group. Highest heart weight (3.2 ± 0.3 g) and lowest (2.60 ± 0.2 g) was observed in groups 2 and 6, respectively.

Table 1. Composition (%) of soybean seed, HESPC and rat diet.

Soybean seed and HESPC						
Composition (%)	Moisture	Protein (Nx6.25)	Fat	Crude fibre	Ash	
Soybean seed	4.6 \pm 0.14	34.52 \pm 0.34	21.03 \pm 0.37	7.41 \pm 0.16	6.15 \pm 0.13	
HESPC	3.91 \pm 0.36	73.08 \pm 0.25	1.88 \pm 0.25	3.27 \pm 0.17	6.93 \pm 0.30	
Rat groups and diet						
Rat groups	Groups 1	Groups 2	Groups 3	Groups 4	Groups 5	Groups 6
Diet type	Standard diet			HESPC diet		
composition (g/100g)	D ₀	D ₁	D ₂	D ₃	D ₄	
Corn starch	39.74	39.74	39.74	39.74	39.74	
Casein proteins	20.30	15.22	10.16	5.09	0.0	
Proteins (HESPC)	0	5.07	10.14	15.21	20.30	
Carbohydrate	13.2	13.2	13.2	13.2	13.2	
Fat	7.0	6.87	6.74	6.61	6.48	
Fat from HESPC	0	0.13	0.26	0.39	0.52	
Sucrose	10	10	10	10	10	
Fibre	5.0	4.73	4.55	4.32	4.09	
Fibre (HESPC)	0	0.27	0.45	0.68	0.91	
Vitamin mixture	1.0	1.0	1.0	1.0	1.0	
Mineral mixture	3.5	3.5	3.5	3.5	3.5	

Casein proteins were replaced with hydrolysate of extruded soy proteins concentrate (HESPC) D₀: Standard diet (casein diet), D₁₋₄: diets containing 5.07, 10.14, 15.21, 20.30 g proteins of HESPC, respectively, replaced with casein proteins. Each group had 12 rats; group 1 was not induced for myocardial infarction (MI) while groups 2-6 were induced for MI.

The previous studies have shown that increased protein content in soy protein diet results in intake of specific amino acids that were useful in mediation of myocardial infarction. The heart weight increased in isoprenaline treated rat groups was significantly

decreased ($p < 0.05$) because of amino acid present in diet given to infarcted rat groups (Moradi-Arzelo *et al.*, 2016). Ahmad and Ahmed also observed weight gain in soybean fed groups of rats from initial to final day of study (Ahmad & Ahmed, 2014).

Table 2. Effect of HESPC intake on feed intake, physiological and biological parameters of rats.

Physical parameters	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6	
Feed Intake (g/day)	16.68±0.58 ^a	16.03±1.26 ^b	16.26±0.94 ^{ab}	16.27±1.3 ^{ab}	16.29 ±0.86 ^{ab}	16.9 ±1.06 ^a	
Body weight (g)	264.7±8.1 ^a	225.3±6.3 ^c	239.6±7.1 ^d	253.31±5.6 ^c	255.73±6.01 ^{bc}	257.94±1.2 ^b	
Heart weight (g)	2.3±0.3 ^d	3.21±0.3 ^a	3.09±0.19 ^a	2.93±0.27 ^b	2.68±0.32 ^c	2.60±0.2 ^c	
Haematological parameters of rat groups							
Blood Cell Count	RBCs (10 ⁶ /μL)	5.67±0.24 ^e	11.25±0.68 ^a	8.61±0.44 ^b	7.95±0.71 ^c	7.12±0.54 ^d	5.73±0.38 ^e
	WBCs (10 ³ /L)	3.57±0.25 ^e	9.2 ± 0.4 ^a	7.56±0.17 ^b	6.31±0.48 ^c	4.01±0.31 ^d	3.95 ±0.06 ^d
	Platelets (10 ⁵ /mm ³)	2.96±0.25 ^e	4.72±0.49 ^a	4.16±0.53 ^b	3.98±0.82 ^b	3.36±0.46 ^c	3.11±0.62 ^d
CMP	Urea (mmol/L)	9.8±1.35 ^f	45.40±2.55 ^a	35.82±1.46 ^b	28.56±1.7 ^c	21.62±2.56 ^d	11.6±1.92 ^e
	Creatinine (μmol/L)	46±0.04 ^a	10.95±2.89 ^f	15.31±3.05 ^e	23.88±2.06 ^d	36.02±3.01 ^c	41.63±2.46 ^b
	ALT (U/L)	20.58±3.16 ^e	48.5±2.3 ^a	41.47±2.39 ^b	37.36±3.26 ^b	28.26±2.18 ^c	22.10±1.07 ^d
	AST (U/L)	48.74±4.00 ^e	96±5.24 ^a	85.47±3.02 ^b	69.42±3.24 ^c	48.47±4.65 ^d	37.01±3.8 ^e
Blood lipids, proteins profile and glucose level							
Serum Glucose (mg/dL)	Glucose	92.91±1.81 ^f	170±5.83 ^a	157.5±5.49 ^b	142.00±5.03 ^c	136.33±4.51 ^d	121.73±4.01 ^e
	TC	82.19±4.13 ^d	141.14±4.1 ^a	130.81±6.14 ^b	124.47±5.95 ^b	115.32±3.90 ^c	110.15±5.8 ^e
Lipid Profile (mg/dL)	HDL-C	56.58±2.14 ^a	18.5±2.1 ^e	44.71±1.82 ^d	48.48±2.97 ^c	50.56±3.18 ^{bc}	53.79±3.25 ^b
	LDL-C	43.28±1.95 ^e	85.6±3.5 ^a	71.12±2.1 ^b	68.79±2.9 ^b	63.57±1.73 ^c	52.47±3.1 ^d
	VLDL	20.84±3.08 ^b	26.4±0.6 ^a	24.97±1.29 ^a	24.01±0.91 ^a	23.02±1.1 ^{ab}	22.2±1.8 ^b
	TG	106±8.03 ^c	132.69±8.1 ^a	126.73±6.3 ^{ab}	121.31± 7.82 ^b	116.71±6.04 ^{bc}	110.5±5.2 ^c
Plasma proteins	Protein (g/L)	6.35±0.29 ^c	4.19±0.13 ^d	6.09±0.21 ^c	7.04± 0.23 ^b	7.98±0.19 ^{ab}	8.8±0.26 ^a
	Homocysteine (μmol/L)	8.9±0.4 ^e	22.72±0.4 ^a	14.9 ±0.2 ^b	12.7±0.71 ^c	11.53±1.53 ^c	9.8±0.82 ^d

Values are expressed as mean ±S.D. Different superscripts show the significant difference ($p < 0.05$) within row. HESPC: hydrolysate extruded soy proteins concentrate. Group 1, 2: Fed on casein diet, Group 3-6: fed on diets containing 5.07, 10.14, 15.21, 20.30 g proteins of HESPC, respectively, replaced with casein proteins. Values are expressed as mean ±SD. Different superscripts show the significant difference ($p < 0.05$) within row. RBCs: Red blood cells, WBCs: White blood cells, ALT: Alanine transaminase, AST: Aspartate transaminase, CMP: Comprehensive metabolic panel, TC: total cholesterol, HDL-C: high density lipoproteins cholesterol, LDL-C: low density lipoproteins cholesterol, VLDL: very low density lipoproteins cholesterol, TG: Triacyl glycerides.

Haematological parameters

The haematological results of WBC, RBC and platelets are presented in Table 2. A significant decrease of WBC, RBC and platelets was observed in MI induced group. Highest level of WBC, RBC and platelets count were $7.56 \pm 1.17 \times 10^3/\square L$, $8.61 \pm 0.44 \times 10^6/\square L$ and $4.16 \pm 0.53 \times 10^5/\text{mm}^3$, respectively in group 3 and lowest ($5.73 \pm 0.38 \times 10^3/\square L$, $3.95 \pm 0.06 \times 10^6/\square L$ and $3.11 \pm 0.62 \times 10^5/\text{mm}^3$) respectively, were observed in group 6. The studies show that increased RBCs, WBCs and platelets on MI induction can be reversed on consumption of soy proteins. In present research, we observed an increase of RBCs, WBCs and platelets after isoprenaline administration were gradually decreased in rats group consuming HESPC diet. Mean values of blood proteins, urea and creatinine are also presented in Table 2. A significant difference ($p < 0.05$) was observed in all MI groups.

The highest and lowest urea level decrease was observed in group 3 and 6, respectively. The mean comparison analysis of MI induced groups fed on HESPC diet show decreasing trend in urea level with increased HESPC proteins in diets. Creatinine level was significantly increased ($p < 0.05$) with increasing HESPC in diet. The highest ($41.63 \pm 2.46 \square \text{mol/L}$) and lowest ($15.31 \pm 3.05 \square \text{mol/L}$) values of creatinine were observed in group 6 and 3, respectively.

The creatinine level in MI infarcted rats was significantly improved on intake of HESPC in all rats groups. Previous studies show that serum creatinine and urea concentrations were significantly lowered in rats fed on soy protein based diet (Aukema and Housini, 2001). They proposed that soybean protein diet decreased urea concentrations in comparison to casein diet (Okafor and Ebuehi, 2016).

Table 3. Effect of HESPC intake on biomarkers in rat groups.

Biomarkers	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6
LDH (IU/L)	362±10.43 ^e	794±11.53 ^a	702±17 ^b	563±21 ^c	474±20 ^d	359±23 ^e
CK-MB (IU/L)	768.1±23.2 ^e	1104.4±64.3 ^a	943±11.8 ^b	886±12.46 ^c	799.53±26.8 ^d	715±20.2 ^f
Troponin (ng/dL)	7.3±0.31 ^e	15±0.23 ^a	11±0.12 ^b	9.3±0.26 ^c	8.9±0.02 ^d	6.4±0.15 ^f
SOD (U/mg prot)	42.6±2.6 ^a	24.61±1.91 ^d	26.9±2.4 ^d	28.6±2.3 ^{cd}	31.16±1.16 ^c	34.20±0.95 ^b
CAT (pmol/mg prot)	83.74±1.48 ^a	22.64±2.40 ^d	75.1±4.7 ^c	77.41±3.61 ^{bc}	78.07±3.55 ^b	79.13±3.64 ^{ab}
GPx (nmol/min/mg prot)	14.6±2.3 ^a	9.54±2.05 ^c	12.4±1.2 ^{bc}	13.11±1.50 ^b	14.08±1.92 ^{ab}	15.42±1.99 ^a

Values are expressed as mean ± S.D. Different superscripts show the significant difference ($p < 0.05$) within row. HESPC: hydrolysate of extruded soy flour proteins, Group 1, 2: Fed on casein diet, group 3-6: fed on diets containing 5.07, 10.14, 15.21, 20.30 g proteins of HESPC, respectively, replaced with casein proteins. LDH: Lactate dehydrogenase, CK-MB: Creatine kinase-MB, SOD: superoxide dismutase, CAT: Catalase, GPx: Glutathione peroxidases.

The significant ($p < 0.05$) decrease in all MI induced groups was observed. Highest values of AST and ALT (85.47±3.02 IU/L) and (41.47±2.39 IU/L), respectively, were observed in group 3 and lowest (37.01 ±3.8 IU/L) and (22.10±1.07IU/L), respectively, were observed in group 6. The serum levels of liver biomarkers in experimental rats after 4 weeks of administration of hard wheat bread and wheat soy breads were significantly decreased in experimental rats fed on wheat-soy bread compared to control group. In present finding, glucose level was increased after isoprenaline treatment that was remarkably decreased on consumption of HESPC diet. Anosike *et al.* described that glucose level in soybean supplemented rats was significantly lower than those without soybean supplementation. This may imply that soybean could reduce the incidence of hyperglycemia through production of antioxidant peptides (Anosike *et al.*, 2008).

Blood lipids, proteins and glucose level

The mean values of blood lipids, plasma proteins in Table 2 show that TC, HDL-C, LDL-C, VLDL and TG levels in all MI groups changed significantly. Highest decrease in TC, LDL-C, VLDL and TG and increase in HDL-C was observed in group 6. Maximum level of TC, LDL-C, VLDL and TG and minimum level of HDL-C was observed in group 3. These results are in accordance with the findings that after feeding on soy proteins result in TC, LDL-C, VLDL, and TG decrease. Our current findings show that increased soy protein in diet increased plasma protein levels, as like the results of Hagen *et al.* proposed that soybean diets

produced significant increases in the total plasma protein especially at 25%, 50% and 75% milk protein replacement (Hagen *et al.*, 2009). The plasma proteins show significant difference ($p < 0.05$) in total proteins and homocysteine levels in all MI groups. Maximum increase in proteins level and maximum decrease in homocysteine were observed in group 6. Minimum level of total proteins and maximum level of homocysteine were observed in group 3.

Cardiac biomarkers

Mean values of cardiac biomarkers LDH, CK-MB and troponin and antioxidant enzymes SOD, CAT and GPx are presented in Table 3. The results show a significant decrease in level of these enzymes. Intake of 20.3 g HESPC diet reduced maximum level of LDH, CK-MB and troponin while maximum level of these enzymes was observed in MI control group. Increase in heart weight might be associated with the increased water content, oedematous intramuscular space, increased protein content and infiltration of inflammatory cells to damaged areas that was decreased in group 3-6. Isoprenaline treated groups have increased the LDH, CK-MB and troponin levels and intake of HESPC diet decreased the level of these enzymes. Previous results showed decreasing trend in heart biomarkers on consumption of soybean flour within one week (Bertinchanta *et al.*, 2000). Hamed *et al.* (2010) also worked on effects of a soybean proteins diet in MI female albino rats. They concluded that soy protein diet intake increases CK-MB resulting in protection of heart against acute MI attack.




Fig. 1. Histopathological higher magnification (40x, 100X) images of cardiac muscles showing MI in control (A) and isoprenaline myocardial infarcted induced rats groups (B-F). A= regular arranged cardiac myocytes with branching of rat group fed on standard casein diet, with a pale acidophilic cytoplasm and single vesicular nucleus of the cardiac muscle fibers. B-F= Rats were isoprenaline induced MI and fed on hydrolysate of extruded soy protein concentrate 5.07, 10.14, 15.21, 20.30 g proteins. Myocardial infarction heart tissues damage was observed showing cardiac myocytes with areas of irregular arrangement and necrosis in B-F that was improved in strength and tissue structure gradually after intake.

Similarly, troponin levels were increased in disease induced group that showed decreased trend when HESPC diets were given to groups 3-6 with different concentrations. Significant difference ($p < 0.05$) was observed in all MI groups on intake of HESPC. Minimum activity of SOD, CAT and GPx (26.9 ± 2.4 U/mg prot), (75.1 ± 4.7 pmol/mg prot) and (12.4 ± 1.2 nmol/min/mg prot), respectively, was observed in group 3. Maximum activity of SOD (34.20 ± 0.95 U/mg prot), CAT (79.13 ± 3.64 pmol/mg prot) and GPx (15.42 ± 1.99 nmol/min/mg prot) was observed in group 6. Antioxidant enzymes of heart tissues such as SOD, CAT, GPx were decreased after MI induction and significantly increased on HESPC intake. Hagen *et al.*, worked on comparative effect of soy protein isolate and casein diet in MI rats. They showed that SOD, CAT and GPx were increased in isolated soy protein fed group of rats as compared to casein.

Histopathological study

The histopathological images (Figure 1) show that the intake of HESPC improves the tissue structure of injured heart. The results indicate that heart tissues of group 1 arranged regularly the cardiac myocytes branching with a pale acidophilic cytoplasm and single vesicular nucleus of the cardiac muscle fibres. The broken tissues due to MI can easily be observed in figure 1B, the heart tissues of group 2 were highly damaged showing irregularly arranged cardiac myocytes with areas of irregular arrangement and necrosis. The figure 1C shows heart tissues of group 3 started to improve the strength that gradually increased in group 4, 5 and 6 with increase intake of HESPC as shown in figure 1(D,E,F), respectively. Soybean protein can be valuable in protecting the heart against the attack of acute MI. A significant improvement in the ventricular function was examined after MI and consequently preventing the progression towards a severe heart failure by using isolated soy proteins in diet. We observed that the preventive effect of HESPC products in the terms of post MI dysfunction concluding that favourable effect is possibly secondary to the antioxidant activity of soy protein and bioactive peptides produced during hydrolysis and extrusion (Hagen *et al.*, 2009).

Conclusion

Soy proteins can be used for preparation of functional food and its efficacy can further be enhanced by food processing technology. The extrusion processing of soy flour and enzymatic hydrolysis result in production of bioactive peptides that can be effectively used for MI ailment. The intake of defatted HESPC significantly improves the biological parameter against MI. HESPC diet containing 5.07, 10.14, 15.21, 20.30 g proteins can effectively decrease glucose, LDL-C, VLDL, TG and TC, increase HDL-C and improved haematological and CMP levels in rats. The MI induced was reversed with the help of HESPC diet intake though improved tissues structures of heart and cardiac health and antioxidant enzymes. It is suggested that HESPC could be used food tool for treating MI induced hyperlipidemia and hyperglycemia.

Acknowledgement

The authors are thankful to Higher Education Commission (HEC) Pakistan, for the support of some experimental analysis, University of Agriculture Faisalabad-Pakistan, and Department of Pharmacy, Government College University Faisalabad-Pakistan and also for animal keeping and handling.

References

AACC. Approved methods of American Association of Cereal Chemists. 2000. 10th ed. St. Paul, Minnesota: The American Association of Cereal Chemists, Inc.

Ahmad S, Ahmed I. 2014. Response of wistar rats to broiler chicken feed and soybean on body weight, obesity and weight of selected visceral organs. *Pakistan Journal of Biochemistry and Molecular Biology*, **47(3-4)**, 137-140.

Anosike CA, Obidoa O, Ezeanyika LUS. 2008. Beneficial effects of soybean diet on serum marker enzymes, lipid profile and relative organ weights of wistar rats. *Pakistan Journal of Nutrition* **7(6)**, 817-822.

<http://dx.doi.org/10.3923/pjn.2008.817.822>

- Aukema HM, Housini I.** 2001. Dietary soy protein effects on disease and IGF-I in male and female Han: SPRD-cy rats. *Kidney International* **59**, 52–61.
<https://doi.org/10.1046/j.1523-1755.2001.00465.x>
- Bancroft JD, Gamble M.** 2002. Theory and practice of histological techniques, (5th ed.) Churchill Livingstone, London, New York and Philadelphia.
- Bertinchanta JP, Roberta E, Polge A, Marty-Doublet C, Fabbro-Peray P, Poirey S, Aya G, Juan JM, Ledermann B, de la Coussaye JE, Dauzat M.** 2000. Comparison of the diagnostic value of cardiac troponin I and T determinations for detecting early myocardial damage and the relationship with histological findings after isoprenaline-induced cardiac injury in rats. *Clinica Chimica Acta* **298**, 13-28.
[https://doi.org/10.1016/S0009-8981\(00\)00223-0](https://doi.org/10.1016/S0009-8981(00)00223-0)
- Bradford MM.** 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72**, 248-254.
[https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- Burr ML, Holliday RM, Fehily AM, Whitehead PJ.** 1992. Haematological prognostic indices after myocardial infarction: evidence from the diet and reinfarction trial (DART). *The European Heart Journal* **13**, 166-170. PMID:1313369.
- Chauhan OP, Chauhan GS.** 2007. Development of anti-nutrients free soy beverage using germinated soybean. *Journal of Food Science and Technology* **44**, 62-65.
- Decroos K, Vanhemmens S, Cattoir S, Boon N, Verstraete W.** 2005. Isolation and characterisation of an equol-producing mixed microbial culture from a human faecal sample and its activity under gastrointestinal conditions. *Archives of Microbiology* **183**, 45-55.
<https://doi.org/10.1007/s00203-004-0747-4>
- Durand P, Fortin LJ, Luissier-Cacan S, Davignon J, Blache D.** 1996. Hyperhomocysteinemia induced by folic acid deficiency and methionine load-application of a modified HPLC method. *Clinica Chimica Acta* **252**, 83-93.
[http://dx.doi.org/10.1016/0009-8981\(96\)06325-5](http://dx.doi.org/10.1016/0009-8981(96)06325-5)
- Hagen MK, Lehenbauer-Lu'dke AR, Paludo AC, Schenkel P, Goncalves L, Fernandes TG, Caron R, Llesuy S, Mill JG, Bello-Klein A.** 2009. Diet with isolated soy protein reduces oxidative stress and preserves ventricular function in rats with MI. *Nutrition, Metabolism & Cardiovascular Diseases* **19**, 91-97.
<http://dx.doi.org/10.1016/j.numecd.2008.03.001>
- Hamed GM, Bahgat NM, EI-Agaty SM, Soliman GZA, Emara MM.** 2010. Effects of a soybean protein diet on ovariectomised female albino rats subjected to MI. *Singapore Medical Journal* **51(10)**, 781-789. PMID:21103813.
- Hasler CM, Bloch AS, Thomson CA, Enrione E, Manning C.** 2004. Position of the American Dietetic Association: functional foods. *Journal of the American Dietetic Association* **104(5)**, 814-26.
<https://doi.org/10.1016/j.jada.2004.03.015>
- Husni A, Purwanti D, Ustadi.** 2016. Blood Glucose Level and Lipid Profile of Streptozotocin-induced Diabetes Rats Treated with Sodium Alginate from *Sargassum crassifolium*. *Journal of Biological Sciences* **16**, 58-64.
<https://doi.org/10.3923/jbs.2016.58.64>
- King J.** 1965. The dehydrogenases or oxidoreductases-lactate dehydrogenase. In: D Van, Ed. *Practical Clinical Enzymology* 93–193. London, Van Nostrand.
- Kloner RA.** 2015. New observations regarding post-ischemia/reperfusion myocardial swelling. *The Journal of the American College of Cardiology* **65**, 324–326.

<https://doi.org/10.1016/j.jacc.2014.11.006>

Milán-Carrillo J, Gutiérrez-Dorado R, Perales-Sánchez JXK, Cuevas-Rodríguez EO, Ramírez-Wong B, Reyes-Moreno C. 2006. The optimization of the extrusion process when using maize flour with a modified amino acid profile for making tortillas. *International Journal of Food Science and Technology* **41**, 727-736.

<https://doi.org/10.1111/j.1365-2621.2005.00997.x>

Moradi-Arzeloo M, Farshid AA, Tamaddonfard E, Asri-Rezaei S. 2016. Effects of histidine and vitamin C on isoproterenol-induced acute MI in rats. *Veterinary Research Forum* **7(1)**, 47-54. PMID: [27226887](https://pubmed.ncbi.nlm.nih.gov/27226887/)

Noori S, Nasir K, Mahboob T. 2009. Effects of cocoa powder on oxidant/antioxidant status in liver, heart and kidney tissues of rats. *Journal of Animal and Plant Sciences* **19(4)**, 174-178.

Okafor HK, Ebuehi OAT. 2016. Defatted Soya Flour Supplementation of Wheat Bread Confers Oxidative, Renal, Hepatic and Cardiovascular Protective Effects in Wistar Rats. *International Journal of Biochemistry Research & Review* **10(1)**, 1-14.

<http://dx.doi.org/10.9734/ijbcr/2016/22912>

Okinaka S, Kumagai H, Ebashi S, Sugata H, Momoi H, Toyokura Y, Fujie Y. 1961. Serum creatine phosphokinase activity in progressive muscular dystrophy and neuromuscular diseases. *Archives of Neurology* **4**, 520-5.

<https://doi.org/10.1001/archneur.1961.00450110050006>

Pihlanto-Leppälä AL, Koskinen P, Piilola K, Tupasela T, Korhonen H. 2000. Angiotensin I-converting enzyme inhibitory properties of whey protein digests: Concentration and characterization of active peptides. *Journal of Dairy Research* **67**, 53-64.

<http://dx.doi.org/10.1017/S0022029999003982>

Sabeena FKH, Anandan R, Kumar SHS, Shiny KS, Sankar TV, Thankappan TK. 2004. Effect of squalene on tissue defense system in isoproterenol-induced MI in rats. *Pharmacological Research* **50**, 231-6.

<https://doi.org/10.1016/j.phrs.2004.03.004>

Seena S, Sridhar KR, Arunb AB, Young CC. 2006. Effect of roasting and pressure cooking on nutritional and protein quality of seeds of mangrove legume *Canavaliacathartica* from southwest coast of India. *The Journal of Food Composition and Analysis* **19**, 284-29.

<https://doi.org/10.1016/j.jfca.2005.05.004>

Steel RGD, Torrie JH, Dickey D. 1997. Principles and procedures of statistics, a biometrical approach. New York: McGraw Hill Book Co Inc: USA.

Wang J, Yuan X, Jin Z, Tian Y, Song H. 2007. Free radical and reactive oxygen species scavenging activities of peanut skins extract. *Food chemistry* **104(1)**, 242-250.

<https://doi.org/10.1016/j.foodchem.2006.11.035>

Wolf BW, Weisbrode SE. 2003. Safety evaluation of an extract from *Salacia oblonga*. *Food and Chemical Toxicology* **41**, 867-874.

[https://doi.org/10.1016/S0278-6915\(03\)00038-3](https://doi.org/10.1016/S0278-6915(03)00038-3)

Zaafan AM, Zaki FH, El-Brairy IA, Kenawy AS. 2013. Protective effects of Atorvastatin and quercetin on Isoprenaline-induced MI in rats. *Bulletin of Faculty of Pharmacy, Cairo University* **51**, 35-41.

<https://doi.org/10.1016/j.bfopcu.2013.03.001>

Zhan S, Ho SC. 2005. Meta-analysis of the effects of soy-protein containing isoflavones on the lipid profile. *The American Journal of Clinical Nutrition* **81**, 397-408.

<https://doi.org/10.1093/ajcn.81.2.397>