



## RESEARCH PAPER

## OPEN ACCESS

## Productive performance of *Achai* and crossbred cows under different farm managements in Dir-Kohistan Hindukush Mountainous Range

Zia ur Rehman Khalil<sup>\*1</sup>, Abdur Rehman<sup>1</sup>, Muhammad Subhan Qureshi<sup>1</sup>,  
Muhammad Saleem<sup>2</sup>, Shakoor Ahmad<sup>3</sup>

<sup>1</sup>Department of Livestock Management, Breeding and Genetics, The University of Agriculture, Peshawar, Pakistan.

<sup>2</sup>Livestock and Dairy Development Department, Government of Khyber Pakhtunkhwa, Pakistan

<sup>3</sup>Department of Animal Health, The University of Agriculture, Peshawar, Pakistan

**Key words:** *Achai* cow, Crossbreeding, Farming systems, Economic traits, Non-genetic factors Hindukush Mountains

<http://dx.doi.org/10.12692/ijb/16.3.576-594>

Article published on March 30, 2020

### Abstract

The study was to understand the prevailing knowledge and applied practices regarding factors affecting the productive performance of *Achai* and Jersey X *Achai* cows under different farming systems in Hindukush Mountains of Pakistan. During survey, two types of rural housing systems were observed which were termed as rural traditional farming system (RFS) and rural progressive farming (PFS) system on the basis on differences in nutrition, health and general management and breeding planning. Overall, the productive performance of crossbred cows (except CSR) was significantly ( $P < 0.05$ ) better than *Achai* cows. Comparing different farm managements, both breeds had significantly ( $P < 0.05$ ) better performance in PFS. Regarding factors effect, the DMY, LL and LY was significantly better in good condition *Achai* cows in PFS particularly during summer season. Parity and age showed no effect on productive performance of *Achai* cows in any farming system. In crossbred cows, all the studied productive traits were significantly ( $P < 0.05$ ) better during summer season. Adult (4-6 years age) crossbred cows had significantly ( $P < 0.05$ ) higher DMY and LY due to longer LL in their 3<sup>rd</sup> and 4<sup>th</sup> parities. On the other side, the daily milk yield, Lactation Length, Lactation Yield and Calf Birth Weight was not affected ( $P > 0.05$ ) by BCS. The study revealed that crossbreeding of *Achai* cows with Jersey semen and improving farm managements markedly improved the productive performance in study area. Further, the introduction of *Achai* cows to intensive farm conditions (as in state farm) significantly affected its productive potentials.

\* Corresponding Author: Zia ur Rehman Khalil ✉ [ziakhalil@sbbu.edu.pk](mailto:ziakhalil@sbbu.edu.pk)

## Introduction

Pakistan holds 42.6 million heads of cattle population among which 70% shares non-descript cattle. These cattle have dynamic economic performances under varying climatic conditions with in Pakistan but mostly have low production record (Sattar *et al.*, 2005). In spite of low performance, livestock plays a significant role in national economy particularly in uplifting the socio-economic status of rural population (Khattak *et al.*, 2018). Farmers of Northern Hindukush region has substantial relation with livestock and rely on dairy products for basic life support (Saleem *et al.*, 2010). In Northern Hindukush region of Pakistan, the production and distribution system of livestock and livestock products are traditional and poorly developed. Livestock are reared under sedentary, semi-nomadic and nomadic systems (Sadiq *et al.*, 2003). Livestock farmers belong to small/landless farmers and remain under extensive production system. On the other side, utilization of inputs obviously influences the profitability of dairy industry because of substantial relationship with economic traits of cattle (Williams *et al.*, 2011).

Therefore selection of economic traits has traditionally paid more attention in breeding policies for profitability of dairy farming (Gonzalez Recio *et al.*, 2014). Recently, genetic improvement in economic traits with better resistance to non-genetic factors affecting its performance has been emphasized in many countries (Miglior *et al.*, 2005).

Lowered feed cost, higher and quality milk production and health issues are key factors for adopting genetic improvement programmes (Bell *et al.*, 2013; Connor, 2015) therefore the inclusion of economic traits in breeding and selection schemes to improve profitability estimates of different cattle breeds had been widely studied (Sölkner *et al.*, 2000; Fernández-Perea and Jiménez, 2004). In response, it was observed that expressing the economic traits primarily depend upon certain factors including basic genetic makeup, feed conversion efficiency, physiological condition of animals, adoptability to climatic conditions and heredity flow of specific breed immunity to certain diseases (Epaphras *et al.*, 2004;

Naceur *et al.* 2014). Husbandry practices, nutrition, lactation, age and calving seasons are other prominent factors affecting productive and reproductive economic traits of cattle (Msanga *et al.*, 2000). *Achai* cattle, a better choice of farmers rearing livestock in remote areas of Himalayan Hindukush Mountains are inherently slow-maturing and low-milk producers (Saleem, 2012). *Achai* cattle termed as an economical livestock in high altitudes of northern areas of Pakistan has some promising characteristics i.e impressive performance on suboptimal quality roughages, enhanced immunity and in some aspect showed comparatively good reproductive record than any other cattle breeds in Pakistan (Saleem *et al.*, 2012). High resistant and well adaptive to harsh climatic conditions make it more favourable to graze on rugged mountain terrain. Priority for increased milk production, progressive use of mechanized tools for agricultural purposes instead of cattle bull as draught animal, and easily accessibility to artificial insemination initiated crossbreeding services with exotic cattle specifically with Jersey breed.

Negligible research has been conducted on economic traits of *Achai* cattle, which is key source of income in the Northern Hindukush Region of Pakistan.

In addition, the increased demand of milk and meat production for human consumption has raised questions that whether *Achai* breed has the potentials to support current demand for milk and meat production or either crossbreeding of *Achai* cow with Jersey breed could be an alternate source for improved performance. Present study was designed with the objectives to acquire the knowledge regarding factors affecting economic traits of *Achai* and Jersey X *Achai* cattle in Hindukush Mountains of northern Khyber Pakhtunkhwa province of Pakistan.


## Materials and methods

### Study area

The broader home tract of the *Achai* cattle is spread over the North-Western Hindu Kush Mountains of Khyber Pakhtunkhwa, Pakistan (Fig. 1). The study area is situated 34° 10N latitude and 72°20E longitude.

The area falls in both subtropical dry temperate zone as well as moist temperate zone of Hindukush series in Pakistan. Geographically, Afghanistan lies in west, Swat in East, District Chitral in North and Dargai council of Malakand Division in South of the study area (Hazrat *et al.*, 2015).

Climatic conditions of the area are moderate. Annual precipitation and relative humidity of the study area ranges from 70-300 mm and 15 to 60%, respectively while temperature ranges from 20 to 33°C and -1 to 15°C during summer and winter, respectively (Fayaz, 2017).



**Fig. 1.** Map of study area

#### *Selection of animals*

Pure *Achai* and Jersey X *Achai* cows were selected to study its productive performance under the effect of different factors i.e. body condition score (BCS), Age, parity and calving season. Age of cow was determined through dentition formula as per the guidelines of Pace and Wakeman, (2003). Parity of animal was calculated as per information provided by respondents during the survey. Body condition score of animal was confirmed on five point scale as described by Peters and Ball (1987).

#### *Classification of fixed factors*

To evaluate the effect of different factors on productive performance of cows, each factor was further subdivided into different levels in each farming system on the basis of 1: Age (young cows; 2-4 years age, adult cows: 5-6 years age,

older cows: 7-8 years age), 2: Parity (early parity cows: 1<sup>st</sup> and 2<sup>nd</sup> parity, mid parities cows; 3<sup>rd</sup> and 4<sup>th</sup> parity, advanced parity cows; 5<sup>th</sup> and 6<sup>th</sup> parity), 3: Calving season (Summer and winter calvers) and 4: BCS (cow with BCS < 2.5 and BCS > 2.5). In addition to state farms, different kinds of rural housing systems were observed on the basis of differences in nutrition, housing design and animal management, approach to address animal health and reproduction and orientation toward livestock marketing. The effect of fixed factors was therefore studied in each farming system.

#### *Attributes of rural farming systems*

Rural farmers adopted various approaches in rearing livestock which resulted in rural traditional farming system (RFS) and rural progressive farming system (PFS).

RFS mainly existed on hilly areas with 6747±265.25ft altitude where farmers preferred mixed type herding with 3-5 cattle heads in extensive management practices and had limited market access due to remote hilly areas. The farming objectives were milk production, animal sale and gifting, ploughing, replacement and meat purposes for home consumption. RFS farmers practiced indiscriminate breeding mostly with *Achai* bull, occasionally inseminated cows with Jersey semen. Animals were kept in sheds made of clay walls or stones with soil floor. Animals chained with tree trunks were considered as open yard. Animals were stall fed during evening and allowed to graze during day time on rugged mountain terrains. Animal received suboptimal quantity and quality nutrition and occasionally supplemented for increased production. In RFS, mainly homemade remedies were used for treating animals. The trend of using de-wormers and vaccination was occasional.

PFS mostly prevailed on plain areas with an altitude of 4343±152.60ft. PFS farmers kept 5-10 cattle per household in an intensive management system focused on milk production for sale. Farmers adopted crossbreeding policy and prefer Jersey semen for insemination of *Achai* cows. For cattle housing, farmers used cemented blocks for walls and concrete or bricks for flooring. Generally wooden planks or

tree branches were used for the construction of open paddock. In PFS, two times stall feeding with optimum quantity and quality feed is provided. Provision of concentrates and supplementation for increased production is common. For animal health, professional veterinarians are preferred however homemade remedies are also used. Contrary to RFS, regular vaccination and use of de-wormers was common in PFS.

#### Nutrition management

In the study area, cattle nutrition varied on the basis of farming system, season and breed of cow. Quantity and type of feed provided to pure *Achai* and crossbred cows in different farming systems during summer and winter seasons is detailed in the Table I. Briefly, crossbred cattle received comparatively more different kind of concentrates, green fodders, weed thinning and tree leaves in rural progressive farming system. While in winter, the quantity of concentrates increased in all farming systems for both breeds.

Further, cow grazing for a specific time in rural traditional farming system and provision of a special energy supplementation containing wheat flour, ghee and different herbs mixed in the fluid of locally produced sweetener from sugarcane as first diet for freshly parturated cows was also common in the surveyed households of study area.

**Table 1.** Quantity of feed (kg) received by animals during different seasons in study area.

| Feed Ingredients (kg) | Summer season     |                   |                   |                   |
|-----------------------|-------------------|-------------------|-------------------|-------------------|
|                       | RAFS <sup>1</sup> | RCFS <sup>2</sup> | PAFS <sup>3</sup> | PCFS <sup>4</sup> |
| Concentrate           | 1                 | 0.86              | 1.53              | 1.95              |
| Dry bread             | 0.50              | 0.53              | 1.06              | 1.00              |
| Green Fodder          | 7.00              | 10.56             | 7.85              | 14.68             |
| Wheat Straw           | 1.00              | 1.03              | 1.00              | 0.50              |
| Weed Thinning         | 0.00              | 0.00              | 1.00              | 0.00              |
| Tree Leaves           | 2.50              | 2.67              | 2.53              | 4.00              |
| Maize Stover          | 0.00              | 0.52              | 1.00              | 0.00              |
| Winter season         |                   |                   |                   |                   |
| Concentrate           | 1.00              | 2.00              | 2.00              | 3.00              |
| Dry bread             | 0.73              | 1.40              | 1.00              | 2.00              |
| Green Fodder          | 0.00              | 1.38              | 0.00              | 0.00              |
| Wheat Straw           | 5.33              | 7.05              | 5.03              | 6.64              |
| Weed Thinning         | 1.73              | 0.52              | 1.50              | 1.39              |
| Tree Leaves           | 0.60              | 0.61              | 1.52              | 1.89              |
| Maize Stover          | 2.60              | 4.57              | 4.20              | 5.50              |

<sup>1</sup>RAFS, Rural *Achai* farming system, <sup>2</sup>RCFS, Rural crossbred farming system, <sup>3</sup>PAFS, Progressive *Achai* farming system, <sup>4</sup>PCFS, Progressive crossbred farming system.

*Collection of feed samples and chemical analysis*

Seventy five (200-250g) representative samples of feed provided in each farming system (N=25 for each farming system) randomly collected in large paper bags were analysed for chemical composition including dry matter, moisture, crude protein and

ash, crude fibre, EE, NFE and TDN content. Feed samples were analysed following the procedure of AOAC, (1995). All standard protocols for feed sample collection, processing and chemical analysis were ensured. The nutritive value of feed samples is detailed in the Table 2.

**Table 2.** Nutritive value of feed ingredients provided to animals in studied area.

| Grasses                     | %DM   | Moisture | %CP   | %CF   | Ash   | %EE   | %NFE  | %TDN  |
|-----------------------------|-------|----------|-------|-------|-------|-------|-------|-------|
| <i>Poa alpine</i>           | 92.73 | 07.27    | 21.20 | 19.67 | 09.01 | 06.20 | 51.74 | -     |
| <i>Trifolium repens</i>     | 90.13 | 09.87    | 22.62 | 19.64 | 08.32 | 04.60 | 44.62 | -     |
| <i>Plectranthus rogusus</i> | 93.20 | 06.80    | 13.11 | 21.63 | 08.87 | 05.40 | 42.50 | -     |
| Concentrates                |       |          |       |       |       |       |       |       |
| Wheat Bran                  | 88.72 | 09.37    | 12.03 | 09.84 | 04.63 | 03.12 | 68.73 | 74.05 |
| Cotton seed cake            | 90.95 | 09.13    | 22.37 | 28.41 | 06.58 | 07.62 | 34.60 | 64.52 |
| Mustard seed cake           | 91.76 | 08.32    | 32.08 | 19.84 | 12.02 | 09.64 | 26.21 | 84.63 |
| Commercial. Concentrates    | 90.73 | 09.63    | 17.17 | 10.16 | 04.14 | 04.95 | 52.97 | 72.48 |
| Crop Residues               |       |          |       |       |       |       |       |       |
| Wheat Straw                 | 89.94 | 9.06     | 03.21 | 41.81 | 10.9  | 00.12 | 44.23 | 43.63 |
| Maize Stover                | 93.66 | 6.42     | 04.60 | 45.72 | 12.3  | 01.75 | 39.72 | 54.38 |
| Fodder                      |       |          |       |       |       |       |       |       |
| Barseem                     | 13.64 | 86.36    | 19.34 | 21.41 | 16.28 | 01.86 | 43.47 | 61.65 |

*Data collection*

Between December 2016 and November 2017, 720 households at Sheringal, Dogdara, Sharmai, and Jandool valley and 356 animals at two state farms i.e Livestock Research and Development Station and *Achai* Cattle Conservation Farm, Dir (Lower), were surveyed for data collection (Fig.1) through a structured questionnaire by face to face interacting farmers with repeated questioning to excerpt concrete information and tracking concerned animals for confirmation of breed, housing pattern and management, use of antiparasitic, growth promoting and milk boosting medicine, nutrition management and productive and reproductive performance. Data on productive performances include daily milk yield (DMY), lactation length (LL), lactation milk yield (LY), calf birth weight (CBW).

Daily milk yield was calculated by emptying cow udder completely through gentle hand milking in morning while practicing same procedure next day at evening and combining the weight of both times milking to determine the daily milk yield. The purpose of such practice was to ensure calf health and reluctance of local farmers to collect milk from cow by prohibiting calf from natural udder suckling. Lactation length was calculated as interval (in days)

from parturition till complete cessation of milk yield either by cow or voluntarily by farmers. For calculation of lactation length, the trust upon the responses of farmers during survey was only option. However, during pooling of data, it was ensured to remove any unusual observation which exceeds normal ranges. Lactation yield was calculated by multiplying daily milk yield of each cow with its lactation length. Calf birth weight was calculated by weighing young calves within three (03) days after births mostly by spring balance due to lack of facilities to use electronic scale. However, electronic scale was also used where possible and average data was compared with records of spring balance before entering in record sheets.

*Statistical analysis*

Initially data was grouped as *Achai* vs crossbred cows to study the difference between breeds irrespective of farming or other fixed factors effect. For the difference between breeds two sample t-test at 5% level of probability was used. Upon significant differences observed in studied parameters between breeds, it was decided to analyse further data separately for each breed under all fixed factors. Therefore, during the 3<sup>rd</sup> step, within breed combined analysis of variance technique was followed

separately for *Achai* and crossbred cows to study the main effect as well as the interaction effect between the farming systems and fixed effects (i.e. body conditions, age, parity and season) according to Annicchiarico (2002). Upon significant results for interaction effect the data was further analysed at individual level in each farming system to study the variation among the levels of fixed effects. In addition, to study the effect of farming systems on each level of fixed effects the data was also analysed within each level of body condition, age, parity and season across different farming systems. Mean separation was carried out using Least Significant

Difference test (LSD) following Steel and Torrie (1985) where required.

## Results

Overall productive performance of *Achai* and crossbred cows

The overall productive performance of *Achai* and crossbred cows studied during this investigation is presented in the Table 3.1. The crossbred cattle had significantly higher daily milk yield ( $P < 0.05$ ), longer lactation length ( $P < 0.05$ ), milk yield per lactation ( $P < 0.05$ ) and heavier calves ( $P < 0.05$ ) at birth than *Achai* cows.

**Table 3.1.** Overall productive performances of *Achai* and crossbred cows.

| Productive traits        | <i>Achai</i> cows         | Crossbred cows             | P-Value |
|--------------------------|---------------------------|----------------------------|---------|
| Daily milk yield (L)     | 3.55±0.07 <sup>b</sup>    | 4.57±0.07 <sup>a</sup>     | 0.00    |
| Lactation length (days)  | 222.94±2.99 <sup>b</sup>  | 257.26±2.93 <sup>a</sup>   | 0.00    |
| Milk yield/lactation (L) | 814.90±21.16 <sup>b</sup> | 1179.38±20.73 <sup>a</sup> | 0.000   |
| Calf birth weight (kg)   | 15.20±0.17 <sup>b</sup>   | 16.29±0.16 <sup>a</sup>    | 0.04    |

Means with different superscripts are significantly different at  $P < 0.05$

### Effect of farming systems on productive performance of *Achai* and crossbred cows

Productive performance of *Achai* and crossbred cows in different farming systems is presented in the Table 3.2. Farming systems showed significant ( $P < 0.00$ ) effect on productive performance of *Achai* and

crossbred cows with better results in rural progressive farming system. It was also revealed from the present.

Study that introducing *Achai* cows to intensive farming systems i.e state farms significantly affected its productive performances.

**Table 3.2.** Productive performance of *Achai* and crossbred cows under different farming systems.

| Performance  | Breed        | SF                  | RFS                        | PFS                        | P-value |
|--------------|--------------|---------------------|----------------------------|----------------------------|---------|
| DMY (litres) | <i>Achai</i> | 2.00 <sup>c</sup>   | 2.40±0.08 <sup>b</sup>     | 4.18±0.23 <sup>a</sup>     | 0.00    |
|              | Crossbred    | *                   | 4.44±0.02 <sup>b</sup>     | 5.53±0.47 <sup>a</sup>     | 0.03    |
| LL (days)    | <i>Achai</i> | 183.37 <sup>b</sup> | 191.73±3.05 <sup>b</sup>   | 244.78±4.04 <sup>a</sup>   | 0.00    |
|              | Crossbred    | *                   | 236.73±5.15 <sup>b</sup>   | 283.47±3.05 <sup>a</sup>   | 0.00    |
| LY (days)    | <i>Achai</i> | 381.87 <sup>b</sup> | 462.50±19.34 <sup>b</sup>  | 1019.56±17.62 <sup>a</sup> | 0.00    |
|              | Crossbred    | *                   | 1048.19±22.66 <sup>b</sup> | 1564.45±24.43 <sup>a</sup> | 0.00    |
| CBW (kg)     | <i>Achai</i> | 15.27 <sup>a</sup>  | 16.51±.08 <sup>b</sup>     | 16.56±.04 <sup>b</sup>     | 0.05    |
|              | Crossbred    | *                   | 15.56±.09 <sup>b</sup>     | 16.66±.07 <sup>a</sup>     | 0.00    |

Means within rows with different superscripts differs significantly at  $P < 0.05$  separately for each breed.

### Factors affecting the daily milk yield of *Achai* and crossbred cows under different management systems

The effect of body condition score, parity, season and age on daily milk yield of *Achai* and crossbred cows in SF, RFS and PFS is presented in the Table 3.3. BCS significantly ( $P < 0.05$ ).

Affect DMY of *Achai* cows reared under PFS with better production in cows with more than 2.50 BCS.

Calving season also had significant ( $P < 0.05$ ) effect on DMY with higher production in *Achai* cows calved in summer season. The effect of parity and age on same trait was not significant ( $P > 0.05$ ). In crossbred cows, parity, season and age had significant ( $P < 0.05$ ) effect on DMY. Higher yield was observed in 3<sup>rd</sup> and 4<sup>th</sup> parity group, summer calvers and 4-6 years age group cows under both RFS and PFS. The effect of BCS on DMY of crossbred cows was not significant ( $P > 0.05$ ).



**Table 3.3.** Effect of BCS, parity, season and age on DMY (kg) of *Achai* and crossbred cows under different management systems.

| <i>Achai</i> cows | Levels                            | SF                      | RFS                     | PFS                      | Mean                    |
|-------------------|-----------------------------------|-------------------------|-------------------------|--------------------------|-------------------------|
| BCS               | <2.5                              | 1.56±0.63 <sup>c</sup>  | 2.81±0.88 <sup>b</sup>  | 3.18±0.58 <sup>a</sup>   | 2.51±0.62               |
|                   | >2.5                              | 1.90±0.48 <sup>c</sup>  | 3.17±0.63 <sup>b</sup>  | 4.18±0.47 <sup>a</sup>   | 3.08±0.47               |
|                   | <i>P-value</i>                    | 0.30                    | 0.06                    | 0.04                     | 0.05                    |
| Parity            | 1 <sup>st</sup> & 2 <sup>nd</sup> | 2.87±1.27 <sup>c</sup>  | 3.78±0.16 <sup>b</sup>  | 4.43±1.22 <sup>a</sup>   | 3.69±1.56               |
|                   | 3 <sup>rd</sup> & 4 <sup>th</sup> | 2.79±0.94 <sup>c</sup>  | 3.89±0.14 <sup>b</sup>  | 4.34±1.06 <sup>a</sup>   | 3.67±0.98               |
|                   | 5 <sup>th</sup> & 6 <sup>th</sup> | 2.92±0.82 <sup>c</sup>  | 3.72±0.18 <sup>b</sup>  | 4.61±0.87 <sup>a</sup>   | 3.75±1.25               |
|                   | <i>P-value</i>                    | 0.10                    | 0.06                    | 0.08                     | 0.14                    |
| Season            | Summer                            | 2.50±0.16 <sup>c</sup>  | 3.03±0.26 <sup>b</sup>  | 3.80±0.76 <sup>a</sup>   | 3.21±0.54               |
|                   | Winter                            | 2.01±0.11 <sup>bc</sup> | 2.37±0.52 <sup>b</sup>  | 3.22±0.51 <sup>a</sup>   | 2.51±0.48               |
|                   | <i>P-value</i>                    | 0.09                    | 0.04                    | 0.02                     | 0.04                    |
| Age               | < 4 years                         | 2.32±0.61 <sup>c</sup>  | 3.31±0.16 <sup>b</sup>  | 3.93±1.06 <sup>a</sup>   | 3.18±0.74               |
|                   | 4-6 years                         | 2.51±0.37 <sup>c</sup>  | 3.18±0.22 <sup>b</sup>  | 4.03±0.73 <sup>a</sup>   | 3.24±0.53               |
|                   | 7-8 years                         | 2.28±0.28 <sup>c</sup>  | 2.95±0.20 <sup>b</sup>  | 3.88±0.50 <sup>a</sup>   | 3.03±0.39               |
|                   | <i>P-value</i>                    | 0.16                    | 0.08                    | 0.18                     | 0.32                    |
| Crossbred cows    |                                   |                         |                         |                          |                         |
| BCS               | <2.5                              | *                       | 4.39±0.77 <sup>b</sup>  | 5.74±1.68 <sup>a</sup>   | 5.06±0.94               |
|                   | >2.5                              | *                       | 4.58±0.82 <sup>b</sup>  | 5.62±0.53 <sup>a</sup>   | 5.10±1.35               |
|                   | <i>P-value</i>                    |                         | 0.09                    | 0.14                     | 0.07                    |
| Parity            | 1 <sup>st</sup> & 2 <sup>nd</sup> | *                       | 3.95±1.06 <sup>bB</sup> | 4.62±0.95 <sup>aB</sup>  | 4.28±1.74 <sup>B</sup>  |
|                   | 3 <sup>rd</sup> & 4 <sup>th</sup> | *                       | 5.41±1.33 <sup>bA</sup> | 6.35±1.72 <sup>aA</sup>  | 5.88±1.26 <sup>A</sup>  |
|                   | 5 <sup>th</sup> & 6 <sup>th</sup> | *                       | 3.24±1.08 <sup>bC</sup> | 4.41±1.24 <sup>aBC</sup> | 3.82±0.95 <sup>BC</sup> |
|                   | <i>P-value</i>                    |                         | 0.03                    | 0.00                     | 0.02                    |
| Season            | Summer                            | *                       | 4.30±0.36 <sup>b</sup>  | 5.80±0.31 <sup>a</sup>   | 5.05±0.44               |
|                   | Winter                            | *                       | 3.51±0.21 <sup>b</sup>  | 4.43±0.47 <sup>a</sup>   | 3.97±0.23               |
|                   | <i>P-value</i>                    |                         | 0.04                    | 0.00                     | 0.02                    |
| Age               | < 4 years                         | *                       | 4.07±0.92 <sup>B</sup>  | 4.62±0.95 <sup>B</sup>   | 4.34±1.86 <sup>B</sup>  |
|                   | 4-6 years                         | *                       | 5.53±1.06 <sup>A</sup>  | 6.15±1.72 <sup>A</sup>   | 5.84±2.43 <sup>A</sup>  |
|                   | 7-8 years                         | *                       | 3.41±1.10 <sup>C</sup>  | 4.41±1.24 <sup>BC</sup>  | 3.91±1.37 <sup>BC</sup> |
|                   | <i>P-value</i>                    |                         | 0.02                    | 0.03                     | 0.03                    |

Significantly different means at  $P < 0.05$  within rows are expressed with small alphabets whereas means if significantly different at  $P < 0.05$  within columns are expressed with capital alphabets. \* Data regarding crossbred cows was not available in state farms. SF=State farms, RFS=Rural traditional farming system, PFS=Rural progressive farming systems

#### *Factors affecting the lactation length of Achai and crossbred cows under different management systems*

The effect of body condition score, parity, season and age on lactation length (LL) of *Achai* and crossbred cows in state farms (SF), rural traditional farming system (RFS) and rural progressive farming system (PFS) in Dir-Kohistan mountains of Hindukush range is presented in the Table 3.4. BCS significantly ( $P < 0.05$ ) affect the LL of *Achai* cows under all farming systems with longer lactation period in good condition (BCS>2.5) cows reared under PFS. Calving season also had significant ( $P < 0.05$ ) effect on LL with longer period in *Achai* cows calved in summer season under PFS and RFS.

The effect of parity and age on same trait of *Achai* cows was not significant ( $P > 0.05$ ). In crossbred cows,

parity, season and age had significant ( $P < 0.05$ ) effect on LL with longer period observed in 1<sup>st</sup> and 2<sup>nd</sup> parity group and age group of less than 4 years in both RFS and PFS and summer calvers in RFS. The effect of BCS on lactation length of crossbred cows was not significant ( $P < 0.05$ ).

#### *Factors affecting the lactation milk yield of Achai and crossbred cows under different management systems*

The effect of body condition score, parity, season and age on lactation yield (LY) of *Achai* and crossbred cows in state farms (SF), rural traditional farming system (RFS) and rural progressive farming system (PFS) in Hindukush mountains of Dir-Kohistan is presented in the Table 3.5. BCS significantly ( $P < 0.05$ ) affect the LY of *Achai* cows under all farming systems.

Higher lactation yield was observed in cows with body condition score more than 2.5. Calving season also had significant ( $P < 0.05$ ) effect on LY with greater yield in *Achai* cows calved in summer season. The effect of parity and age on same trait of *Achai* cows was not significant ( $P < 0.05$ ).

In crossbred cows, parity, season and age had significant ( $P < 0.05$ ) effect on LY with greater yield observed in 3<sup>rd</sup> and 4<sup>th</sup> parity group, summer calvers and age group of 4-6 years. The effect of BCS on lactation yield of crossbred cows was not significant ( $P < 0.05$ ).

**Table 3.4.** Effect of BCS, parity, season and age on lactation length (days) of *Achai* and crossbred cows under different management systems.

| <i>Achai</i> cows     | Levels                            | SF                         | RFS                         | PFS                        | Mean                       |
|-----------------------|-----------------------------------|----------------------------|-----------------------------|----------------------------|----------------------------|
| BCS                   | <2.5                              | 186.37±12.74               | 190.73±13.05                | 198.78±14.04               | 191.96±8.32                |
|                       | >2.5                              | 215.33±13.07               | 220.99±13.03                | 234.32±14.06               | 223.54±5.24                |
|                       | <i>P-value</i>                    | 0.00                       | 0.00                        | 0.01                       | 0.02                       |
| Parity                | 1 <sup>st</sup> & 2 <sup>nd</sup> | 234.31±7.87                | 213.67±4.54                 | 225.33±11.37               | 224.43±8.41                |
|                       | 3 <sup>rd</sup> & 4 <sup>th</sup> | 218.44±6.33                | 221.46±5.72                 | 215.62±09.32               | 218.17±11.90               |
|                       | 5 <sup>th</sup> & 6 <sup>th</sup> | 211.73±8.36                | 228.35±5.08                 | 239.45±13.58               | 229.84±6.24                |
|                       | <i>P-value</i>                    | 0.14                       | 0.21                        | 0.12                       | 0.26                       |
| Season                | Summer                            | 236.82±14.32 <sup>bc</sup> | 241.91±11.85 <sup>b</sup>   | 259.72±09.11 <sup>a</sup>  | 249.08±12.56               |
|                       | Winter                            | 221.35±12.51               | 223.41±15.55                | 232.35±09.21               | 225.37±13.43               |
|                       | <i>P-value</i>                    | 0.07                       | 0.04                        | 0.03                       | 0.04                       |
| Age                   | < 4 years                         | 232.56±12.24 <sup>c</sup>  | 244.33±14.22 <sup>ab</sup>  | 247.71±13.45 <sup>a</sup>  | 241.53±15.77               |
|                       | 4-6 years                         | 228.87±09.46 <sup>c</sup>  | 241.56±12.11 <sup>b</sup>   | 254.84±15.33 <sup>a</sup>  | 245.75±09.26               |
|                       | 7-8 years                         | 219.98±08.68 <sup>c</sup>  | 245.38±15.81 <sup>ab</sup>  | 239.22±11.12 <sup>b</sup>  | 234.86±11.48               |
|                       | <i>P-value</i>                    | 0.10                       | 0.08                        | 0.16                       | 0.09                       |
| <b>Crossbred cows</b> |                                   |                            |                             |                            |                            |
| BCS                   | <2.5                              | *                          | 246.52±7.13 <sup>b</sup>    | 285.47±6.07 <sup>a</sup>   | 265.99±9.23                |
|                       | >2.5                              | *                          | 253.43±5.12 <sup>b</sup>    | 290.82±8.23 <sup>a</sup>   | 272.12±7.78                |
|                       | <i>P-value</i>                    |                            | 0.06                        | 0.07                       | 0.12                       |
| Parity                | 1 <sup>st</sup> & 2 <sup>nd</sup> | *                          | 266.53±09.14 <sup>bA</sup>  | 281.11±09.14 <sup>aA</sup> | 273.93±15.41 <sup>A</sup>  |
|                       | 3 <sup>rd</sup> & 4 <sup>th</sup> | *                          | 247.26±13.11 <sup>bBC</sup> | 268.37±17.51 <sup>ab</sup> | 257.81±10.54 <sup>BC</sup> |
|                       | 5 <sup>th</sup> & 6 <sup>th</sup> | *                          | 251.71±11.21 <sup>B</sup>   | 261.02±15.00 <sup>BC</sup> | 258.36±17.21 <sup>B</sup>  |
|                       | <i>P-value</i>                    |                            | 0.02                        | 0.01                       | 0.04                       |
| Season                | Summer                            | *                          | 262.36±16.71 <sup>b</sup>   | 286.82±17.26 <sup>a</sup>  | 269.59±15.60               |
|                       | Winter                            | *                          | 241.47±11.47 <sup>b</sup>   | 275.35±15.52 <sup>a</sup>  | 253.41±13.42               |
|                       | <i>P-value</i>                    |                            | 0.00                        | 0.07                       | 0.04                       |
| Age                   | < 4 years                         | *                          | 269.53±09.14 <sup>A</sup>   | 284.21±09.91 <sup>A</sup>  | 276.87±14.59               |
|                       | 4-6 years                         | *                          | 253.32±11.25 <sup>B</sup>   | 270.53±12.32 <sup>B</sup>  | 261.92±13.26               |
|                       | 7-8 years                         | *                          | 245.71±08.33 <sup>C</sup>   | 258.74±11.10 <sup>C</sup>  | 252.22±09.50               |
|                       | <i>P-value</i>                    |                            | 0.04                        | 0.03                       | 0.05                       |

Significantly different means at  $P < 0.05$  within rows are expressed with small alphabets whereas means if significantly different at  $P < 0.05$  within columns are expressed with capital alphabets. \* Data regarding crossbred cows was not available in state farms. SF=State farms, RFS=Rural traditional farming system, PFS=Rural progressive farming systems.

**Table 3.5.** Effect of BCS, parity, season and age on lactation yield (L) of *Achai* and crossbred cows under different management systems.

| <i>Achai</i> cows | Levels                            | SF                         | RFS                       | PFS                        | Mean         |
|-------------------|-----------------------------------|----------------------------|---------------------------|----------------------------|--------------|
| BCS               | <2.5                              | 390.87±13.76 <sup>c</sup>  | 438.50±19.34 <sup>b</sup> | 629.56±17.62 <sup>a</sup>  | 452.97±21.77 |
|                   | >2.5                              | 516.77±12.43 <sup>c</sup>  | 719.10±22.64 <sup>b</sup> | 978.73±22.97 <sup>a</sup>  | 738.20±12.45 |
|                   | <i>P-value</i>                    | 0.04                       | 0.01                      | 0.00                       | 0.00         |
| Parity            | 1 <sup>st</sup> & 2 <sup>nd</sup> | 624.88±14.37 <sup>c</sup>  | 719.99±18.99 <sup>b</sup> | 996.75±17.22 <sup>a</sup>  | 780.54±15.37 |
|                   | 3 <sup>rd</sup> & 4 <sup>th</sup> | 608.22±16.08 <sup>c</sup>  | 727.21±34.90 <sup>b</sup> | 928.76±23.64 <sup>a</sup>  | 754.73±21.40 |
|                   | 5 <sup>th</sup> & 6 <sup>th</sup> | 645.32±13.06 <sup>c</sup>  | 745.55±28.87 <sup>b</sup> | 1007.40±15.48 <sup>a</sup> | 799.42±9.53  |
|                   | <i>P-value</i>                    | 0.22                       | 0.16                      | 0.09                       | 0.33         |
| Season            | Summer                            | 530.67±53.42 <sup>c</sup>  | 748.71±25.36 <sup>b</sup> | 961.87±56.02 <sup>a</sup>  | 767.08±27.61 |
|                   | Winter                            | 444.71±45.46 <sup>bc</sup> | 540.56±43.91 <sup>b</sup> | 747.44±39.26 <sup>a</sup>  | 577.57±16.36 |
|                   | <i>P-value</i>                    | 0.08                       | 0.03                      | 0.02                       | 0.03         |
| Age               | < 4 years                         | 539.53±23.59 <sup>c</sup>  | 808.73±18.99 <sup>b</sup> | 973.50±13.17 <sup>a</sup>  | 773.92±14.21 |
|                   | 4-6 years                         | 574.46±18.30 <sup>c</sup>  | 768.16±34.90 <sup>b</sup> | 1027.05±23.73 <sup>a</sup> | 789.89±19.60 |
|                   | 7-8 years                         | 501.55±20.54 <sup>c</sup>  | 723.87±28.87 <sup>b</sup> | 958.17±15.50 <sup>a</sup>  | 717.86±26.13 |



| <i>Achai</i> cows | Levels                            | SF   | RFS                         | PFS                         | Mean                       |
|-------------------|-----------------------------------|------|-----------------------------|-----------------------------|----------------------------|
|                   | <i>P-value</i>                    | 0.06 | 0.11                        | 0.17                        | 0.06                       |
| Crossbred cows    |                                   |      |                             |                             |                            |
| BCS               | <2.5                              | *    | 1079.52±19.51 <sup>b</sup>  | 1493.40±14.32 <sup>a</sup>  | 1286.46±20.21              |
|                   | >2.5                              | *    | 1158.74±23.31 <sup>b</sup>  | 1629.80±27.54 <sup>a</sup>  | 1394.27±29.50              |
|                   | <i>P-value</i>                    |      | 0.07                        | 0.13                        | 0.60                       |
| Parity            | 1 <sup>st</sup> & 2 <sup>nd</sup> | *    | 1030.95±13.73 <sup>bB</sup> | 1298.95±11.95 <sup>aB</sup> | 1164.95±19.55 <sup>B</sup> |
|                   | 3 <sup>rd</sup> & 4 <sup>th</sup> | *    | 1336.27±10.95 <sup>bA</sup> | 1648.27±09.61 <sup>aA</sup> | 1492.27±13.26 <sup>A</sup> |
|                   | 5 <sup>th</sup> & 6 <sup>th</sup> | *    | 826.27±14.96 <sup>bC</sup>  | 1151.01±15.11 <sup>aC</sup> | 988.64±21.78 <sup>C</sup>  |
|                   | <i>P-value</i>                    |      | 0.02                        | 0.00                        | 0.01                       |
| Season            | Summer                            | *    | 1126.67±23.42 <sup>b</sup>  | 1660.67±24.42 <sup>a</sup>  | 1393.67±24.61              |
|                   | Winter                            | *    | 845.71±13.26 <sup>b</sup>   | 1213.25±19.26 <sup>a</sup>  | 1009.83±15.48              |
|                   | <i>P-value</i>                    |      | 0.01                        | 0.03                        | 0.04                       |
| Age               | < 4 years                         | *    | 1096.98±19.51 <sup>B</sup>  | 1313.28±21.44 <sup>B</sup>  | 1205.13±21.37 <sup>B</sup> |
|                   | 4-6 years                         | *    | 1399.07±17.50 <sup>A</sup>  | 1645.30±15.20 <sup>A</sup>  | 1522.18±16.22 <sup>A</sup> |
|                   | 7-8 years                         | *    | 837.87±12.22 <sup>C</sup>   | 1141.04±25.34 <sup>C</sup>  | 989.45±20.68 <sup>C</sup>  |
|                   | <i>P-value</i>                    |      | 0.02                        | 0.00                        | 0.02                       |

Significantly different means at  $P < 0.05$  within rows are expressed with small alphabets whereas means if significantly different at  $P < 0.05$  within columns are expressed with capital alphabets. \* Data regarding crossbred cows was not available in state farms. SF=State farms, RFS=Rural traditional farming system, PFS=Rural progressive farming systems

**Table 3.6.** Effect of BCS, parity, season and age on calf birth weight (kg) of *Achai* and crossbred cows under different management systems.

| <i>Achai</i> cows | Levels                            | SF                       | RFS                      | PFS                     | Mean       |
|-------------------|-----------------------------------|--------------------------|--------------------------|-------------------------|------------|
| BCS               | <2.5                              | 15.27±0.26 <sup>c</sup>  | 15.59±0.15 <sup>b</sup>  | 16.56±0.04 <sup>a</sup> | 15.80±0.22 |
|                   | >2.5                              | 15.42±0.43 <sup>bc</sup> | 15.50±0.11 <sup>ab</sup> | 16.36±0.93 <sup>a</sup> | 15.68±0.09 |
|                   | <i>P-value</i>                    | 0.06                     | 0.10                     | 0.21                    | 0.11       |
| Parity            | 1 <sup>st</sup> & 2 <sup>nd</sup> | 15.41±0.15               | 16.20±2.27               | 16.43±1.31              | 16.01±0.15 |
|                   | 3 <sup>rd</sup> & 4 <sup>th</sup> | 16.95±2.14               | 15.58±1.72               | 15.97±2.75              | 16.16±0.62 |
|                   | 5 <sup>th</sup> & 6 <sup>th</sup> | 15.78±3.27               | 15.73±2.24               | 16.04±0.98              | 15.85±0.75 |
|                   | <i>P-value</i>                    | 0.06                     | 0.06                     | 0.11                    | 0.26       |
| Season            | Summer                            | 16.04±0.38               | 15.84±0.46               | 16.24±0.32              | 16.04±0.40 |
|                   | Winter                            | 15.95±0.35               | 16.15±0.67               | 15.89±0.49              | 15.99±0.53 |
|                   | <i>P-value</i>                    | 0.22                     | 0.13                     | 0.06                    | 0.35       |
| Age               | < 4 years                         | 16.31±0.15               | 16.18±2.32               | 16.03±1.78              | 16.17±1.54 |
|                   | 4-6 years                         | 16.15±2.14               | 16.24±1.50               | 15.97±2.33              | 16.12±1.70 |
|                   | 7-8 years                         | 15.98±3.27               | 16.53±2.28               | 16.24±1.64              | 16.25±2.04 |
|                   | <i>P-value</i>                    | 0.08                     | 0.17                     | 0.19                    | 0.33       |
| Crossbred cows    |                                   |                          |                          |                         |            |
| BCS               | <2.5                              | *                        | 16.41±0.37               | 15.94±0.57              | 16.17±1.04 |
|                   | >2.5                              | *                        | 16.36±0.24               | 16.04±0.63              | 16.20±0.88 |
|                   | <i>P-value</i>                    |                          | 0.23                     | 0.13                    | 0.47       |
| Parity            | 1 <sup>st</sup> & 2 <sup>nd</sup> | *                        | 15.93±0.14 <sup>b</sup>  | 16.73±0.43 <sup>a</sup> | 16.33±0.82 |
|                   | 3 <sup>rd</sup> & 4 <sup>th</sup> | *                        | 15.62±0.58 <sup>b</sup>  | 16.92±0.84 <sup>a</sup> | 16.27±0.56 |
|                   | 5 <sup>th</sup> & 6 <sup>th</sup> | *                        | 16.07±0.36 <sup>b</sup>  | 16.71±0.66 <sup>a</sup> | 16.39±0.77 |
|                   | <i>P-value</i>                    |                          | 0.30                     | 0.15                    | 0.19       |
| Season            | Summer                            | *                        | 16.04±0.51               | 16.21±0.45              | 16.12±0.48 |
|                   | Winter                            | *                        | 15.42±0.23               | 15.55±0.71              | 15.48±0.47 |
|                   | <i>P-value</i>                    |                          | 0.03                     | 0.04                    | 0.05       |
| Age               | < 4 years                         | *                        | 15.88±0.47               | 16.53±0.70              | 16.20±0.86 |
|                   | 4-6 years                         | *                        | 16.32±0.23               | 16.87±0.49              | 16.55±0.59 |
|                   | 7-8 years                         | *                        | 16.13±0.77               | 16.64±0.31              | 16.38±0.42 |
|                   | <i>P-value</i>                    |                          | 0.11                     | 0.27                    | 0.36       |

Significantly different means at  $P < 0.05$  within rows are expressed with small alphabets whereas means if significantly different at  $P < 0.05$  within columns are expressed with capital alphabets. \* Data regarding crossbred cows was not available in state farms. SF=State farms, RFS=Rural traditional farming system, PFS=Rural progressive farming systems.

### *Factors affecting the calf birth weight of Achai and crossbred cows under different management systems*

The effect of body condition score, parity, season and age on calf birth weight (CBW) of *Achai* and crossbred cows in state farms (SF), rural traditional farming system (RFS) and rural progressive farming system (PFS) in Hindukush mountains of Dir-Kohistan is presented in the Table 3.6. BCS, parity, calving season and age had no significant ( $P>0.05$ ) effect on the CBW of *Achai* cows under all farming systems. In crossbred cows, calving season had significant ( $P<0.05$ ) effect on CBW. Heavier calves were born in summer season under both RFS and PFS. The effect of BCS, parity and age on CBW of crossbred cows was not significant ( $P<0.05$ ).

### **Discussion**

#### *Overall productive performance of Achai and crossbred cows*

Crossbreeding of indigenous cows with exotic breeds improved productive performance (Mulugeta and Belayneh, 2013). Previous study (Buckley *et al.*, 2014) has demonstrated remarkable benefits of additive and non-additive genetic effects of crossbreeding on overall dairy cattle performance. Several studies reported significantly higher daily milk yield (Wakchaure *et al.*, 2015), longer lactation length (Rokonuzzaman *et al.* 2009), greater lactation yield (Mulugeta and Belayneh, 2013) and calf birth weight (Prendiville *et al.*, 2010) in crossbred cows as compared to local cows. Mean daily milk yield recorded in present study was significantly ( $P<0.001$ ) lower than crossbred cows. For *Achai* cows, Hayazuddin *et al.* (2014) reported slightly lower ( $2.81\pm 0.12$  litres/day) DMY than  $3.55\pm 0.77$  litres in the present study. Several factors like animal diseases, unavailability of feeds in terms of quality and quantity, poor management are responsible for low production of local cattle (Hassan *et al.*, 2016). The extended lactation length observed in crossbred cows as compared to local cows confirms the findings of Hayazuddin *et al.* (2014) regarding improvement in lactation length of local cattle through crossbreeding. However, the observed lactation period in both *Achai* as well as Jersey crossbred cows reported in this study is shorter than the

recommended value of 305 days (Habib *et al.*, 2003) which need improvement (De Vries, 2000). Lactation milk yield of *Achai* cow observed in this study was significantly lower than crossbred. In another study, Hayazuddin *et al.* (2014) recorded  $813.07\pm 113.39$  litres in a lactation period of  $263.14\pm 24.53$  days. Low lactation milk yield in local cattle are also reported by Amin, (2007) and Mulugeta and Belayneh, (2013) as compared to their crossbred with high yielding dairy cows. Average birth weight of crossbred calves ( $16.29\pm 0.16$ ) was significantly higher than *Achai* calves ( $15.20\pm 0.17$ ). *Achai* is light weight multipurpose breed reared under extensive production system (Khan, D., 2004) as compared to Jersey breed. Several researchers (Sørensen *et al.*, 2008; Prendiville *et al.*, 2010) worked on crossbreeding to improve calf birth weight in addition to other long term economic traits. The aim of dairy producers in practicing crossbreeding is to improve calving ease and to produce calf with optimum weight for cow to birth (Weigel and Barlass, 2003). In other perspective, many researchers concluded that crossbreeding has significant impact in reducing calving difficulties (Heins *et al.*, 2006a). Dhakal, *et al.*, (2013) also reported less calving difficulty in crossbred as compared to purebred due to calf birth weight ratio to dam size.

#### *Effect of farming systems on the productive performance of Achai and crossbred cows*

Very few studies described the impact of housing systems on the performance of dairy cattle (Fregonesi and Leaver, 2002; Klopchich *et al.*, 2007; Simensen *et al.*, 2010). However, it is clearly evident that improving animal welfare through proper farming practices and housing conditions considerably improves dairy cattle performance (Lambertz, *et al.*, 2014). Several researchers (Fregonesi and Leaver, 2001; de Vries *et al.*, 2011) has determined direct relationship between housings conditions and milk production. Different housing systems increase or decrease the impact of various environmental conditions which directly affect production of dairy cattle (Bohmanova *et al.*, 2007). Simensen *et al.*, (2010) reported significant impact of tie and free-stall herding on milk production.

On the other hand, improved milk production in individual cows was observed when cows were moved from loose housing barns to tie stall barns (Hovinen *et al.* (2009). Several factors including more attention in milking practices (Hovinen *et al.* 2009), nutrition (Brun-Lafleur *et al.* 2010), bedding material effect on various genotypes (Mark and Lassen, 2007), housing design (Hristov *et al.*, 2014) in tie-stall are responsible for higher milk production. In modern dairying, 305 days in lactation is considered as standard. However, such value is not practicable in rural and conventional dairy farmers where lactation length considerable extends or reduces from normal values (Msangi *et al.*, 2005). Regarding farming practices, (Eicher, *et al.*, 2013) analysed that improved animal welfare observed in grazing system compared to zero grazing and cows housed in straw yards compared to free stall significantly affected dairy cattle performance including extended milk yield duration and animal health. Lehmann *et al.*, (2017) observed prematurely dry off due to insufficient feeding resulting short lactation length in dairy cattle. Albarrán-Portillo and Pollott, (2011) reported significant effect of general management on milk production and lactation period in dairy animals. Calf birth weight is complex phenomenal trait associated with genetic and non-genetic factors (Kamal *et al.*, 2014). Besides genetic factors, nutritional limitations significantly affect embryo and foetus development during gestation (Funston and Summers, 2013). Several researchers (Zhang *et al.*, 2002; Symonds *et al.*, 2010) demonstrated environmental and nutritional effect on calf birth. Other studies (Lundborg *et al.*, 2003; Swali and Wathes, 2006) reported dam morphometrics as source of influence on birth weight of their calves. Similarly, early breeding management in heifers has potential effect on birth size and weight of calves (González-Recio *et al.*, 2014).

#### *Factors affecting the daily milk yield of Achai and crossbred cows under different management systems*

The significant effect of BCS on DMY of *Achai* cows in PFS with higher production in cows with BCS of more than 2.5 may be due better nutrition in PFS. It has been observed that optimum nutrition result in better

BCS (reflected by the degree of subcutaneous fatness) is correlated with energy balance and enhanced cow productivity (Gaillard, C. 2016). Nickerson (1995) observed strong association of milk production and body frame of cattle and revealed significant support of serum precursors for milk synthesis. Higher milk yield in good condition cows have also been reported in zebu and crossbred cows (Samarutel *et al.*, 2009; Singh *et al.*, 2015) as well as high producing cows (Roche *et al.*, 2007). In crossbred cows, parity had significant effect on DMY with higher yield observed in 3<sup>rd</sup> and 4<sup>th</sup> parity group under all management systems. Sum higher yield in crossbred cows might be due to complete development and adoption of lactation physiology (Nickerson, 1995). Devery-Pocius and Larson (1983) further explained that better immunological adaptation in multiparous cows is responsible for increased milk production as compared to primiparous cows which yet to be adopted for certain physiological processes. The incomplete structural growth of udder and teats of cow is another reason for low in milk production in primiparous cows. According to Ihsanullah and Qureshi, (2019), the lower milk yield in crossbred cows of early and late parities compared to mid parities cows may possibly be due to the consumptions of more nutrients for maintenance instead of production (Ihsanullah and Qureshi, 2019).

Several studies (Lakshmi, *et al.*, 2009; Islam and Kundu, 2011) have also reported significant effect of parity on DMY in crossbred cows from various regions of the world. Calving season had significant effect on DMY of *Achai* and crossbred cows with higher production in both breed cows calved in summer season under PFS while the effect of calving season on DMY of both breed was non-significant under state farming system. The significantly higher DMY of *Achai* and crossbred cows calved in RFS and PFS during summer season may be due to better feed availability in the summer season (Ihsanullah and Qureshi, 2019) as the winter season in the study area has harsh environmental conditions, severe shortage of fodders and the animals mostly rely on roughages (Saleem *et al.*, 2012). Seasonal effect on DMY with higher production during the months of higher fodder

availability has also been reported by Lakshmi, *et al.*, (2010) and Das *et al.*, (2011). The significant increase in DMY of crossbred cows at the age of 4-6 years might be due to maturity of cattle body to support lactation stress while reduction in daily yield of older animals is due to ageing affect (Ihsanullah and Qureshi, 2019). The number and activity of alveoli in mammary gland reduces with age which causes reduction in milk yield (Auldism, *et al.* 2007). Brscic *et al.*, (2015) revealed that initial increase in serum protein and glucose concentrations up to certain age followed by gradual decrease results variation in milk yield. Goff, (2008) further explained the phenomena of reduced milk production in young cows characterized by immunological suppression due to lack of adaptation to physiological stress leading to decline in several protein fractions.

The absence of age effect on DMY of *Achai* cows may be contributed due to breed specific characters. Local breeds which are usually low producers due to multipurpose selection are adapted without particular attention for milk production (Abera, 2016). Another reason for the absence of age effect on production performance may be such level of low production that further couldn't be affected by age (Amasiab *et al.*, 2011). Constantly low milk production without any significant effect of age has also been reported by Kurtu, (2003) and Gurmesa *et al.*, (2012).

#### *Factors affecting the lactation length of Achai and crossbred cows under different management systems*

The significantly longer LL in good condition (BCS >2.5) *Achai* cows may be due to improved nutrition reflected by degree of subcutaneous fatness providing significant amount of serum precursors to support milk production for longer period (Gillard, C., 2016). In study area, farmers don't follow the standard 305-day lactation length rather milk the cows till its production capacity which is greater in good condition cows. Janu's *et al.* (2007) observed that length of lactation period significantly depend upon postpartum body fat reserves. The authors noted that cows gained greater body reserves during dry period had longer lactation period.

Longer LL has also been reported in local and crossbred cows with good body condition scores by Samarutel *et al.*, (2009) and Singh *et al.*, (2015). The significant effect of parity on LL of the crossbred cows with longer LL in 1<sup>st</sup> and 2<sup>nd</sup> parity group may be due to the readily available body reserves of primiparous cows enabling them to produce milk for longer time (Ratnayake *et al.*, 1998).

Furthermore, better nutrition improves body condition of heifers which results in milk production for longer durations after parturition as compared to primiparous cows (Ihsanullah and Qureshi, 2019). Significant effect of parity on LL with longer LL in early parity cows have also been reported by many authors (Ahmad *et al.*, 2007; Kolver *et al.*, 2007; Phyn *et al.*, 2008). The significant effect of calving season on LL of *Achai* as well as crossbred cows with longer LL in summer calvers seems to be due to availability of quality fodders during summer season (Amasiab *et al.*, 2008; Ihsanullah and Qureshi, 2019). Several studies revealed extended lactations with provision of quality feed (Auldism *et al.*, 2007; Butler, *et al.*, 2010; Kolver, *et al.*, 2007; Phyn *et al.*, 2008; Grainger *et al.*, 2009). The non-significant effect of age on LL of *Achai* cows is consistent with the early reports of Habib *et al.*, (2010). However, the significant effect of age on LL in crossbred cows with longer LL in the age group <4 years and a decline afterward seems to be the development of size and activity of mammary glands up to this age followed by a decline in the number and progress of alveoli in mammary glands with the advancement of age of the cattle (Nickerson, 1995; Amasiab *et al.*, 2011). Several secondary factors including imbalance metabolism (Meikle *et al.*, 2004), DMI (Maekawa *et al.*, 2002) and physiological condition (Ingvarsten, 1994) has been strongly associated with age which causes variation in productive performance of dairy cattle. Some studies confirm gradual decrease in milk production of crossbred cows due to shorter lactation period with advancement up to 6 years (Mohamed, 2004; Kurtu, 2003). The effect of age on productive performance including LL has also been reported by Brscic *et al.*, (2015), Abera, (2016) and Gurmesa, *et al.*, (2012).

*Factors affecting the lactation yield of Achai and crossbred cows under different management systems*

The significant effect of BCS on LY of *Achai* cows with higher production in cows with BCS more than 2.5 may be due to higher DMY for longer lactation period (Gillard, C., 2016) because of better nutrition and management practices in PFS. Better nutrition associated with body condition provides significant amount of serum precursors to support higher milk yield for longer period (Nickerson (1995). According to Van Kneysel, *et al.*, (2014) body reserves significantly supports lactogenesis for the entire lactation thus cows with good body condition results higher yield per lactation. The effect of BCS on LY with higher production in good condition cows has also been reported by Dechow *et al.* (2002), Bewley and Schutz (2008) and Loker *et al.* (2012). The completion of mammary gland development with experiencing lactations might be the reason of higher LY in crossbred cows of 3<sup>rd</sup> and 4<sup>th</sup> parity group. In addition, the immunological adaptation to various physiological processes increases up to certain age with the experience of lactations which results higher LY in mid parities as compared to primiparous or older cows (Devery-Pocius and Larson, 1983). Bajwa *et al.*, (2004) reported that LY increases up to 5<sup>th</sup> parity in Sahiwal cows followed by gradual decrease in later parities. The effect of parity on LY of cattle with higher yield up to 6<sup>th</sup> parity has also been reported in Pakistan (Dahlin, 1998; Ahmad *et al.*, 2004) and other countries (Dhumal *et al.*, 1989; Deshpande and Sakhare, 1984).

The availability of quality feed during favorable climatic conditions at the initiation of lactation stage results higher daily yield (Ihsanullah and Qureshi, 2019) for longer lactation period (Amasiab, *et al.*, 2008) might be reason for higher lactation yield in summer calved cows. The nutrients availability supports the provision of volatile fatty acids and serum glucose for milk synthesis. The effect of calving season with higher yield in favorable climatic conditions due to better fodder availability has been reported by Bajwa *et al.*, (2004), Dahlin (1998) and Talbott (1994) in Pakistan as well as other countries by Sorenson, *et al.*, (2008), Jankowska, *et al.*, (2012)

and Steri, *et al.*, (2012). The significant effect of age on lactation yield with higher production in 4-6 years age cows has been reported by Briscic *et al.*, (2015) and Abera, (2016). The conclusion of these authors for higher milk yield per lactation was the development of size and activity of alveoli in mammary gland with advancement in age. According to Ihsanullah and Qureshi (2019) maturity of cattle significantly improves the production performance of dairy cattle. The immunological suppression in young and old cows also significantly affect milk yield of dairy cattle (Goff, 2008). Some studies confirmed the gradual decrease in lactation yield of crossbred cows particularly after 6 years (Mohammed, 2004; Kurtu, 2003). Gurmessa, *et al.*, (2012) concluded similar reports regarding age effect on production traits of dairy cattle. To justify the variation in milk yield due to age, Briscic *et al.*, (2015) observed significant increase in serum protein and glucose concentrations up to five years which then gradually decreased.

*Factors affecting the calf birth weight (CBW) of Achai and crossbred cows under different management systems*

The non-significant effect of BCS on CBW of *Achai* and crossbred cows might be due to greater proportion of nutrient partitioning for body maintenance during dry period which is common characteristic of all low producers (Reference). Furthermore, improving cows BCS during pre-partum nutrition planning has very little or no effect on CBW (Renquest *et al.*, 2005). The non-significant effect of body condition of cattle at calving associated with nutrition has also been reported in many studies (Mulliniks *et al.*, 2015; Lake *et al.*, 2005). Parity had no significant ( $P>0.05$ ) effect on the CBW of *Achai* and crossbred cows under all farming systems. The non-significant effect of parity on CBW has been reported by Addisu *et al.*, (2010), Melaku *et al.*, (2011a) and Tesfa, *et al.*, (2016). Despite of significantly positive correlation between age of cow and internal and external pelvic measurements, Bures, *et al.*, (2008) found insignificant effect of parity on weight of calves born. Similar results were reported by Nogalski (2003) for HF calves. The significant effect of calving season on CBW of



crossbred cows with heavier calves born in summer might be due seasonal fodder availability (Renquest *et al.*, 2005) because proper climatic conditions significantly improves the nutrient composition of fodders in addition to greater quantity fodders production (Lammoglia *et al.*, 1996). Some studies revealed that concentrate supplementation during winter season have significant effect of CBW (Giday, 2001; Melaku *et al* 2011a; Almaz 2012). On the other side, restricted metabolism of pregnant cow due to fodder scarcity in extreme weather conditions may also results in lighter calf births (Kumar, *et al.*, 2017). The significant effect of calving season on CBW has also been reported by Habtamu, *et al*, (2010), Aynalem *et al.*, (2010), Demissu *et al*, (2013) and Tesfa *et al.*, (2016). However, some studies also reported non-significant effect of calving season on CBW (Getinet *et al.*, 2009; Addisu *et al.*, 2010).

### Conclusions

1. *Achai* x Jersey (crossbreds) cows had significantly better DMY, LL, MYL and CBW than pure *Achai* cows.
2. Improving management practices significantly improved the productive performance of both breeds as observed in rural progressive farming system (PFS).
3. Age and parity had no effect on productive performance of *Achai* cows.
4. BCS had no effect on productive performance of crossbred cows.
5. Economics traits of both breeds were better in summer season.
6. Adult (4-6 years age) crossbred cows had better DMY and LY in their 3<sup>rd</sup> and 4<sup>th</sup> parities while longer LL was observed in young (<4 years age) cows in its early (1<sup>st</sup> and 2<sup>nd</sup> parity group) parities.
7. Introducing *Achai* cows to state farming systems significantly affected its performance.

### References

**Abera M.** 2016. Reproductive and productive performances of crossbred and indigenous dairy cattle under rural, peri-urban and urban dairy farming systems in West Shoa Zone, Oromia, Ethiopia. (pp. 20-21).

**Addisu B, Mengistie T, Adebabay K, Getinet M, Asaminew T, Tezera M, Gebeyehu G.** 2010. Milk yield and calf growth performance of cattle under partial suckling system at Andassa Livestock Research Centre, North West Ethiopia. Livestock Research for Rural Development **22(8)**, 55-61.

**Ahmad A, Ahmad MK, Teirab B, Lutfi MA, Peters J.** 2007. Milk production and reproduction trait of different grades of zebu x Friesian crossbred under semi-arid conditions. Archives Tierz Dummerstorf **50(3)**, 240-249

**Ahmad I, Lodhi LA, Qureshi ZI, Younis M.** 2004. Studies on blood glucose, total proteins, urea and cholesterol levels in cyclic, non-cyclic and endometritic crossbred cows. Blood **24(2)**, 2-4

**Albarrán-Portillo B, Pollott GE.** 2011. Environmental factors affecting lactation curve parameters in the United Kingdom's commercial dairy herds. Arch. Med. Vet **43**, 145-153.

**Almaz B.** 2012. Genetic parameter estimation of growth and reproduction traits of Fogera cattle at Metekel Ranch, Amhara Region, Ethiopia. MSc thesis, Bahir Dar university college of agriculture and environmental science, Bahir Dar, Ethiopia. pp 108.

**Amin AA, Nahar A.** 2007. Productive and reproductive performance of Non-descript (Local) and crossbred dairy cows in coastal area of Bangladesh. Asian Journal of. Animal and Veterinary Sciences **2(1)**, 46-49.

**Annicchiarico P.** 2002. Genotype × environment interaction: Challenges and opportunities for plant breeding and cultivar recommendations. FAO Plant Production and Protection Paper 174. FAO of the United Nations, Rome.

**Auldism MJG, O'Brien D, Cole KL, Macmillan, Grainger C.** 2007. Effects of varying lactation length on milk production capacity of cows in pasture based dairying systems, Journal of Dairy Science **90**, 3234-3241.



- Aynalem H, Joshi BK, Workinehi A, Azage T, Singh A.** 2010. Genetic evaluation of Ethiopian Boran cattle and their crosses with Holstein Friesian for growth performance in central Ethiopia. *Journal of Animal Breeding and Genetics* **1(4)**, 0931-0938.
- Bajwa IR, Khan MS, Khan MA, Gondal KZ.** 2004. Environmental factors affecting milk yield and lactation length in Sahiwal cattle. *Pakistan Veterinary Journal* **24(1)**, 23-27.
- Bell MJ, Eckard RJ, Haile-Mariam M, Pryce JE.** 2013. The effect of changing cow production and fitness traits on net income and greenhouse gas emissions from Australian dairy systems. *Journal of Dairy Science* **96**, 7918-7931.
- Bewley JM, Schutz MM.** 2008. Review: an interdisciplinary review of body condition scoring for dairy cattle. *The Professional Animal Scientist* **24(6)**, 507-529.
- Bohmanova J, Misztal I, Cole JB.** 2007. Temperature humidity indices as indicators of milk production losses due to heat stress. *Journal of Dairy Science* **90**, 1947-1956.
- Brcsic M, Cozzi G, Lora I, Stefani AL, Contiero B, Ravarotto L, Gottardo F.** 2015. Short communication: Reference limits for blood analytes in Holstein late-pregnant heifers and dry cows: Effects of parity, days relative to calving, and season. *Journal of Dairy Science* **98**, 7886-7892.
- Brun-Lafleur L, Delaby L, Husson F, Favardin P.** 2010. Predicting energy  $\times$  protein interaction on milk yield and milk composition in dairy cows. *Journal of Dairy Science* **93**, 4128-4143.
- Buckley F, O'Sullivan K, Mee JF, Evans RD, Dillon P.** 2014. Relationships among milk yield, body condition, cow weight and reproduction in spring-calved Holstein-Friesians. *Journal of Dairy Science* **86**, 2308-2319.
- Bures D, Barton L, Zahradkova R, Teslik V, Fiedlerova M.** 2008. Calving difficulty as related to body weights and measurements of cows and calves in a herd of Gascon breed. *Czech Journal of Animal Sciences* **53(5)**, 187-194.
- Connor EE.** 2015. Invited review: Improving feed efficiency in dairy production: Challenges and possibilities. *Animal* **9**, 395-408.
- Das AM, Gupta D, Khan MKI, Miah G.** 2011. Effect of non-genetic factors on the productive and reproductive traits of Friesian crossbred dairy cows. *Wayamba Journal of Animal Science* **4**, 62-64.
- De Vries M, Bokkers E, Dijkstra T, van Schaik G, de Boer I.** 2011. Invited review: Associations between variables of routine herd data and dairy cattle welfare indicators. *Journal of Dairy Science* **94**, 3213-3228.
- De Vries MJ, Veerkamp RF.** 2000. Energy Balance of Dairy Cattle in Relation to Milk Production Variables and Fertility. *Journal of Dairy Science* **83(1)**, 62-69.
- Demissu H, Fekadu B, Gemedu D.** 2013. Early growth and reproductive performances of Horro cattle and their F1 Jersey crosses in and around Horro-Guduru Livestock Production and Research Center, Ethiopian Scientific Technological and Arts Research Journal **2(3)**, 134-141.
- Dhakal K, Maltecca C, Cassady JP, Baloch G, Williams CM, Washburn SP.** 2013. Calf birth weight, gestation length, calving ease, and neonatal calf mortality in Holstein, Jersey, and crossbred cows in a pasture system. *Journal of Dairy Science* **96(1)**, 690-698.
- Eicher SD, Lay D, Arthington JD, Schutz MM.** 2013. Effects of rubber flooring during the first 2 lactations on production, locomotion, hoof health, immune functions, and stress. *Journal of Dairy Science* **96(6)**, 3639-3651.

- Epaphras A, Karimuribo ED, Msellem SN.** 2004. Effect of season and parity on lactation of crossbred Ayrshire cows reared under coastal tropical climate in Tanzania. *Livestock Research for Rural Development* **(16)6**, 205-213.
- Fernández-Perea MT, Jimenez RA.** 2004. Economic weights for a selection index in Avileña purebred beef cattle. *Livestock Production Science* **89**, 223-233.
- Fetrow J, Stewart S, Eicker S, Rapnicki P.** 2007. 'Reproductive Health Programs for Dairy Herds: Analysis of Records for Assessment of Reproductive Performance' in *Current Therapy in Large Animal Theriogenology*, eds Youngquist, R.S. & Threlfall, W.R., Saunders, Philadelphia, pp. 473-489.
- Fodor I, Ozsvári L.** 2015. The evaluation of reproductive performance in dairy herds. 085
- Fregonesi JA, Leaver JD.** 2001. Behaviour, performance and health indicators of welfare for dairy cows housed in strawyard or cubicle systems. *Livestock Production Science* **68**, 205-216.
- Fregonesi JA, Leaver JD.** 2002. Influence of space allowance and milk yield level on behavior, performance and health of dairy cows housed in strawyard and cubicle systems. *Livestock Production Science* **78**, 245-257.
- Funston RN, Summers AF.** 2013. Effect of prenatal programming on heifer development. *Veterinary Clinics North American Food Animal Practitioners* **29**, 517-536.
- Gaillard C.** 2016. Extended lactation and feeding strategies in dairy cows extended lactation and feeding strategies in dairy cows. Aarhus University, Foulum, Denmark. PhD Thesis, pp. 18-23.
- Getinet M, Workneh A, Hegde BP.** 2009. Growth and reproductive performance of Ogaden cattle at Haramaya University, Ethiopia. *Ethiopian Journal of Animal Production* **9(1)**, 13-38.
- Giday Y.** 2001. Assessment of calf crop productivity and total herd life of Fogera cows at Andassa ranch, Northwestern Ethiopia. M.Sc. Thesis. Alemaya University, Alemaya, Ethiopia.
- Goff JP.** 2008. Transition cow immune function and interaction with metabolic diseases. Pages 45-57 in *Proc. 17<sup>th</sup> Annu. Tri-State Dairy Nutr. Conf.* The Ohio State University, Columbus.
- Gonzalez-Recio O, Pryce JE, Haile-Mariam M, Hayes BJ.** 2014. Incorporating heifer feed efficiency in the Australian selection index using genomic selection. *Journal of Dairy Science* **97**, 3883-3893.
- Grainger C, Auldish MJ, O'Brien G, Macmillan KL, Culley C.** 2009. Effect of type of diet and energy intake on milk production of Holstein-Friesian cows with extended lactations. *Journal of Dairy Science* **92**, 1472-1492.
- Gurmessa J, Melaku A.** 2012. Effect of Lactation Stage, Pregnancy, Parity and Age on Yield and Major Components of Raw Milk in Bred Cross Holstein Friesian Cows. *World Journal of Dairy & Food Science* **7(2)**, 146-149.
- Habtamu A, Solomon A, Yoseph M.** 2012. Influence of non-genetic factors on growth traits of Horro (Zebu) and their crosses with Holstein Friesian and Jersey cattle. *International Journal of Livestock Production* Vol. **3(7)**, pp. 72-77.
- Hazrat A, Nisar M, Sher K, Zaman S.** 2015. Role of economic plants in the community development of Dir valley, Khyber Pakhtunkhwa, Pakistan. *FUUAJST. Journal of Biology*. **5(1)**, 137-143
- Heins BJ, Hansen LB, Seykora AJ.** 2006a. Calving difficulty and stillbirths of pure Holsteins versus crossbreds of Holstein with Normande, Montbeliarde, and Scandinavian Red. *Journal of Dairy Science* **89**, 2805-2810.
- Hovinen M, Rasmussen DM, Pyrala S.** 2009. Udder health of cows changing from tie stalls to free stalls with conventional milking to free stalls either with conventional or automatic milking. *Journal of Dairy Science* **92**, 3696-3703.

- Hristov S.** 2014. The relationship between rearing system, animal needs index and dairy cows milk traits. *Mljekarstvo* **64(3)**, 186-194.
- Ihsanullah, Qureshi MS.** 2019. Effect of Stress on reproductive physiology of various graes of crossbred cattle under subtropical conditions of North-Western Pakistan. PhD Dissertation, FAHVS, The University of Agriculture, Peshawar. Pakistan.
- Islam SM, Kundu SK.** 2011. Effect of genotypes and parity on reproductive and productive attributes in dairy cattle of Natore Districts Bangladesh. *Pakistan Journal of Biological Sciences* **7**, 1269-1276.
- Jankowska M, Neja W, Krezel-Czopek S.** 2012. Effect of extended lactations on milk and reproductive performance of cows. *Acta Scientiarum Polonorum – Zootechnica* **11**, 15-22.
- Janu's E, Borkowska D, Grzesik K.** 2007. Relationship between body condition of cows in dryperiod and level of their energy reserves and productivity in the next lactation. *Rocz. Nauk. PTZ* **3(4)**, 149-156.
- Kamal MM, Van Eetvelde M, Depreester E, Hostens M, Vandaele L, Opsomer G.** 2014. Age at calving in heifers and level of milk production during gestation in cows are associated with the birth size of Holstein calves. *Journal of Dairy Science* **97(9)**, 5448-5458.
- Khan D,** 2004. Breed characteristics and relative performance of *Achai* cattle at Dir, (unpublished M.Sc thesis, Department of Agriculture Sciences, Allama Iqbal Open University Islamabad, Pakistan
- Khattak AHK, Islam MN, Khan MS, Iqbal M, Shah IA, Ghani S, Ali A, Hameed Z, Hassan MF, Ali T.** 2018. Comparative performance of Jersey sired calves from *Achai* dams and Azakheli buffalo calves fed with milk replacer. *Pakistan of Journal Zoology* **50(5)**, 1987-1990.
- Klopchich M, Hepon MC, Osterc J, Kompan D, Halchmi I, Løvendahl P.** 2007. A comparison between housing systems of dairy cows with regard to milk quality, animal welfare and animal health. EAAP, Book of abstracts, No. 13, Dublin, Ireland, 26-29 August 2007, 136.
- Kolver ES, Roche JR, Burke CR, Kay JK, Aspin PW.** 2007. Extending lactation in pastured based dairy cows; 1. Genotype and diet effect on milk and reproduction. *Journal of Dairy Science* **90**, 5518-5530.
- Kumar N, Chandrahas, Gaur GK, Sahoo SP, Devi LS, Tripathi AK.** 2017. Factors affecting growth and mortality pattern of Tharpakar calves at organized dairy farm. *International Journal of Livestock Research* **7(5)**, 87-92.
- Kurtu MY.** 2003. Certain aspects of the dairy system in the Harar milkshed, eastern Ethiopia. PhD thesis, University of the Free State, Bloemfontein, Faculty of Natural and Agricultural Sciences, Department of Animal, Wildlife and Grassland Sciences, South Africa. 195 pp.
- Lake SL, Scholljegerdes EJ, Atkinson RL, Navigihugu V, Paisley SI, Rule DC, Moss GE, Robinson TJ, Hess BW.** 2005. Body condition score at parturition and postpartum supplemental fat effects on cow and calf performance. *Journal of Animal Science* **83**, 2908-2917.
- Lakshmi S, Gupta BR, Sudhakar K, Prakash MG, Sharma S.** 2009. Genetic analysis of production performance of Holstein Friesian x Sahiwal cows. *Tamilnadu Journal of Veterinary and Animal Sciences* **5(4)**, 143-148.
- Lakshmi SB, Romesh B, Gupta R, Prakash MG, Sudhakar K, Sharma S.** 2010. Genetic analysis of production performance of Frieswal cattle. *Tamilnadu Journal of Veterinary and Animal Sciences* **6(5)**, 215-222.
- Lambertz C, Sanker C, Gaully M.** 2014. Climatic effects on milk production traits and somatic cell score in lactating Holstein-Friesian cows in different housing systems. *Journal of Dairy Science* **97(1)**, 319-329.
- Lammoglia MA, Willard ST, Oldham JR, Randel RD.** 1996. Effects of dietary fat and season on steroid hormonal profiles before parturition and on hormonal, cholesterol, triglycerides, follicular patterns, and post-partum reproduction in Brahman cows. *Journal of Animal Sciences* **74**, 2253-2262.

- Lehmann JO, Mogensen L, Kristensen T.** 2017. Early lactation production, health, and welfare characteristics of cows selected for extended lactation. *Journal of Dairy Science* **100(2)**, 1487-1501.
- Loker S, Bastin C, Miglior F, Sewalem A, Schaeffer LR, Jamrozik J, Ali A, Osborne V.** 2012. Genetic and environmental relationships between body conditionscore and milk production traits in Canadian Holsteins. *Journal of Dairy Science* **95(10)**, 410-419.
- Lundborg GK, Oltenacu PA, Maizon DO, Svensson EC, Liberg PGA.** 2003. Dam-related effects on heart girth at birth, morbidity and growth rate from birth to 90 days of age in Swedish dairy calves. *Previews of Veterinary Medicine* **60**, 175-190.
- Maekawa M, Beauchemin KA, Christensen DA.** 2002. Effect of concentrate level and feeding management on chewing activities, saliva production, and ruminal pH of lactating dairy cows. *Journal of Dairy Science* **85**, 1165-1175.
- Meikle A, Kulcsar M, Chilliard Y, Febel H, Delavaud C, Cavestany D, Chilbroste P.** 2004. Effects of parity and body condition at parturition on endocrine and reproductive parameters of the cow. *Reproduction* **127**, 727-737.
- Melaku M, Zeleke M, Getinet M, Mengistie T.** 2011a. Pre-weaning growth performances of Fogera calves at Metekel cattle improvement and multiplication ranch, North West Ethiopia. *Livestock Research for Rural Development* **23**, 78-86.
- Miglior F, Muir BL, Van Doormaal BJ.** 2005. Selection indices in Holstein cattle of various countries. *Journal of Dairy Science* **88**, 1255-1263.
- Mohamed AM.** 2004. Studies of some performance traits of Butana cattle in Atbara live-stock research station. MSc thesis University of Khartoum, Sudan.
- Msangi BSJ, Bryant MJ, Thorne PJ.** 2005. Some factors affecting variation in milk yield in crossbred dairy cows on smallholder farms in North-east Tanzania. *Tropical Animal Health and Production* **37**, 403-412.
- Mulliniks JT, Cox SH, Kemp ME, Endecott RL, Waterman RC, Vanleeuwen DM.** 2015. Relationship between body condition score at calving and reproductive performance in young postpartum cows grazing native range **1**, 2811-2817.
- Naceur M, Bouallegue M, Frouja S, Ressaissi Y.** 2012. Effects of Environmental Factors on Milk Yield, Lactation Length and Dry Period in Tunisian Holstein Cows. In N. Chaiyabutr (Ed.), *Milk production-An Up-to-Date Overview of Animal Nutrition, Management and Health* (1st ed., pp. 2-13). London: Intech Publisher. UK.
- Nickerson SC.** 1995. *Milk Quality: Factors affecting milk composition.* (F. Harding, Ed.) (1<sup>st</sup> Edition). Chapman & Hall, Wiltshire, UK.
- Pace JE, Wakeman DL.** 2003. Determining the age of cattle by their teeth. CIR253. Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences. University of Florida, Gainesville, FL.
- Phyn CVC, Clark C, Aspin PW, Kolver ES.** 2008. Effect of genotype and diet on feed conversion efficiency of dairy cows during a 600-day extended lactation. *Proceedings of the New Zealand Society of Animal Production* **68**, 100-104.
- Prendiville R, Pierce KM, Buckley F.** 2010. A comparison between Holstein-Friesian and Jersey dairy cows and their F1 cross with regard to milk yield, somatic cell score, mastitis, and milking characteristics under grazing conditions. *Journal of Dairy Science*. **93**, 2741-2750.
- Ratnayake D, Berglund B, Bertilsson J, Forsberg M, Gustafsson H.** 1998. Fertility in dairy cows managed for calving intervals of 12, 15 or 18 months. *Acta Veterinaria Scandinavica* **39(1)**, 215-228.
- Roche JR, Lee JM, Macdonald KA, Berry DP.** 2007. Relationships among body condition score, body weight, and milk production variables in pasture-based dairy cows. *Journal of Dairy Science* **90**, 3802-3815.

- Saleem M, Rahim I, Rueff H, Khan M, Maselli D, Wiesmann U, Muhammad S.** 2012. Effect of management on reproductive performances of the *Achai* cattle in the Hindu Kush (Northern Pakistan). *Tropical Animal Health and Production* **41(3)**, 137-142.
- Samarutel J, Ling K, Jaakson H.** 2006. Effect of body condition score at parturition on the production performance, fertility and culling in primiparous Estonian Holstein cows. *Veterinaija IR Zootechnikaal T.* **36(58)**, 69-74.
- Sattar A, Mirza RH, Niazi AAK, Latif M.** 2005. Productive and reproductive performance of Holstein Friesian cows in Pakistan. *Pakistan Veterinary Journal* **25(2)**, 75-82.
- Simensen E, Østerås O, Bøe KE, Kielland C, Ruud EL, Naess G.** 2010. Housing system and herd size interactions in Norwegian dairy herds; associations with performance and disease incidence. *Acta. Veterinary Scandinavia* **16(52)**, 14.
- Singh CV.** 2015. Cross-breeding in Cattle for Milk Production: Achievements, Challenges and Opportunities in India-A Review. *Advances in Dairy Research* **4(3)**, 1-14.
- Sölkner J, Miesenberger J, Willam A, Fuerst C, Baumung R.** 2000. Total merit indices in dual purpose cattle. *Arch. Tierz.* **43**, 597-608.
- Sørensen MK, Norberg E, Pedersen J, Christensen LG.** 2008. Invited review: Crossbreeding in dairy cattle: A Danish perspective. *J. Dairy Sci.* **91**, 4116-4128.
- Swali A, Wathes DC.** 2006. Influence of the dam and sire on size at birth and subsequent growth, milk production and fertility in dairy heifers. *Theriogenology* **66**, 1173-1184.
- Symonds ME, Sebert SP, Budge H.** 2010. Nutritional regulation of fetal growth and implications for productive life in ruminants. *Animal* **4**, 1075-1083.
- Tesfa A, Kumar D, Abegaz S, Mekuriaw G, Bimerew T, Kebede A, Bitew A, Ferede Y, Mazengia H, Tilahun M.** 2016. Growth and reproductive performance of Fogera cattle breed at Andassa Livestock Research Centre. *Livestock Research for Rural Development* **28(1)**, 56-62.
- Van Knegsel ATM, Rummelink GJ, Jorjong S, Fievez V, Kemp B.** 2014. Effect of dry period length and dietary energy source on energy balance, milk yield, and milk composition of dairy cows. *Journal of Dairy Science* **97(3)**, 1499-1512.
- Wakchaure R, Ganguly S, Para PA, Kumar P, Kumar A.** 2015. Development of Crossbred Cattle in India: A Review International. *Journal of Emerging Technology and Advance Engineering* **5**, 75.
- Weigel KA, Barlass KA.** 2003. Results of a producer survey regarding crossbreeding on US dairy farms. *Journal of Dairy Science* **86**, 4148-4154.
- Zhang WC, Nakao T, Kida K, Moriyoshi M, Nakada K.** 2002. Effect of nutrition during pregnancy on calf birth weights and viability and fetal membrane expulsion in dairy cattle. *J. Reprod. Dev.* **48**, 415-422.