



## Influence of different ratio of soil and organic fertilizers on growth and yield of strawberry and marigold for rooftop garden

Alak Barman<sup>1</sup>, Swarna Shome<sup>2\*</sup>

<sup>1</sup>Soil Science Division, Bangladesh Agricultural Research Institute, Gazipur, Bangladesh

<sup>2</sup>Department of Agronomy, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh

**Key words:** Vermicompost, Compost, Cowdung, Days to 1<sup>st</sup> flowering, Roof garden.

<http://dx.doi.org/10.12692/ijb/16.3.448-453>

Article published on March 29, 2020

### Abstract

Urban farming with organic fertilizer produces safe food and meet family nutrition. Again, flower production fulfil aesthetic purpose. Considering nutritional and aesthetic purpose, an experiment was conducted in Gazipur (rooftop of Soil Science Division) during the year of 2018-19 to scrutinize the effect of different ratio of soil and organic materials on the growth and yield of fruits and flowers for rooftop garden. The experiment was laid out in a completely randomized design with three replications and consisted of six different treatments *viz.* T<sub>1</sub> = 1 kg Compost for 1 kg soil, T<sub>2</sub> = 1 kg Compost for 2 kg soil, T<sub>3</sub> = 1 kg Cowdung for 1 kg soil, T<sub>4</sub> = 1 kg Cowdung for 2 kg soil, T<sub>5</sub> = 1 kg Vermicompost for 1 kg soil & T<sub>6</sub> = 1 kg Vermicompost for 2 kg soil. Pot was used for fruit and flower ((v Strawberry (BARI Strawberry -3) and Marigold (Hybreed Inca). Prior to setting the experiments initial soil samples as well as organic fertilizers were analyzed and nutrient status was determined. Pot was used for fruit and flower *viz.* Strawberry (BARI Strawberry -3) and Marigold (Hybrid Inca). Results revealed that marigold performed better in T<sub>5</sub> treatment (1 kg vermicompost for 1 kg soil) in case of all parameters while strawberry showed best performance for all parameters except days to 1<sup>st</sup> flowering compared to others in the experiments. The lowest yield was recorded from the T<sub>4</sub> treatment (1 kg cowdung for 2 kg soil).

\* **Corresponding Author:** Swarna Shome ✉ [swarnashome.sau@gmail.com](mailto:swarnashome.sau@gmail.com)

## Introduction

Urban agriculture is a global and growing pursuit for economic development, job creation, food security and community building. But major constraints for this are competition for space with other forms of urban development, a lack of formalized land use rights, and health hazards related to food contamination. The use of green roof technology in urban agriculture has the potential to alleviate some of these problems, without adversely affecting the benefits provided by urban agriculture. It would not only enable the use of land for development and agriculture, but may facilitate the formation of formal space and water use agreements and enable redistribution of ground level resources among urban farmers (Whittinghill and Rowe, 2012). This could decrease the use of contaminated land and water at ground level and alleviate health concerns.

Rooftop gardening is going to be popular in urban areas of Bangladesh. Bangladesh is one of the main victims of climate change. A country needs 25% of its land to be occupied by forests to maintain its ecological balance, but here the percentage is less than 8% (BBS, 2015). Due to the urbanization, our cultivable lands decrease day by day. As there is limited scope for horizontal expansion of agriculture, vertical expansion is one of the major ways to increase crop productivity. Rooftop gardening is one of the potential areas for vertical expansion. As it is estimated that, there are about 2 lac house buildings in Dhaka city. A rooftop garden not only can be a source of agricultural production but also can be able to fix CO<sub>2</sub> and some other gases causing greenhouse effect (BARI, 2019). So, there are many scopes for rooftop gardening.

Organic fertilizers are utilized globally to protect the soils against deterioration and food pollution. Organic nutrients increase soil enzyme activity, available nitrates, carbon to total organic carbon ratio and metabolic quotients resulting in enhanced soil fertility (Okwuagwu *et al.*, 2003). Organic fertilizers improve soil fertility by modifying soil structure, pH, biophysical conditions and availability of essential

nutrients (Atiyeh *et al.*, 2002). Though some interested people are producing vegetables, fruits and flowers on their rooftop but research information on nutrient management for a rooftop garden is not available. Moreover, extensive literature review showed very little or no information regarding fertilizer trials on rooftop garden. Therefore, there is a possibility to increase the yield of a rooftop garden through integrated nutrient management approach. Proper nutrient management can produce maximum yield of a rooftop garden. Considering the above facts the experiment was conducted to find out the optimum soil and manure ratio as a media for better growth and development of crops under rooftop garden, to increase yield and ensure family nutrition, to focus on the promotional and awareness aspects by providing incentives for environmental and aesthetic purpose.

## Material and methods

Experiment was conducted at the rooftop of Soil Science Division, BARI, Gazipur during 2018-2019. The initial soil sample, vermicompost, compost and cowdung will be collected before establishing the experiment and was analyzed in the laboratory following standard methods. Chemical properties of soil and nutrient status of vermicompost, compost and cowdung were illustrated in Table 1 & 2.

Organic based research (Only used organic fertilizers) for fruit and flower production in the rooftop garden was laid out in a completely randomized design with three replications. Six different treatments *viz.* T<sub>1</sub> = 1 kg Compost for 1 kg soil, T<sub>2</sub> = 1 kg Compost for 2 kg soil, T<sub>3</sub> = 1 kg Cowdung for 1 kg soil, T<sub>4</sub> = 1 kg Cowdung for 2 kg soil, T<sub>5</sub> = 1 kg Vermicompost for 1 kg soil & T<sub>6</sub> = 1 kg Vermicompost for 2 kg soil will be selected. Pot was used for fruit and flower *viz.* Strawberry (BARI Strawberry -3) and Marigold (Hybreed Inca).

1/2 organic fertilizer was applied at the time of final soil organic ratio preparation. The remaining half organic fertilizer was applied as top dress. Irrigations and other intercultural operation were done as when

required.

### Statistical analysis

All the collected data were analyzed following the analysis of variance (ANOVA) technique and using Statistix 10 package and the mean differences were adjudged by LSD technique (Gomez and Gomez, 1984).

## Results and discussion

### Effect of soil and organic fertilizer ratio on strawberry

The influence of different ratio of soil and organic matter on the growth and yield of Strawberry are presented in Table 3. The highest plant height (23.9 cm) was obtained from T<sub>5</sub> treatment (1 kg Vermicompost for 1 kg soil) whereas the lowest plant

height (19.3 cm) was obtained from T<sub>4</sub> treatment (1 kg cowdung for 2 kg soil). Significantly maximum number of runner plant<sup>-1</sup> (7.4) was obtained from T<sub>5</sub> treatment whereas the minimum number of runner plant<sup>-1</sup> (3.4) was obtained from T<sub>4</sub> treatment (1 kg cowdung for 2 kg soil). The longest period was required for flowering (74.8 days) in T<sub>4</sub> treatment while shortest period (65.4 days) in T<sub>5</sub> treatment. Significantly maximum number of fruits plant<sup>-1</sup> (24.4) was counted in T<sub>5</sub> treatment followed by T<sub>1</sub> treatment while the minimum number of fruits plant<sup>-1</sup> (12.1) was obtained from T<sub>4</sub> treatment (1 kg Cowdung for 2 kg soil). The maximum individual fruit weight (16.3 g) was obtained from T<sub>5</sub> treatment which was statistically similar with T<sub>1</sub> treatment followed by T<sub>3</sub> treatment whereas minimum individual fruit weight (11.9 g) was recorded from T<sub>4</sub> treatment.

**Table 1.** Chemical properties of soil (Initial) used in the rooftop experiment.

Soil properties	pH	OM	Total N	K	Ca	Mg	P	S	Zn	B	Cu	Fe	Mn
Analytical value	6.7	1.00	0.07	0.19	4.9	2.2	13.0	14	1.5	0.19	5.0	39	2.9
Critical level	-	-	0.12	0.12	2.0	0.5	7	10	0.6	0.2	0.2	4.0	1.0

**Table 2.** Nutrient status of vermicompost, compost and cowdung used in the rooftop experiment.

Name of the manure	pH	OC	Ca	Mg	K	Total N	P	S	Zn	B	Pb	Cd	As
Compost	7.1	16.3	1.50	2.10	1.17	1.15	0.79	0.50	0.14	0.013	2.89	2.11	1.72
Cowdung	7.5	15.4	2.23	0.44	0.69	1.02	0.57	0.36	0.15	0.011	3.10	2.84	1.26
Vermicompost	7.2	17.9	2.10	2.60	1.94	1.35	1.26	0.89	0.16	0.015	2.61	2.19	1.14

Moisture content of Compost = 12.15 %, Cowdung = 12.46 % and Vermicompost = 11.96 %.

The highest fruit yield plant<sup>-1</sup> (390.7 g) was recorded from T<sub>5</sub> treatment followed by T<sub>1</sub> and T<sub>3</sub> treatment whereas the lowest yield (143.9 g) was recorded in T<sub>4</sub> treatment. Kumar *et al.* (2015a) and Joshi and Vig (2010) also recorded significant influence of inclusion of vermicompost with soil on plant height, first flowering and flowers plant<sup>-1</sup>. Mehraj *et al.* (2014) reported significant influence on maximum fruits number and fruit weight. Bhattacharyya *et al.* (2003) stated the reason that vermicompost improved vegetative growth characters in strawberry by increasing soil enzyme activity and improving soil aeration. Tagliavini *et al.* (2005) found that the optimum level of nutrients as N, P and K and

hormones provided by vermicompost played a significant role in increasing gibberellic acid in roots thus breaking bud dormancy and increasing flowering buds and fruiting sites, total number of flowers was produced in strawberry. Arancon *et al.* (2004) reported that Vermicompost applications increased strawberry growth and yields significantly. According to Nourbakhsh (2007), vermicompost enhances soil aeration, builds up water retention capacity of soil because of its high organic matter content and promotes better root growth and nutrient absorption. Vermicompost is the builder of protein and is the main constituent of protoplasm in plants thus; the increase in nitrogen supply accelerates synthesis of

amino acids which might have indirectly exhibited increase in plant height of strawberry plant (Kumar *et al.*, 2015b). Singh *et al.* (2008) observed significant increase in fruit yield and flowering of strawberry with vermicompost based fertilizer and concluded that increase in plant growth might be due to the improvement in physio-chemical properties of soil; increase in enzymatic activity, microbial population

and also increase in plant growth hormones by application of vermicompost. Singh *et al.* (2011) opined that increase in number of runners per plant might be due to increased growth of plant in the form of height and number of leaves, which accumulated more photosynthates and thereby increased runners and leaf area per plant.

**Table 3.** Influence of different ratio of soil and organic matters on the growth and yield of Strawberry.

Treatment	Plant height (cm)	Number of runner plant <sup>-1</sup>	Days to 1 <sup>st</sup> flowering	No. of fruits plant <sup>-1</sup>	Individual fruit wt. (g)	Fruit yield plant <sup>-1</sup> (g)
T <sub>1</sub>	22.9ab	6.5b	67.2c	20.2b	15.3ab	318.7b
T <sub>2</sub>	20.6cd	4.8d	72.5ab	14.4cd	12.9cd	183.4e
T <sub>3</sub>	22.5ab	6.3bc	68.3bc	17.5bc	14.2bc	266.5c
T <sub>4</sub>	19.3d	3.4e	74.8a	12.1d	11.9d	143.9f
T <sub>5</sub>	23.9a	7.4a	65.4c	24.4a	16.3a	390.7a
T <sub>6</sub>	21.8bc	5.5cd	69.2bc	16.3c	14.1bc	229.8d
CV (%)	4.58	8.85	4.45	11.44	7.13	7.92

Means followed by the same letter in a column are not statistically significant at 5% level.

T<sub>1</sub> = 1 kg compost for 1 kg soil, T<sub>2</sub> = 1 kg compost for 2 kg soil, T<sub>3</sub> = 1 kg cowdung for 1 kg soil, T<sub>4</sub> = 1 kg cowdung for 2 kg soil, T<sub>5</sub> = 1 kg vermicompost for 1 kg soil & T<sub>6</sub> = 1 kg vermicompost for 2 kg soil.

These results gain support from the findings of Tripathi *et al.* (2015) noted earliest flowering and maximum berry weight with vermicompost. This increase in fruit size and weight during the present investigation might be due to the increased photosynthetic ability of plants fertilized with vermicompost, which in turn might have favored an

increased accumulation of dry matter (Beer *et al.*, 2017).

#### *Effect of soil and organic fertilizer ratio on marigold*

The influence of different ratio of soil and organic matter on the growth and yield of Marigold is presented in Table 4.

**Table 4.** Influence of different ratio of soil and organic matters on the growth and yield of Marigold.

Treatment	Plant height (cm)	Number of branches plant <sup>-1</sup>	Number of flowers plant <sup>-1</sup>	Flower diameter (cm)	Flower wt. plant <sup>-1</sup> (kg)
T <sub>1</sub>	32.2a	10.5ab	29.2ab	8.5a	0.53b
T <sub>2</sub>	30a	9.7abc	24.1b	8.3a	0.41d
T <sub>3</sub>	31.9a	10.8ab	27.3ab	8.5a	0.49c
T <sub>4</sub>	23b	8.6c	9.8c	6.7b	0.21e
T <sub>5</sub>	32.3a	11.3a	31.6a	8.6a	0.58a
T <sub>6</sub>	31.9a	10.8ab	27.3ab	8.4a	0.49c
CV (%)	6.98	10.46	13.27	3.80	4.85

Means followed by the same letter in a column are not statistically significant at 5% level

T<sub>1</sub> = 1 kg compost for 1 kg soil, T<sub>2</sub> = 1 kg compost for 2 kg soil, T<sub>3</sub> = 1 kg cowdung for 1 kg soil, T<sub>4</sub> = 1 kg cowdung for 2 kg soil, T<sub>5</sub> = 1 kg vermicompost for 1 kg soil & T<sub>6</sub> = 1 kg vermicompost for 2 kg soil.

The highest plant height (32.3 cm) was recorded in T<sub>5</sub> treatment whereas the lowest plant height (23 cm) was observed in T<sub>4</sub> treatment. The highest number of branches plant<sup>-1</sup> (11.3) was recorded in T<sub>5</sub> treatment

whereas the lowest number of branches plant<sup>-1</sup> (8.6) was observed in T<sub>8</sub> treatment. The maximum flower diameter (8.6 cm) was observed in T<sub>5</sub> treatment whereas the minimum flower diameter (6.7 cm) was

observed in T<sub>4</sub> treatment. The maximum flower wt. plant<sup>-1</sup> (0.58 kg) was observed in T<sub>5</sub> treatment followed by whereas the minimum flower wt. plant<sup>-1</sup> (0.21 kg) was observed in T<sub>4</sub> treatment. Arancon *et al.* (2008) showed that application of vermicompost increased yield and improved nitrogen and phosphorus uptake. Vermicompost has been reported to have effects similar to plant growth regulating substances and hormones (Muscolo, 1999). Sardohei *et al.* (2014) documented that vermicompost had significant positive effects on flower numbers, leaf growth and shoot fresh and dry weights. Gupta *et al.* (2014) noted enhanced effect of vermicompost on growth and productivity of marigold plants.

### Conclusion

Results of the experiment revealed that strawberry and marigold performed better in case of growth and yield with 1 kg vermicompost for 1 kg soil. So, 1 kg vermicompost for 1 kg soil may be a good practice for strawberry and marigold production on rooftop garden which meets nutrition and aesthetic purpose.

### Acknowledgment

Authors are immensely grateful to Ministry of Science and Technology, Bangladesh for supporting this research work. Authors would like to express their sincere gratitude to concern personnel of the Soil Science Division, Bangladesh Agricultural Research Institute, Gazipur for valuable guide and support.

### References

**Arancon NQ, Edwards CA, Babenko A, Cannon J, Galvis P, Metzger JD.** 2008. Influences of vermicomposts, produced by earthworms and microorganisms from cattle manure, food waste and paper waste, on the germination, growth and flowering of petunias in the greenhouse. *Applied Soil Ecology* **39**(1), 91-99. <http://dx.doi.org/10.1016/j.apsoil.2007.11.010>

**Arancon NQ, Edwards CA, Bierman P, Welch C, Metzger JD.** 2004. The influence of vermicompost applications to strawberries: Part 1. Effects on growth and yield. *Bioresource Technology*

**93**, 145-153.

<http://dx.doi.org/10.1016/j.biortech.2003.10.014>.

**Atiyeh RM, Edwards CA, Metzger JD, Lee S, Arancon NQ.** 2002. The influence of humic acids derived from earthworm-processed organic wastes on plant growth. *Bioresource Technology* **84**, 7-14. [https://doi.org/10.1016/S0960-8524\(02\)00017-2](https://doi.org/10.1016/S0960-8524(02)00017-2).

**BARI (Bangladesh Agricultural Research Institute).** 2019. BARI Annual Report. Gazipur, Dhaka- 1701 p 232.

**Beer K, Santosh K, Gupta AK, Syamal MM.** 2017. Effect of Organic, Inorganic and Bio-Fertilizer on Growth, Flowering, Yield and Quality of Strawberry (*Fragaria × Ananassa Duch.*) cv. Chandler. *International Journal of Current Microbiology and Applied Science* **6**(5), 2932-2939. <https://doi.org/10.20546/ijcmas.2017.605.332>

**BBS (Bangladesh Bureau of Statistics).** 2015. Statistical Yearbook of Bangladesh, Statistics Division, Ministry of Planning, Govt. of the People's Republic of Bangladesh, p 90.

**Bhattacharyya P, Chakrabarti K Chakraborty A.** 2003. Effect of MSW compost on microbiological and biochemical soil quality indicators. *Compost Science and Utilization* **11**(3), 220-227. <https://doi.org/10.1080/1065657X.2003.10702130>.

**Gomez KA, Gomez AA.** 1984. Statistical procedure for agricultural research. Second Edn. Intl. Rice Res. Inst., John Wiley and Sons. New York, p 1-340.

**Gupta R, Yadav A, Garg VK.** 2014. Influence of vermicompost application in potting media on growth and flowering of marigold crop. *International Journal of Recycling of Organic Waste in Agriculture* **3**, 47. <https://doi.org/10.1007/s40093-014-0047-1>.

**Joshi R, Vig PA.** 2010. Influence of vermicompost on growth, yield and quality of tomato (*Lycopersicon esculentum* L). *African Journal of Basic and Applied*

Sciences **2**, 117-123.

**Kumar N, Singh HK, Mishra PK.** 2015a. Impact of Organic Manures and Biofertilizers on Growth and Quality Parameters of Strawberry cv. Chandler. Indian Journal of Science and Technology **8(15)**, 1-6. <http://dx.doi.org/10.17485/ijst/2015/v8i15/51107>.

**Kumar N, Ram RB, Mishra PK.** 2015b. Effect of vermicompost and azotobacter on quality parameters of strawberry (*Fragaria x ananassa* Duch.) cv. sweet Charlie. International Journal of Agricultural Science and Research **5(4)**, 269-276.

**Mehraj H, Ahsan MK, Hussain MS, Rahman MM, Jamal UFM.** 2014. Response of different organic matters in strawberry. Bangladesh Research Publication Journal **10(2)**, 151-161. <http://www.1410019/1410019>.

**Muscolo AF, Bovalo F, Gionfriddo F, Nardi S.** 1999. Earthworm humic matter produces auxin-like effects on *Daucus carota* cell growth and nitrate metabolism. Soil Biology and Biochemistry **31**, 1303-1311. [https://doi.org/10.1016/S0038-0717\(99\)00049-8](https://doi.org/10.1016/S0038-0717(99)00049-8)

**Nourbakhsh F.** 2007. Influence of vermicomposting on solid wastes decomposition kinetics in soils. Journal of Zhejiang University Science **8(10)**, 725-730. <http://dx.doi.org/10.1631/jzus.2007.B0725>.

**Okwuagwu MI, Alleh ME, Osemwota IO.** 2003. The effects of organic and inorganic manure on soil properties and yield of okra in Nigeria. African Crop Science Conference Proceeding **6**, 390-393.

**Sardoei AS, Roien A, Sadeghi T, Shahadadi F, Mokhtari TS.** 2014. Effect of Vermicompost on the Growth and Flowering of African Marigold (*Tagetes*

*erecta*). American-Eurasian Journal of Agricultural and Environmental Science **14(7)**, 631-635. <http://dx.doi.org/10.5829/idosi.ajeaes.2014.14.07.12366>.

**Singh BK, Pathak KA, Verma AK, Verma VK, Deka BC.** 2011. Effects of vermicompost, fertilizer and mulch on plant growth, nodulation and pod yield of French bean (*Phaseolus vulgaris* L). Vegetable Crops Research Bulletin **74**, 153-165. <https://doi.org/10.2478/v10032-011-0013-7>

**Singh R, Sharma RR, Kumar S, Gupta RK, Patil RT.** 2008. Vermicompost substitution influences the physiological disorders, fruit yield and quality of strawberry (*Fragaria ananassa* Duch). Bioresource Technology **99**, 8507-8511. <https://doi.org/10.1016/j.biortech.2008.03.034>

**Tagliavini ME, Baldi E, Lucchi P, Antonelli M, Sorrenti G, Baruzzi G, Faedi G.** 2005. Dynamics of nutrient uptake by strawberry plants (*Fragaria x Ananassa* Duch.) grown in soil and soilless culture. European Journal of Agronomy **23**, 15-25. <http://dx.doi.org/10.1016/j.eja.2004.09.002>

**Tripathi VK, Kumar S, Gupta AK.** 2015. Influence of Azotobacter and Vermicompost on Growth, Yield and Quality of Strawberry Cv. Chandler. Indian Journal of Horticulture **72(2)**, 201-205. <http://dx.doi.org/10.5958/0974-0112.2015.00039.0>

**Whittinghill LJ, Rowe DB.** 2012. The role of green roof technology in urban agriculture. Renewable Agriculture and Food Systems **27(4)**, 314-322. <http://dx.doi.org/10.1017/S174217051100038X>