



RESEARCH PAPER

OPEN ACCESS

In vitro antidiabetic, antiobesity and antioxidant activities of selected endemic plants from Mount Mayon and Mount Malinao Albay, Philippines

Lilibeth A Cajuday^{*1}, Daile Meek S Membreve¹, Michael V Montealegre²,
Jocelyn E Serrano¹, Diomerl Edward Baldo³

¹Department of Biology, Faculty of College of Science Bicol University, Legazpi City, Philippines

²Department of Chemistry, Faculty of College of Science Bicol University, Legazpi City, Philippines

³Department of Biology, Graduate Student of Bicol University Graduate School, Legazpi City Philippines

Key words: Phytochemical components, Antioxidant activity, Alpha-glucosidase, Lipase inhibition

<http://dx.doi.org/10.12692/ijb/16.3.648-661>

Article published on March 30, 2020

Abstract

The present study investigated the *in vitro* biological activities of the ethanol extracts of thirteen endemic medicinal plants in Albay against diabetes, obesity and as free radical scavenger. Alpha-glucosidase (AG), porcine pancreatic lipase (PPL) inhibition tests and DPPH radical scavenging assay were used to determine the antidiabetic, antiobesity and antioxidant properties of the extracts. Quantitative phytochemical analysis of the extracts was performed using standard methods. Of the 13 endemics, 6 plants exhibited potent inhibitory activity (>70%) against AG enzyme. *Hydnocarpus alcalae*, *Merremia peltata*, *Trema orientalis*, *Cascabela thevetia*, *Stachytarpheta jamaicensis*, and *Ficus septica*. *S. jamaicensis* has the lowest computed IC₅₀ (0.8ug/ml) indicating highest AG inhibitory activity compared to the other plants and the drug *Acarbose* (1.88ug/ml). A total of 7 plants were found to have strong inhibitory activity of >70% against PPL: *Leea guineensis*, *Solanum torvum*, *Cheilocostus speciosus*, *Melastoma malabathricum*, *Dendrocnide meyeniana*, *Cinnamomum mercadoi*, and *Gmelina arborea*. *L. guineensis* which recorded the highest activity against PPL, contains very high amount of alkaloids (237.49mg/g AE), flavonoids (490.54mg/g QE), and phenols (220.59mg/g GAE) and also exhibited strong antioxidant activity in the DPPH assay. *M. malabathricum* has the lowest IC₅₀ suggesting that it has the highest activity against PPL compared to the other plants and the drug *Orlistat* (1.81ug/ml). DPPH radical scavenging assay showed that all plant extracts possessed statistically (p≤0.05) higher antioxidant activity against ascorbic acid suggesting promising potent source of antioxidants. There is a need to conduct further study on the toxicity of the evaluated plants.

*Corresponding Author: Lilibeth A. Cajuday ✉ lacajuday@bicol-u.edu.ph

Introduction

Non-communicable diseases (NCDs), where diabetes mellitus and obesity are leading causes of mortality have now become the main threat to global health (Perk 2017). In the country, six million Filipinos have been diagnosed to have diabetes and this fig. could double to 12 million by 2040 because of undiagnosed diabetes cases as reported by the Philippine Center for Diabetes Education Foundation (2016). Concurrently, 18 million Filipinos are obese and overweight, according to a report released in 2016 by Asia Roundtable on Food Innovation for Improved Nutrition. Statistics likewise shows that 22.3% of Filipino adults are overweight and 6.1% are obese (FNRI, 2011). In the region particularly, there is high incidence of lifestyle related illnesses and metabolic disorders which is further aggravated by expensive medicines and inadequate health services (RDP, 2011). It is therefore necessary to address the health situation of the Albayanos and seek affordable alternative therapies from available natural local sources.

Investigation of natural products is a research field with great potential and is especially important in developing countries possessing great biodiversity, like the Philippines, having 2/3 of the earth's biodiversity and about 70-80% of the world's plant and animal species (CBD, 2009). To date, despite increasing research efforts leading to the discovery of many bioactive compounds from terrestrial plants including flavonoids, terpenoids, carotenoids, phyosterols, isothiocyanates and other phytochemicals (Lynn *et al.*, 2006; Hayes and Eggleston, 2008; Ragasa *et al.*, 2009; Raga *et al.*, 2011; Macabeo *et al.*, 2013), it is understandable that there is still a huge number of plant compounds that are not well investigated pharmacologically in the approximately 310,000 plant species described so far (IUCN, 2015). In the province of Albay, Mount Malinao and Mount Mayon are considered as key biodiversity areas that support endemics, abundant, indigenous and medicinal species. Phytochemical and pharmacological studies of species that are abundant in the locality and with reported medicinal uses are therefore promising areas for exploration.

This research is one of the first assessments of the *in vitro* biological activities of these endemic medicinal plants: (1) *Gmelina arborea* (Lamiaceae); (2) *Solanum torvum* (Solanaceae); (3) *Stachytarpheta jamaicensis* (Verbenaceae); (4) *Cascabela thevetia* (Apocynaceae); (5) *Melastoma malabathricum* (Melastomataceae); (6) *Dendrocide meyeniana* (Urticaceae); (7) *Cinnamomum mercadoi* (Lauraceae); (8) *Cheilocostus speciosus* (Costaceae); (9) *Trema orientalis* (Ulmaceae); (10) *Leea guineensis* (Vitaceae); (11) *Ficus septica* (Moraceae); (12) *Hydnocarpus alcalae* (Achariaceae); and (13) *Merremia peltata* (Convolvulaceae).

In this study, the 13 endemic medicinal plants were evaluated for their phytochemical components, antioxidant property, alpha-glucosidase and lipase inhibition activities using a simple, fast, efficient, and reliable spectrophotometric method. The plant extracts were also compared with standard chemical agents (e.g. Ascorbic acid, Acarbose and Orlistat) in order to assess their potential use as an alternative to these drugs.

Materials and methods

The study conducted comparative *in vitro* screening of the antioxidant, anti-diabetic and anti-obesity activity of thirteen (13) endemic medicinal plants from Mount Mayon and Mount Malinao Albay. The thirteen plants were selected because of their medicinal use, availability and endemism.

Collection of plant samples and preparation of extracts

Leaves of the thirteen endemic medicinal plants were obtained from Mt. Mayon and Mt. Malinao in Albay. The plants were identified and authenticated at Jose Vera Santos Memorial Herbarium Institute of Biology, University of the Philippines, Diliman Quezon City. Preparation of extracts and all analyses were done at College of Science Bicol University, Legazpi City. Prior to extraction, the different leaf samples were washed, air-dried and reduced to fine particles using Osterizer™ blender. The pulverized samples, soaked in 95% ethanol (1:2 w/v) for 72 hours were filtered and concentrated by evaporation under reduced pressure at 45°C using rotary vacuum evaporator.

The total ethanol extract concentrate yield per gram of dried plant material was determined using the formula: weight (g) of dried extract/dry-weight (g of plant material) x 100

Antioxidant Activity

DPPH assay

To evaluate free radical scavenging activity, sample stock solutions (1mg/ml) were serially diluted to prepare concentrations of 1000µg/ml, 500µg/ml, 250µg/ml, 125µg/ml, 62.5µg/ml and 31.5µg/ml of the extracts. About 50µL various concentrations of plant ethanolic extracts were mixed with 150 µL 0.1mM DPPH-ethanol solution. Ascorbic acid was used as a positive control and blank control was prepared with ethanol and DPPH. After 30 minutes of incubation at room temperature in the dark, the absorbance at 517nm was measured using BIOBASE-EL10B ELISA reader. The DPPH radical scavenging activity of the samples was calculated using the following formula described by Sroka and Cisowski (2005):

DPPH scavenging activity (%) = (Absorbance of sample)/(Absorbance of control) X 100.

Concentrations of 250, 125, 62.5, and 31.25µg/ml were used to determine the effective concentration (EC₅₀) to scavenge the DPPH radical by 50%. EC₅₀ were calculated using linear regression analysis described by Mensor *et al.* (2001) where plots were generated with the abscissa represented the concentration of tested plant extracts and the ordinate the scavenging activity from three independent experiments.

Anti-diabetic Activity

α-Glucosidase inhibition assay

Crude extracts from 13 identified plants were investigated for its α-glucosidase inhibitory activity using a Glucosidase Activity Assay Kit (Sigma Aldrich Co Ltd St. Louis, MO, USA). Samples were dissolved in dimethyl sulfoxide (DMSO) at various concentrations. 20 µl of each sample extracts were transferred at separate wells. The samples were treated with 200 µl Assay buffer, pH 7.0 and 8 µl (4-

Nitrophenyl-b-D- glucuronide) a-NPG Substrate. The initial absorbance of the released p-nitrophenol was measured at 405nm using ELISA microplate reader (Biobase EL10B). Samples were incubated at 37 °C and final absorbance was measured at 405nm after 20 minutes. The inhibition of α-glucosidase activity in the sample was calculated based from the formula described by Hyun *et al.*, (2015)

Inhibition rate (%) = $1 - \frac{\text{Abs sample} - \text{Abs blank}}{\text{Abs control}} \times 100$

Where Abs sample represents the absorbance of the extracts after 20 minutes, Abs blank denotes the absorbance of water added with calibrator, and Abs control represents the absorbance of the water (control).

Anti-obesity Activity

Lipase inhibition assay

The inhibition of lipase by the ethanolic extracts of the selected plant species was determined using a Lipase Activity Assay Kit (Sigma Aldrich Co Ltd St. Louis, MO, USA). Lipase activity was measured using a coupled enzyme reaction, which results in a colorimetric (570nm) product proportional to the enzymatic activity present. One unit of lipase is the amount of enzyme that will generate 1.0mmole of glycerol from triglycerides per minute at 37°C. Plant samples were diluted in DMSO and centrifuged at 10,000 x g for 10 minutes to remove insoluble materials. The inhibitory activities of the plant extracts and Orlistat were measured at concentrations of 1000, 500, 250, 125, 62.5 and 31.25µg/ml. Samples and standard solutions were adjusted to final volume of 50ul with Lipase Assay Buffer into a 96 well plate. 100ul of the reaction mix was added to each well by pipetting. After the mixture was incubated at 37°C for 2-3 minutes, the initial measurement was read at 570nm using ELISA EL10B. The mixture was further incubated until it reached the final measurement. The final absorbance measurement for calculating the enzyme activity was the penultimate reading at 30 minutes. The plate was protected from light during the incubation. The measurements were performed in triplicate. The inhibitory activity (%I) was calculated according to the following formula (Dechakhamphu and Wongchum, 2015), where A is the activity of the

enzyme initial absorbance, and a is the o blank control initial absorbance; B is the activity of the enzyme final absorbance, and b is the o blank control final absorbance:

$$1\% = \left(1 - \frac{B-b}{A-a}\right) \times 100$$

The half maximal inhibitory concentration (IC₅₀) determination

The IC₅₀ value of the extracts was determined at a concentration of 1000, 500, 250, 125, 62.5, and 31.25 µg/mL. *Acarbose* and *Orlistat* were used as a positive control. IC₅₀ value was calculated by the following formula (Dechakhamphu and Wongchum, 2015), where Low Inh%/HighInh% signify% inhibition directly below/above 50% inhibition, and Low Conc/High Conc are the corresponding concentrations of extract.

$$IC_{50} = \left(\frac{50\% - \text{low Inh}\%}{\text{high Inh}\% - \text{low Inh}\%}\right) \times (\text{high conc} - \text{low conc}) + \text{low conc}$$

Phytochemical Screening

Determination of Total Alkaloids

Spectrophotometric method was used to determine the total alkaloids using a Perkin Elmer Lambda 35 Double-beam UV/Vis Spectrometer. A 1000 ppm atropine stock solution was prepared to determine the standard calibration curve. Samples were prepared by dissolving 0.1 g of dried extract with 10mL 200-proof ethanol solvent. Sample, standard (20-150 µg/mL) and blank (ethanol) solutions were prepared by mixing 1mL aliquot with 5mL phosphate buffer (pH 4.7) and 5mL 0.1mM Bromocresol Green solution. The mixture was transferred to a funnel and three successive portions of 5, 3, then 2mL chloroform were used for solvent extraction. The chloroform portions were combined to make up a 10mL solution. Absorbance of each solution in quartz sample cell was read against the blank at 470nm. Concentrated plant extract was prepared for sample with absorbance below the linear range of the calibration curve. The total alkaloids are reported with mean values of three replicates in milligrams atropine equivalent (AE) per gram of dried extract.

Determination of Total Flavonoids

The total flavonoid contents of the extracts were measured by the aluminum chloride colorimetric assay as adopted from Sulaiman and Balachandran (2012). Briefly, a 1000 ppm quercetin stock solution was prepared for the standard calibration curve. Samples were then prepared by dissolving 0.1 g of dried extract with 25mL 200 proof ethanol solvent. Sample, standard (10-80 µg/mL) and blank (ethanol) solutions were added with 4mL of distilled water and 300 µL 5% NaNO₂. After five minutes, 300 µL 10% AlCl₃ was added to the mixture. For another 5 minutes, 2mL 1M NaOH was added to the mixture and diluted to 10mL with distilled water. Absorbance was read at 550nm using Biobase EL10B Microplate Reader. The total flavonoids are reported as milligrams quercetin equivalent (QE) per gram of dried extract.

Determination of Total Phenols

The total phenolic composition of the ethanolic extracts was determined using the Folin-Ciocalteu reagent as first described by Singleton *et al.* (1999). Briefly, a 1000ppm gallic acid stock solution was prepared to determine the standard calibration curve. Samples were then prepared by dissolving 0.1 g of dried extract with 25mL 200-proof ethanol. Sample, standard (10-50 µg/mL) and blank (ethanol) solutions were added with 0.4mL Folin-Ciocalteu reagent. After 5 minutes, 4mL 7% Na₂CO₃ was added to the mixture. The mixture was incubated for 90 minutes before diluting to 10mL with distilled water. Sample solutions were further filtered before reading the absorbance at 550nm using Biobase EL10B Microplate Reader. Solutions not within the linear range of the curve were either diluted or a concentrated. Total phenols are reported as milligrams gallic acid equivalent (GAE) per gram of dried extract.

Statistical Analysis

Statistical analysis of the data was performed using the SPSS 21.0 program. Data are expressed as mean ± SEM. Statistical comparisons were assessed by the Mann–Whitney U test. Values of $p < 0.05$ were considered significant.

Results and discussion

Phytochemical screening

Preliminary phytochemical analysis of the ethanolic extracts of the 13 endemic medicinal plants was determined by qualitative analyses. Phytoconstituents namely alkaloids, flavonoids and phenols were detected in the extracts. Results of the present study reflected that the quantity of alkaloids, flavonoids and phenols vary from one plant to another, and this finding is in agreement with previous studies (Srivastava *et al.*, 2012 and Liu 2008).

The data obtained after analysis of total alkaloids as shown in Table 1 was largely variable among the plants. The highest alkaloid content was found in *Stachytarpheta jamaicensis* (344.50mg/g) followed by *Trema orientalis*, *Hydnocarpus alcalae*, *Leea guineensis*, *Solanum torvum*, *Cinnamomum mercadoi* and *Gmelina arborea* with the values being 316.80, 271.06, 237.49, 154.23, 109.19 and 97.89mg/g, respectively.

Comparison of the flavonoid content of the 13 plants revealed that the leaves of *L. guineensis* were found to have maximum flavonoid content followed by leaves of *T. orientalis*, *M. peltata*, *H. alcalae*, *M. malabathricum*, *C. mercadoi*, *C. thevetia*, *G. arborea*, *D. meyeniana*, *C. speciosus*, *S. torvum*, *S. jamaicensis* and *F. septica*, with the minimum flavonoid content. Measurement of the total phenolic composition of the extracts suggested that the maximum phenolics (ug/ml) was found in the leaves of *C. thevetia* (237.80) followed by *L. guineensis* (220.59), *S. jamaicensis* (187.13), *M. peltata* (141.41), *D. meyeniana* (104.68), *C. speciosus* (94.53), *S. torvum* (89.48), *G. arborea* (86.16), *C. mercadoi* (70.43), *M. malabathricum* (63.90), *H. alcalae* (57.12). Relatively low level of phenols was detected in the leaves of *T. orientalis* and *F. septica*.

Table 1. Extraction yield, total alkaloids, flavonoids and phenols content of leaf extracts of endemic medicinal plants in Albay.

Scientific Name	Family	Extraction yield (EtOH)	Total Alkaloids	Total Flavonoids	Total Phenols
		% Yield (W/W)	AE (mg/g)	QE (mg/g)	GAE (mg/g)
<i>Gmelina arborea</i>	Lamiaceae	4.32	97.89±5.61	202.58±1.36	86.16±4.49
<i>Solanum torvum</i>	Solanaceae	9.9	154.23±5.0	127.02±1.82	89.48±6.51
<i>Stachytarpheta jamaicensis</i>	Verbenaceae	1.16	344.50±44.54	104.49±0.79	187.13±7.83
<i>Cascabela thevetia</i>	Apocynaceae	5.23	53.86±6.46	203.69±3.59	237.80±6.06
<i>Melastoma malabathricum</i>	Melastomataceae	2.49	31.12±12.67	267.87±2.88	63.90±1.58
<i>Dendrocnide meyeniana</i>	Urticaceae	7.85	46.80±3.98	201.22±1.71	104.68±7.73
<i>Cinnamomum mercadoi</i>	Lauraceae	7.11	109.19±3.00	221.28±1.33	70.43±1.05
<i>Cheilocostus speciosus</i>	Costaceae	3.1	18.63±2.14	156.74±3.12	94.53±2.52
<i>Trema orientalis</i>	Ulmaceae	10.67	316.80±41.19	391.39±5.53	35.32±0.62
<i>Leea guineensis</i>	Vitaceae	4.79	237.49±21.17	490.54±1.52	220.59±1.87
<i>Ficus septica</i>	Moraceae	2.99	62.78±6.57	77.11±4.11	5.01±0.02
<i>Hydnocarpus alcalae</i>	Achariaceae	1.73	271.06±2.68	329.90±1.14	57.12±1.87
<i>Merremia peltata</i>	Convolvulaceae	10.02	62.36±13.33	338.24±1.44	141.41±0.73

AE – Atropine equivalent, QE – Quercetin Equivalent, GAE – Gallic Acid Equivalent All result are reported as mean ± SEM in milligrams per gram of dried extract, N = 3

Antioxidant activity

The ethanol extracts of 13 endemic medicinal plants showed statistically higher antioxidant activity towards the DPPH radical compared with the standard ascorbic acid at higher concentrations (1000ug/ml-2500ug/ml) and comparable to ascorbic acid at lower doses (Table 2). At high dose (10000ug/ml) higher scavenging activity was observed

in *M. peltata* followed by *S. jamaicensis* and *G. arborea*. For EC₅₀, lowest EC₅₀ was observed in *M. peltata* followed by *C. speciosus*, *S. jamaicensis*, *C. mercadoi*, *G. arborea*, *D. meyeniana*, *M. malabathricum*, *T. orientalis*, *C. peruviana*, *S. torvum*, *F. septica*, *L. guineensis*, and *H. alcalae* with the highest EC₅₀. An EC₅₀ value is the concentration that is required to scavenge 50% of the free radicals in

the system that is inversely proportional to the antioxidant activity (Sahu, Kar and Routay, 2013). Thus, *M. peltata* with the lowest EC₅₀ is shown to possess the highest activity compared to other plants. To the authors' knowledge, this is the first report of

the antioxidant activity of *M. peltata* collected in Bicol region or in the Philippines. Antioxidants compounds such as alkaloids, flavonoids and saponins were observed in *M. peltata* collected in Iligan, Philippines (Perez *et al.*, 2015).

Table 2. DPPH scavenging activity of 13 Endemic Medicinal plants of Albay at different concentration.

Plant extracts	1000ug/ml	500ug/ml	25ug/ml	125ug/ml	62.5ug/ml	31.25ug/ml	*EC ₅₀ ±SEM (ug/ml)
<i>G. arborea</i>	293.00±29.00*	227.52±37.68*	181.12±32.66*	98.83±16.42*	57.51±8.23*	34.28±4.93	95.07±60.04
<i>S. torvum</i>	194.72±55.01*	174.1±55.87*	123.53±40.30*	72.45±19.16	43.13±7.96	22.79±1.51	243.03±129.71
<i>S. jamaicensis</i>	299.25±12.02*	212.08±23.13*	123.10±7.99*	74.89±4.26*	45.49±3.00*	31.35±2.21*	74.77±11.09
<i>C. thevetia</i>	185.38±30.12*	128.88±25.36*	73.29±10.19*	42.79±4.00*	28.74±2.39	23.35±1.33	193.46±66.41
<i>M. malabathricum</i>	241.06±19.16*	133.02±13.31*	75.89±5.15*	50.42±2.45*	38.70±1.52*	33.32±1.34	127.71±23.09
<i>D. meyeniana</i>	164.91±42.24*	140.42±22.29*	64.21±14.02*	37.92±6.20	36.53±1.17	39.06±4.62	113.38±70.68
<i>C. mercadoi</i>	162.38±39.55	129.21±17.04*	62.97±11.25*	43.16±5.38*	37.18±2.93	41.73±3.75	76.92±33.12
<i>C. speciosus</i>	131.85±37.81*	161.57±33.38*	49.89±8.16*	35.23±2.00	33.22±4.74	41.10±10.65	73.18±74.27
<i>T. orientalis</i>	179.53±31.17*	203.21±28.69*	83.01±5.98*	53.59±2.01	62.26±9.45*	48.94±10.09	157.17±61.75
<i>L. guineensis</i>	213.93±32.83*	131.28±31.59*	102.79±9.89*	134.32±28.80	99.93±25.54*	74.08±17.29	274.51±168.01
<i>F. septica</i>	180.84±31.18*	95.14±15.50*	54.91±5.73*	35.75±3.01	26.85±2.00	17.49±2.54*	255.40±79.65
<i>H. alcalae</i>	179.47±36.61*	125.41±40.82*	95.32±30.52*	52.29±14.60	34.14±7.79	25.95±4.69	312.69±135.17
<i>M. peltata</i>	312.69±3.87*	229.22±15.07*	128.13±10.16*	76.46±4.14	53.52±2.56	37.76±2.87*	61.99±12.64
Ascorbic acid	29.79±6.40	25.89±2.60	27.18±3.34	28.23±3.24	27.03±2.89	30.23±3.44	

*significant to ascorbic acid

*values obtained from regression lines with 95% of confidence level

Anti-diabetic activity

The alpha-glucosidase (AG) inhibitory activity of the plant extracts was determined using Acarbose as the standard drug (Table 3.) A total of six (6) plants exhibited potent inhibitory activity (>70%) against alpha-glucosidase enzyme. *Hydnocarpus alcalae* belonging to Family Achariaceae showed the highest inhibitory activity (91.67%), the other plants include: *Merremia peltata* belonging to Family Convolvaceae (89.76%); *Trema orientalis* belonging to Family Ulmaceae (84.56%); *Cascabela thevetia* from Family Apocynaceae (79.59%); *Stachytarpheta jamaicensis* from Family Verbenaceae (75.08%); and *Ficus septica* from Family Moraceae (73.93%). As per the results of the phytochemical screening, all these plants contained high amount of total flavonoids (391.39 to 77.11mg/g QE) together with the other phytoconstituents considered in the study, alkaloids and phenols at varying concentration. Interestingly, *H. alcalae* has a total flavonoid content of 329.9mg/g QE and total alkaloid value of 271.06mg/g AE.

Moderate AG inhibition was observed in the extracts of *Cheilocostus speciosus* (67.66%) and *Solanum torvum* (52.45%). Weak inhibitory activity of <50% against AG was recorded in the leaf extracts of *Leea guineensis* (44.11%), *Cinnamomum mercadoi* (32.57%), *Melastoma malabathricum* (21.03%), and *Gmelina arborea* (1.91%). Negative AG inhibitory effect was detected from the extracts of *Dendrocnide meyeniana*, notwithstanding the presence of high amount of total flavonoids (201.22mg/g QE) and total phenols (104.68mg/g GAE) quantified in the phytochemical test.

The different concentration of the ethanol extracts were measured for IC₅₀ at a concentration of 1000, 500, 250, 125, 62.5 and 31.25ug/ml. *S. jamaicensis* has the lowest IC₅₀ value of 0.8ug/ml suggesting that it has the highest activity compared to the other plants and the drug Acarbose with IC₅₀ value of 1.88ug/ml.

Table 3. α -Glucosidase inhibitory effects of 13 selected medicinal plants.

Scientific Name	Family	Inhibition (%)
<i>Gmelina arborea</i>	Lamiaceae	1.91±6.55*
<i>Solanum torvum</i>	Solanaceae	52.45±14.19
<i>Stachytarpheta jamaicensis</i>	Verbenaceae	75.08±1.53
<i>Cascabela thevetia</i>	Apocynaceae	79.59±1.0
<i>Melastoma malabathricum</i>	Melastomataceae	21.03±14.67
<i>Dendrocnide meyeniana</i>	Urticaceae	-0.76±50.84*
<i>Cinnamomum mercadoi</i>	Lauraceae	32.57±14.66
<i>Cheilocostus speciosus</i>	Costaceae	67.66±3.97
<i>Trema orientalis</i>	Ulmaceae	84.56±7.64
<i>Leea guineensis</i>	Vitaceae	44.11±3.05
<i>Ficus septica</i>	Moraceae	73.93±1.07
<i>Hydnocarpus alcalae</i>	Achariaceae	91.67±0.61
<i>Merremia peltata</i>	Convolvulaceae	89.76±1.90
Acarbose		99.47±0.27

*P <0.05 compared to *Acarbose*, data were presented as mean \pm SEM (n = 3). The final concentration of the extracts used in this experiment was 125 μ g/mL.

Anti-obesity activity

The porcine pancreatic inhibitory (PPL) activity of the plant extracts was determined using Orlistat as the standard drug (Table 4). Among the 13 endemics, 7 plants were found to have strong inhibitory activity of >70% against porcine pancreatic lipase (PPL): *Leea guineensis* (89.67%), *Solanum torvum* (89.07%), *Cheilocostus speciosus* (86.15%), *Melastoma malabathricum* (85.36), *Dendrocnide meyeniana* (85.12%), *Cinnamomum mercadoi* (81.35%), and *Gmelina arborea* (71.49%). The strong PPL inhibitory activity is associated with the presence of flavonoids, phenols and alkaloids in these plants, consistent with the results of the phytochemical test. Likewise, study results revealed that, *L. guineensis* which recorded the highest activity against PPL, contains very high amount of alkaloids, flavonoids and phenols and also exhibited strong antioxidant activity in the DPPH assay (Table 2).

Table 4. Lipase inhibitory effects of 13 selected medicinal plants.

Scientific Name	Family	Inhibition (%)
<i>Gmelina arborea</i>	Lamiaceae	71.49±7.52
<i>Solanum torvum</i>	Solanaceae	89.07±4.58
<i>Stachytarpheta jamaicensis</i>	Verbenaceae	57.48±8.14
<i>Cascabela thevetia</i>	Apocynaceae	15.30±20.41*
<i>Melastoma malabathricum</i>	Melastomataceae	85.36±8.50
<i>Dendrocnide meyeniana</i>	Urticaceae	85.12±11.32
<i>Cinnamomum mercadoi</i>	Lauraceae	81.35±2.94
<i>Cheilocostus speciosus</i>	Costaceae	86.15±6.35
<i>Trema orientalis</i>	Ulmaceae	18.28±8.47*
<i>Leea guineensis</i>	Vitaceae	89.67±1.11
<i>Ficus septica</i>	Moraceae	53.00±16.09
<i>Hydnocarpus alcalae</i>	Achariaceae	42.99±2.055*
<i>Merremia peltata</i>	Convolvulaceae	49.90±0.69*
Orlistat		96.97±17.81

*P <0.05 compared to *Orlistat*, data were presented as mean \pm SEM (n = 3). The final concentration of the extracts used in this experiment was 125 μ g/mL.

Previous studies supported the positive correlation between alkaloid, flavonoid and phenolic contents and PPL inhibition activity (Chedda *et al.*, 2016; Adnyana 2014; Gupta *et al.*, 2012). Moderate PPL inhibition was observed in the extracts of *Stachytarpheta jamaicensis* (57.48%), and *Ficus septica* (53%). The extracts of *Merremia peltata* and *Hydnocarpus alcalae*, recorded <50% inhibition of PPL. The 2 plants: *Trema orientalis* and *Cascabela thevetia* exhibited weak inhibitory activity of <20% against porcine pancreatic lipase. Surprisingly, these 2 plants contained sufficient quantity of the major phytoconstituents (Table 1) present in combination of either: high alkaloids/flavonoids and low phenols or high flavonoids/phenols and low alkaloids. The findings of the study propose that high phenolic or alkaloid content and strong antioxidant activity is not necessarily connected with relevant anti-lipase activity. Moreover, plants contain other bioactive components including polyphenols, terpenes, and saponins.

Recent phytochemical studies revealed the presence of many saponins (Marrelli *et al.*, 2016; Hwang *et al.*, 2013; Hernandez-Carlos *et al.*, 2012) and terpenes (Bustanji *et al.*, 2011) with anti-lipase properties.

The different concentration of the plant extracts were measured for IC₅₀ at a concentration of 1000, 500, 250, 125, 62.5 and 31.25ug/ml. The IC₅₀ is defined simply as the inhibitor concentration that decreases the biotransformation of a substrate at a single, specified concentration by 50% (Pharmacology, 2009). The extracts of *M. malabathricum*, *D. meyeniana*, *C. mercadoi*, *G. arborea*, *C. speciosus*, *S. torvum*, and *L. guineensis* had IC₅₀ values of 1.03, 1.07, 1.19, 1.2, 1.31, 1.35, and 2.14ug/ml, respectively. Whereas, *Orlistat* had IC₅₀ value of 1.81ug/ml. *M. malabathricum* has the lowest IC₅₀ suggesting that it has the highest activity compared to the other plants and the drug *Orlistat*.

Conclusion and recommendations

The thirteen (13) endemic medicinal plants collected from Mount Mayon and Mount Malinao Albay showed strong *in vitro* antioxidant capacity consistent with the results of the phytochemical screening. The general assessment of the analytical results for the plant extracts definitely shows the individual specificity of each plant sample and a broad range of alkaloid, flavonoid and phenolic compounds. To the best of our knowledge, the plants studied in this paper have not been screened earlier for *in vitro* biological activities. Results of the study propose that all the plant extracts can be considered as good source of natural antioxidants. Six plants are potent alpha glucosidase inhibitors and seven plants are strong inhibitors of porcine pancreatic lipase.

Conflict of interest

The authors declare that there was no conflict of interest

Acknowledgements

This work was funded by Department of Science and Technology-Philippine Council for Health Research and Development (DOST-PCHRD).

References

- Aboaba SA, Choudhary IM.** 2015. Chemical composition and biological activities of the volatile oils of *Palisota hirsuta* (Thunb) K. Schum and *Trema orientalis* (L) Blume. International Journal of Chemistry **Vol. 7**, No.
- Ahmed D, Fatima M, Saeed S.** 2014. Phenolic and flavonoid contents and anti-oxidative potential of epicarp and mesocarp of *Lagenaria siceraria* fruit: a comparative study. Asian Pacific Journal of Tropical Medicine **7 (Suppl 1)**, S249-S255.
- Ajaiyeoba EO.** 1999. Comparative phytochemical and antimicrobial studies of *Solanum macrocarpum* and *Solanum torvum* leaves, Fitoterapia **70**, 184-186.
- Al –Kattan MO, Khayat SA.** 2017. Antimicrobial activity and Chemical analyses of oil constituents of Medicinal Plant *Costus speciosus* (Koen.) Biomedical Research **28(2)**, 734-739.
- Alvarez E, Leiro JM, Rodríguez M, Orallo F.** 2004. Inhibitory effects of leaf extracts of *Stachytarpheta jamaicensis* (Verbenaceae) on the respiratory burst of rat macrophages. Phytotherapy Research **18(6)**, 457-462.
- Ambujakshi HR, Heena T.** 2009. Anthelmintic activity of *Gmelina arborea* leaves extract. International Journal of Pharmaceutical Research and Development **1(9)**, 1-3.
- Andrade-Cetto A, Becerra-Jimenez J, Cardenas-Vazquez R.** 2008. Alfa- glucosidase-inhibiting activity of some Mexican plants used in the treatment of type 2 diabetes. Journal of Ethnopharmacology **116**, 27-32.
- Arthan D, Svasti J, Kittakoop P, Pittayakhachonwut D, Tanticharoen M, Thebtaranonth Y.** 2002. Antiviral isoflavonoid sulfate and steroidal glycosides from the fruits of *Solanum torvum*. Phytochemistry **59**, 459-463.
- Atanasov AG, Waltenberger B, Pferschy-Wenzig EM, Linder T, Wawrosch, C, Uhrin P, Stuppner H.** 2015. Discovery and resupply of pharmacologically active plant-derived natural products: A review. Biotechnology Advances **33(8)**, 1582-1614.

- Babatunde S, Michael WO, Oyindamola A.** 2015. Bio-guided investigation of the antimalarial activities of *Trema orientalis* (L.) Blume leaves. African Journal of Biotechnology **14(43)**, 2966-2971.
- Bhuyan B, Zaman K.** 2008. Evaluation of hepatoprotective activity of rhizomes of *Costus speciosus* (J. Konig.) Smith. Pharmacology online **4(3)**, 119-126.
- Brewer M.** 2011. Natural antioxidants: sources, compounds, mechanism of action and potential applications. Comprehensive reviews in food science and food safety **10**, 221-247.
- Bueno PR, Buno CB, Santos DL, Santiago LA.** 2013. Antioxidant activity of *Ficus pseudopalma* Blanco and its cytotoxic effect on hepatocellular carcinoma and peripheral blood mononuclear cells. Current Research in Biological and Pharmaceutical Sciences **2(2)**, 14-21.
- Carriere F, Gargouri Y, Moreau H, Ransac S, Rogalska E, Verger R in: Wooley P, Petersen SB. (Eds.).** 1994. Lipases, their structure, biochemistry and application, Book Cambridge University Press pp. 181-205.
- Chan MFE, Geronimo AJO, Aspiras APF, Busing EJW, Dato RJB, Calumpit DMA, Lafuente AT.** 2016. Analysis of phytochemical, antimicrobial, and antioxidant properties of *Sarcandra glabra* (Thunb.) Nakai in relation to its ethnomedicinal relevance in Cordillera, Philippines. Indian Journal of Traditional Knowledge **15(3)**, 411-416.
- Chang YT, Shen JJ, Wing WR, Yen HR.** 2009. Alternative therapy for autosensitization dermatitis. Chang Gung Medicinal Journal **32(6)**, 668-673.
- Chanmee W, Chaicharoenpong C, Petsom A.** 2013. Lipase Inhibitor from fruits of *Solanum stramonifolium* Jacq. Food and Nutrition Sciences **4**, 554-558.
- Convention on Biological Diversity.** 2009. Assessing progress towards the 2010 biodiversity target: The 4th National Report to the Convention on Biological Diversity The 4th National Report to the Convention on Biological Diversity.
- Dechakhamphu A, Wongchum N.** 2015. Screening for anti-pancreatic lipase properties of 28 traditional Thai medicinal herbs. Asian Pacific Journal of Tropical Biomedicine **5**, 1042-1045.
- Floegel A, Kim DO, Chung SJ, Koo SI, Chun OK.** 2011. Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. Journal of Food Composition and Analysis **24**, 1043-1048.
- Florido LV.** 1978. Vegetative propagation by cuttings of yemane (*Gmelina arborea* Roxb.) using growth hormones. Sylvatrop Journal **3(2)**, 115-122.
- Food and Nutrition Research Institute (FNRI).** 2011. 2nd National nutrition summit: 8th National nutrition survey. Food and Nutrition Research Institute Department of Science and Technology.
- Fuentes RG, Diloy FN, Tan IL, Balanquit BJ.** 2010. Antioxidant and Antibacterial Properties of Crude Methanolic Extracts of *Cinnamomum mercadoi* Vidal. Phil Journal of Natural Sciences **15**, 9-15.
- Galvez MA.** 2015. Evaluation of DPPH free radical scavenging activity and phytochemical screening of selected folkloric medicinal plants in Tinoc, Ifugao, Cordillera Administrative Region, Philippines. International Journal of Scientific Research Publication **5(12)**, 440-445.
- Gandhi GR, Ignacimuthu S, Paulrajmg, Sasikumar P.** 2011. Antihyperglycemic activity and antidiabetic effect of methyl caffeate isolated from *Solanum torvum* Swartz. fruit in streptozotocin induced diabetic rats, European Journal of Pharmacology **30(23)**, 623-31.
- Gerrath JM, Lacroix CR.** 1997. Heteroblastic Sequence and Leaf Development in *Leea guineensis*. International Journal of Plant Sciences **158(6)**, 747-756. DOI: 10.1086/297486.

- Gondoin A, Grussu D, Stewart D, McDougall GJ.** 2010. White and green tea polyphenols inhibit pancreatic lipase in vitro. *Food Research International* **43**, 1537-1544.
- Gorgonio SR, Fuentes RG.** 2011. Antidiarrheal Activity of *Cinnamomum mercadoi* methanolic leaf and bark extracts. *Philippine Journal of Natural Sciences* **16-1**, 43-47.
- Gutiérrez SP, Sánchez MAZ, Gonzlález CP, García LA.** 2007. Antidiarrhoeal activity of different plants used in traditional medicine. *African Journal of Biotechnology* **6(25)**, 2988-2994.
- Hamuel J.** 2012. Phytochemicals: Extraction methods, basic structures and mode of action as potential chemotherapeutic Agents, in Rao, V. (Ed.) *Phytochemicals - A Global Perspective of Their Role in Nutrition and Health*. InTech Open p. 538.
DOI: 10.5772/26052
- Hayes JD, Eggleston IM.** 2008. The cancer chemopreventive actions of phytochemicals derived from glucosinolates. *European Journal of Nutrition* **47 Suppl 2**, 73-88.
- Hu L, Yu W, Li Y, Prasad N, Tang Z.** 2014. Antioxidant activity of extract and its major constituents from okra seed on rat hepatocytes injured by carbon tetrachloride. *Biomed Research International* **2014**, 1-9.
- Idu M, Erhabor JO, Odia EA.** 2009. Morphological and anatomical studies of the leaf and stem of some medicinal plants: *Stachytarpheta jamaicensis* (L.) Vahl. And *S. cayennensis* (LC Rich) schau. *Ethnobotanical Leaflets* **13(11)**, 417-1425.
- Ikeda K, Takahashi M, Nishida M.** 2000. Homonojirimycin analogues and their glucosides from *Lobelia sessilifolia* and *Adenophora* spp. (Campanulaceae), *Carbohydrate Research* **323**, 73-80.
- Ikewuchi JC, Okaraonye CC, Ogbonnaya EA.** 2009. Time course of the effect *Stachytarpheta jamaicensis* L. (Vahl.) on plasma sodium and potassium levels of normal rabbits. *Journal of Applied Sciences Research* **5(10)**, 1741-1743.
- Institute for Clinical Systems (ICSI).** 2013. Prevention and management of obesity for children and adolescents. Bloomington: Institute for Clinical Systems Improvement; 2013 [Accessed on 31 January 2018]. p. 41. Available at: www.icsi.org/_asset/tn5cd5/ObesityChildhood.pdf.
- International Union for Conservation of Nature (IUCN).** 2015. Numbers of threatened species by major groups of organisms (1996-2015) http://cmsdocs.s3.amazonaws.com/summarystats/2015_2_Summary_Stats_Page_Documents/2015_2_RL_Stats_Table_1.pdf.
- Jadav HR, Galib R, Harish CR, Kumar PP.** 2016. Preliminary Pharmacognostical profile of Tuvaraka (*Hydnocarpus laurifolia* (Dennst) Sleummer.) seeds. *Medical Journal of Dr. D.Y. Patil University* **9(2)**, 219-223.
- Jagadish NRN, Gopalkrishna B.** 2008. Evaluation of analgesic activity of different extracts of *Stachytarpheta indica* L. (Vahl). *Biomed* **3(3-4)**, 229-233.
- Jaiswal BS.** 2012. *Solanum torvum*: A review of its traditional uses, phytochemistry and pharmacology. *International Journal of Pharma and Bio Sciences* **3(4)**, 104-111.
- Kandakumar S, Sathya V, Manju V.** 2014. Synthesis and characterization of silver nanoparticle using *Hydnocarpus alpine*, its application as a potent antimicrobial and antioxidant agent- A novel study. *International journal of Chem Tech Research* **6(11)**, 4770-6.
- Kang JG, Park CY.** 2012. Anti-obesity drugs: a review about their acts and safety," *Diabetes & Metabolism Journal* **36(1)**, 13-25.

- Kar A.** 2007. Pharmacognosy and pharmaco biotechnology. 2nd edn. New Delhi: New Age International (P) Ltd.
- Kareru PG, Keriko JM, Kenji GM, Gachanja AN.** 2010. Anti-termite and antimicrobial properties of paint made from *T. peruviana* (pers.) Schum. oil extract. African Journal of Pharmacy and Pharmacology **4(2)**, L087- 089.
- Karthikeyan R, Solaimuthu C, Balakrishnan N.** 2013. A study of performance and emissions of diesel engine fuelled with neat diesel and neat *Hydnocarpus pentandra* biodiesel. IOSR Journal of Mechanical and Civil Engineering **10(2)**, 53-57.
- Katalinic V, Milos M, Kulisic T, Jukic M.** 2006. Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. Food Chemistry **94**, 550-557.
- Kaur S, Mondal P.** 2014. Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants. Journal of Microbiology and Experimentation **1(1)**, 1-6.
- Kimura K, Lee JH, Lee IS, Lee HS, Park KH, Chiba S.** 2004. Two potent competitive inhibitors discriminating alpha-glucosidase family I from family II. Carbohydrate Research **339**, 1035-40.
- Kopelman PG.** 2000. Obesity as a medical problem. Nature **404 (6778)**, 635-43.
- Kumar S, Narwal S, Kumar V, Prakash O.** 2011. α -glucosidase inhibitors from plants: A natural approach to treat diabetes. Pharmacognosy Review, Jan-Jun **5(9)**, 19-29. doi: 10.4103/0973-7847.79096
- Lacroix CR, Gerrath JM, Posluszny U.** 1990. The Developmental morphology of *Leea guineensis*. I. Vegetative Development. Botanical Gazette **151(2)**, 204-209. DOI: 10.1086/337819.
- Lahlou M.** 2007. Screening of natural products for drug discovery. Expert Opinion on Drug Discovery **2(5)**, 697-705. DOI: 10.1517/17460441.2.5.697.
- Lanting MV, Palaypayon CM.** 2002. Forest tree species with medicinal uses. DENR Recommends Vol. 11. Ecosystems Research and Development Bureau, Department of nvironment and Natural Resources, College, Laguna p24.
- Latayada FS, Uy MM.** 2016. Screening of the antioxidant properties of the leaf extracts of Philippine medicinal plants *Ficus nota* (Blanco) Merr., *Metroxylon sagu* Rottb., *Mussaenda philippica* A. Rich., *Inocarpus fagifer*, and *Cinnamomum mercadoi* Vidal. Bulletin of Environment, Pharmacology and Life Sciences **5**, 18-24.
- Lawag IL, Aguinaldo AM, Naheed S, Mosihuzzaman M.** 2012. α -Glucosidase inhibitory activity of selected Philippine plants. Journal of Ethnopharmacology **144(1)**, 217-9.
- Lawrence L, Menon S, Vincent S, Sivaram VP, Padikkala V.** 2016. Radical scavenging and gastroprotective activity of methanolic extract of *Gmelina arborea* stem bark. Journal of Ayurveda and Integrative Medicine **7**, 78-82.
- Lebovitz HE.** 1997. α -Glucosidase inhibitors. Endocrinology and Metabolism Clinics of North America **26**, 539-55.
- Lee EM, Lee SS, Chung BY, Cho JY, Lee IC, Ahn SR.** 2000. Pancreatic lipase inhibition by C-glycosidic flavones isolated from *Eremochloa ophiuroides*. Molecules **15**, 8251-9.
- Liew PM, Yong YK.** 2016. *Stachytarpheta jamaicensis* (L.) Vahl: From traditional usage to pharmacological evidence. Evidence-Based Complementary Alternative Medicine. 7842340. <https://doi.org/10.1155/2016/7842340>
- Liu H, Qiu N, Ding H, Yao R.** 2008. Polyphenols contents and antioxidant capacity of 68 Chinese herbals suitable for medical or food uses. Food Research International **41(4)**, 363-370.
- Luna B, Pharm D, Feinglos MN.** 2001. Oral agents in the management of Type 2 Diabetes Mellitus. American Family Physician **63**, 1747-56, 1759-80.

- Lunagariya NA, Patel NK, Jagtap SC, Bhutani KH.** 2014. Inhibitors of pancreatic lipase: state of the art and clinical perspectives. *Experimental and Clinical Sciences (EXCLI J)* **13**, 897-921.
- Lynn A, Collins A, Fuller Z, Hillman K, Ratcliffe B.** 2006. Cruciferous vegetables and colorectal cancer. *Proceedings of the Nutrition Society* **65(1)**, 135-44.
- Ma J, Liu SJ.** 2006. Research progress of DNJ in mulberry twig *Food Science Technology* **9**, 112- 114.
- Macabeo APG, Lopez ADA, Schmidt S, Heilmann J, Dahse HM, Alejandro GJD.** 2013. Antitubercular and cytotoxic constituents from *Goniothalamus gitingensis*. *Natural Products* **8(1)**, 41-45.
- Maqsood M, Dildar Ahmed D, Atique I, Malik W.** 2017. Lipase inhibitory activity of *Lagenaria siceraria* fruit as a strategy to treat obesity. *Asian Pacific Journal of Tropical Medicine* **10(3)**, 305-310.
- Maurya S, Kushwaha A, Singh S, Singh G.** 2014. An overview on antioxidative potential of honey from different flora and geographical origins. *Indian Journal of Natural Products and Resources* **5**, 9-19.
- Mazura MP, Susanti D, Rasadah MA.** 2007. Anti-inflammatory action of components from *Melastoma malabathricum*. *Pharmaceutical Biology* **45(5)**, 372-375.
- McDougall GJ, Kulkarni NN, Stewart D.** 2009. Berry polyphenols inhibit pancreatic lipase activity in vitro. *Food Chemistry* **115**, 193-199.
- Moreno DA, Ilic N, Poulev A, Brasaemle DL, Fried SK, Raskin I.** 2003. Inhibitory effects of grape seed extract on lipases. *Nutrition* **19**, 876-879.
- Nahak G, Kantasahu R.** 2011. Free radical scavenging activity of rhizome of *Costus speciosus* (Koen). *International Journal of Institute of Pharmacy and Life Sciences* **1**, 62-67.
- Ndebia EJ, Kanga R, Nchunga-Anye Nkeh B.** 2007. Analgesic and anti-inflammatory properties of aqueous extract from leaves of *Solanum torvum* (Solanaceae). *African Journal of Traditional and Complementary Medicine* **4(2)**, 240-244.
- Ng M, Fleming T, Robinson M.** 2014. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: A systematic analysis for the Global Burden of Disease Study 2013. *The Lancet* **384(9945)**, 766-781.
- Oke JM, Hamburger MO.** 2002. Screening of some Nigerian medicinal plants for antioxidant activity using 2, 2, Diphenyl-Picryl-Hydrazyl radical. *African Journal of Biomedical Research* **5**, 77-79.
- Oladunmoye MK, Kehinde FY.** 2011. Ethno botanical survey of medicinal plants used in treating viral infections among Yoruba tribe of south west Nigeria. *African Journal of Microbiology Research* **5**, 2991-3004.
- Patil RK, Makari HK, Gurumurthy H.** 2008. *In vitro* antimicrobial activity of ethanol extract of *Thevetia peruviana*, ejeafche. *Biotechnology An Indian Journal* **2(1)**, 5-7.
- Perez KJ, Jose MA, Aranico E, Madamba RZ.** 2015. Phytochemical and antibacterial properties of the ethanolic leaf extract of *Merremia Peltata* (L.) Merr. and *Rubus* sp. *Advances in Environmental Biology* **9(19)**, 50-56.
- Perk J.** 2017. Non-communicable diseases, a growing threat to global health. *E-Journal of Cardiology Practice* **15(14)**.
- Peteros NP, Uy MM.** 2010. Antioxidant and cytotoxic activities and phytochemical screening of four Philippine medicinal plants. *Journal of Medicinal Plants Research* **4(5)**, 407-414.
- Pharmacology.** 2009. Drug-drug interactions with an emphasis on drug metabolism and transport, principles and practice. Chapter **12**, 303-325.

- Playford RJ, Pither C, Gao R.** 2013. Use of the α -glucosidase inhibitor acarbose in patients with 'Middleton syndrome': normal gastric anatomy but with accelerated gastric emptying causing postprandial reactive hypoglycemia and diarrhea, Canadian Journal of Gastroenterology **27**, 403-404.
- Poddar K, Kolge S, Bezman L, Mullin GE, Cheskin LJ.** 2011. Nutraceutical supplements for weight loss: a systematic review. Nutrition Clinical Practice **26(5)**, 539-52.
- Putera I, Shazura KA.** 2010. Antimicrobial activity and cytotoxic effects of *Stachytarpheta jamaicensis* (L.) Vahl crude plant extracts [Master dissertation], Universiti Teknologi Malaysia.
- Quisumbing E.** 1978. Medicinal Plants of the Philippines, Katha Publishing, Manila.
- Raga DD, Espiritu RA, Shen CC, Ragasa CY.** 2011. A bioactive sesquiterpene from *Bixa orellana*. Journal of Natural Medicines **65(1)**, 206-211.
- Ragasa CY, Macuha MR, De Los Reyes MM, Mandia EH, Van Alenas IA.** 2016. Chemical constituents of *Ficus septica* Burm. F. International Journal of Pharmaceutical and Clinical Research **8(11)**, 1464-1469.
- Ragasa CY, Tsai P, Shen CC.** 2009. Terpenoids and sterols from the endemic and endangered Philippine trees, *Ficus pseudopalma* and *Ficus ulmifolia*. Philippine Journal of Science **138(2)**, 205-209.
- Rajbhar N, Kumar A.** 2014. Pharmacological importance of *Thevetia peruviana*. International Journal of Pharmaceutical and Chemical Sciences **3(1)**, 260-263.
- Regional Development Plan (RDP).** 2011. Providing for basic needs: health. Bicol Regional Development Plan 2011-2016 pp. 45.
- Roberts DM, Southcott E, Potter JM, Roberts MS, Eddleston M, Buckley NA.** 2006. Pharmacokinetics of digoxin cross reacting substances in patients with acute yellow oleander (*T. peruviana*) poisoning, including the effect of activated charcoal. Therapeutic Drug Monitoring **28(6)**, 784-792.
- Ruma OC, Zipagang TB.** 2015. Determination of secondary metabolites and antibacterial property of extract from the leaves of *Stachytarpheta jamaicensis* (L.) Vahl. Journal of Medicinal Plants and Studies **3(4)**, 79-8.
- Saeed N, Khan MR, Shabbir M.** 2012. Antioxidant activity, total phenolic and total flavonoids contents of whole plant extracts *Torilis leptophylla* L. BMC Complementary and Alternative Medicine **12**, 221.
- Sahu RK, Kar M, Routay R.** 2013. DPPH free radical scavenging activity of some leafy vegetables used by Tribals of Odisha, India. Journal of Medicinal Plant Studies **1**, 21-27.
- Saroya AS.** 2011. Herbalism, Phytochemistry and Ethnopharmacology. Enfield, New Hampshire: Science Publishers.
- Schaich KM, Tian X, Xie J.** 2015. Hurdles and pitfalls in measuring antioxidant efficacy: a critical evaluation of ABTS, DPPH, and ORAC assays. Journal of Functional Foods **14**, 111-125.
- Seyedan A, Alshawsh MA, Alshagga MA, Koosha S, Mohamed Z.** 2015. Medicinal plants and their inhibitory activities against pancreatic lipase: A Review. Evidence-Based Complementary and Alternative Medicine **1-13**.
- Shibano M, Kakutani K, Taniguchi M.** 2008. Antioxidant constituents in the dayflower (*Commelina communis* L.) and their α -glucosidase-inhibitory activity. Journal of Natural Medicine **62**, 349-353.
- Singleton VL, Orthofer R, Lamuela-Raventos RM.** 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods of Enzymology **299**, 152-178.
- Sivapriya M, Srinivas L.** 2007. Isolation and purification of a novel antioxidant protein from the water extract of Sundakai (*Solanum torvum*) seeds. Food Chemistry **104**, 510-517.

- Sjostrom L, Rissanen A, Andersen T, Boldrin M, Golay A, Koppeschaar HPF, Krempf M.** 1998. Randomised placebo-controlled trial of orlistat for weight loss and prevention of weight regain in obese patients. *Lancet* **352**, 167-172.
- Sravani P, Murali CM, Sadik BS, Soubia SN.** 2011. Evaluation of antidiuretic activity of *Gmelina arborea*. *International Journal of Advances in Pharmacological Research* **2(4)**, 157-161.
- Srinivasan GV, Sharanappa P, Leela NK, Sadashiva CT, Vijayan KK.** 2009. Chemical composition and antimicrobial activity of *Leea indica* (Burm. F.) Merr flowers. *Natural Product Radiance* **8(5)**, 488-493.
- Srivastava N, Chauhan AS, Sharma B.** 2012. Isolation and characterization of some phytochemicals from Indian traditional plants. *Biotechnology Research International*, 1-8.
- Straughan DW, Fentem JE, Balls M.** 1996. Replacement alternative and complimentary in vitro methods in pharmaceutical research. Pages 1-13 in JV Castel and MJG Lechon, Eds. *In vitro* methods in pharmaceutical research. Academic press.
- Sulaiman CT, Balachandran I.** 2012. Total phenolics and total flavonoids in selected Indian medicinal plants. *Indian Journal of Pharmaceutical Science* **74(3)**, 258-260.
- Sulaiman MR, Zakaria ZA, Chiong HS, Lai SK, Israf DA, Azam Shah TM.** 2009. Antinociceptive and anti-inflammatory effects of *Stachytarpheta jamaicensis* (L.) Vahl (Verbenaceae) in experimental animal models. *Medical Principles and Practice* **18(4)**, 272-279.
- Tabopda TK, Ngoupayo J, Awoussong PK.** 2008. Triprenylated flavonoids from *Dorstenia psilurus* and their α -glucosidase inhibition properties. *Journal of Natural Products* **71**, 2068-2072.
- Thaipong K, Boonprakob U, Crosby K, Cisneros-Zevallos L, Byrne D.** 2006. Comparison of ABTS, DPPH, FRAP and ORAC assays for estimating the antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis* **19**, 669-675.
- Thomas RP, Thomas M, Paul J, Mohan M.** 2013. Antifungal activity of Verbenaceae. *Biosciences Biotechnology Research Asia* **10(1)**, 355-360.
- Tiwari BK, Brunton NP, Brennan CS.** 2013. *Handbook of Plant Food Phytochemicals: Sources, Stability and Extraction*. Wiley-Blackwell. ISBN: 978-1-444-33810-2.
- Torres RC, Sison FM, Israel MC.** 2003. Phytochemical screening and biological studies on the crude methanol extract of *Cinnamomum mercadoi*, Vidal. *Philippine Journal of Science* **132(2)**, 27-32.
- Varghese B, Sandhya S, Kavitha MP, Krishnakumar K.** 2016. Genus *Hydnocarpus*: A Review. *International Journal of Phytomedicine* **7(3)**, 143-154.
- Vital PG, Velasco, RNJ, Demigillo JM, Rivera WL.** 2010. Antimicrobial activity, cytotoxicity and phytochemical screening of *Ficus septica* Burm and *Sterculia foetida* L. leaf extracts. *Journal of Medicinal Plants Research* **4(1)**, 58-63.
- Weigle DS.** 2003. Pharmacological therapy of obesity: past, present, and future. *Journal of Clinical Endocrinology and Metabolism* **88(6)**, 2462-2469.
- Yahia EM (ed.)** 2018. *Fruit and Vegetable Phytochemicals: Chemistry and Human Health*. 2nd Editio. John Wiley & Sons.
- Yin Z, Zhang W, Feng F, Zhang Y, Kang W.** 2014. α -Glucosidase inhibitors isolated from medicinal plants. *Food Science and Human Wellness* **3**, 136-174.
- Yun JW.** 2011. Possible anti-obesity therapeutics from nature – A review. *Phytochemistry* **71**, 1625-41.