



Cytotoxic effect of ethanolic extract of *Aloe castellorum* on human breast carcinoma MCF-7 cells

Awad A. Algarni

Department of Biology, College of Science and Arts, Albaha University, Baljurashi, Saudi Arabia

Key words: *Aloe castellorum*; Cytotoxicity, Cancer, Cell line, Anticancer activity.

<http://dx.doi.org/10.12692/ijb/19.4.162-165>

Article published on October 27, 2021

Abstract

Cancer is the world's most serious disease. New and safe anticarcinogens can be found in natural products. *Aloe castellorum* is an Aloe plant species native to Saudi Arabia. In the present study, the cytotoxic activity of the methanol extract of *Aloe castellorum* leaves was evaluated using breast adenocarcinoma cells (MCF-7), assessed by sulforhodamine B (SRB) cell viability assays. The results indicate that the *Aloe castellorum* showed weak cytotoxic effects against tested cancer cell lines, with an IC_{50} value of $341\mu\text{g/ml}$. In conclusion, the in vitro cytotoxic evaluation of *Aloe castellorum* showed low cytotoxicity on MCF-7 cell line.

* **Corresponding Author:** Awad A Algarni ✉ awad-algarni-j@hotmail.com

Introduction

Breast cancer affects women all over the world more than any other form of cancer. After lung cancer, breast cancer is the second leading cause of mortality (Ferlay *et al.*, 2015). Breast cancer accounts for 28.7% of all newly diagnosed cancers in Saudi women, which makes it the most prevalent cancer in Saudi Arabia (Ibrahim *et al.*, 2008). Current constraints in cancer treatment include toxic effects and tumor resistance to chemotherapeutic agents. Therefore, finding new and more effective treatment options is considered a priority (Gordaliza, 2007). Plant extracts are rich in compounds such as flavonoids and polyphenols, which play an important role in drug discovery and development (Newman and Cragg, 2012). *Aloe castellorum* is an Aloe plant species native to Saudi Arabia. Several studies indicate that certain species of *Aloe* have antioxidant and anti-carcinogenic properties (Majumder *et al.*, 2019). Therefore, this study aimed to investigate the cytotoxic effect of the methanolic extract of *Aloe castellorum* against human breast carcinoma MCF-7 cells.

Materials and methods

Plant Material and Extraction

Samples of the *Aloe castellorum* were collected in August 2020 from the Albaha region of Saudi Arabia. The authenticity and identification of the plant were made by taxonomist Abdulwali Al-Khulaidi, Department of Biology, University of Albaha.

The plant samples were dried in a vacuum oven at 40 °C for 72 h and then ground with a mortar and pestle, which were extracted with 70 % methanol (3 × 60 mL fresh solvent, 30 min ultrasonic each). Then, the mixture was filtered and the extract was collected and evaporated under vacuum at 50°C.

Cancer cell culture

Human breast carcinoma cells (MCF-7) were received from Nawah Scientific Inc. (Egypt) and stored in the laboratory. Cell lines were cultured in a DMEM medium containing 100 mg/mL streptomycin, 100 U/mL penicillin and 10% FBS at 37°C in an incubator humidified with 5% CO₂.

SRB assay

The viability of the cells was assessed by colorimetric sulphorhodamine-B (SRB) assay. In brief, 4 × 10⁴ MCF-7 cells were added to 6-well plates and incubated for 24 hours in complete media. Cells were treated with another aliquot of 100 µL media containing plant extract at 0-3000 µg/ml concentrations. After 72 h, cells were fixed with ice-cold 10% TCA at 4 °C for 1 h. The TCA solution was removed, and the cells were washed 5 times with distilled water. Aliquots of 70 µL SRB solution (0.4% w/v) were added and incubated in a dark place at room temperature for 10 min.

After three washes with 1% acetic acid, plates were air-dried overnight. After that, In order to dissolve the protein-bound SRB stain, 150 µL of TRIS (10 mM) was added. Using a BMG LABTECH® - FLUOstar Omega microplate reader, the absorbance was measured at 540 nm.

Statistical Analysis

Using one-way ANOVA, the results were reported as mean ± standard deviation (Mean ± SD). P-values < 0.05 indicate significant differences in the data.

Results and discussion

The cytotoxic effect of the methanol extract of *Aloe castellorum* on human breast adenocarcinoma (MCF-7) was carried out using the SRB assay.

The results indicate that the *Aloe castellorum* showed weak cytotoxic effects against MCF-7 cell lines, with an IC₅₀ value of 341 µg/ml (Fig. 1). *In vitro*, plant extract produced limited reductions in cell viability and limited cytotoxicity.

These results are in line with other studies reporting weak cytotoxicity of other *Aloe* species. *Aloe niebuhriana* latex extract showed a weak cytotoxic effect against MCF-7, HepG2 and HCT-116 tumor cells (Moharram *et al.*, 2020). According to results from MTT assay, Fox *et al.* (2017) found that *Aloe vera*, *Aloe ferox*, and *Aloe marlothii* had weak cytotoxic effects on HaCaT cells. Furthermore, some

studies showed the *in vitro* activity of some extracts of *Aloe* on survival of human-cell lines was not significant. In 2014, the study of DuPlessis and Hamman included *Aloe vera*, *Aloe marlothii*, *Aloe*

speciosa and *Aloe ferox*, which all have not recorded any activity on SH-SY5Y (human neuroblastoma), HepG2 (human hepatocellular carcinoma) and HeLa (human epithelial adenocarcinoma).




Fig. 1. Effect of *Aloe castellanum* on cell viability and growth of MCF-7 cells.

Conclusion

In this study, we tested the cytotoxic activity of *Aloe castellanum* against the MCF-7 cell line using the SRB assay. The results show that the methanolic extract of the plant leaves has a weak cytotoxic effect on the tested cancer cell line.

Conflict of interest

The author declares that there is no conflict of interest.

Funding

This research work has not received any funding at all in any part of the study.

Acknowledgments

The author would like to thank the Department of Biology, College of Science and Arts, Albaha University, Baljurashi, Saudi Arabia.

References

Du Plessis LH, Hamman JH. 2014. *In vitro* evaluation of the cytotoxic and apoptogenic

properties of *aloe* whole leaf and gel materials. Drug and chemical toxicology **37(2)**, 169-177.

<https://doi.org/10.3109/01480545.2013.834356>

Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. 2015. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. International journal of cancer **136(5)**, E359-E386.

<https://doi.org/10.1002/ijc.29210>

Fox LT, Mazumder A, Dwivedi A, Gerber M, Du Plessis J, Hamman JH. 2017. *In vitro* wound healing and cytotoxic activity of the gel and whole-leaf materials from selected *aloe* species. Journal of ethnopharmacology **200**, 1-7.

<https://doi.org/10.1016/j.jep.2017.02.017>

Gordaliza M. 2007. Natural products as leads to anticancer drugs. Clinical and Translational Oncology **9(12)**, 767-776.

<https://doi.org/10.1007/s12094-007-0138-9>

Ibrahim EM, Zeeneldin AA, Sadiq BB, Ezzat AA. 2008. The present and the future of breast cancer burden in the Kingdom of Saudi Arabia. *Medical Oncology* **25(4)**, 387-393.

<https://doi.org/10.1007/s12032-008-9051-5>

Majumder R, Das CK, Mandal M. 2019. Lead bioactive compounds of *Aloe vera* as potential anticancer agent. *Pharmacological research* **148**, 104416.

<https://doi.org/10.1016/j.phrs.2019.104416>

Moharram BA, Al-Doaiss AA, Al-Mahbashi HM, Al-Kahtani MA, Alfaifi MA, Elbehairi SE, Saif-Ali R. 2020. Hepatoprotective, Antioxidant and

Cytotoxic Potential of *Aloe niebuhriana* Latex Extract from Yemen. *Indian Journal of Pharmaceutical Sciences* **82(3)**, 514-521.

<https://doi.org/10.36468/pharmaceutical-sciences.675>

Newman DJ, Cragg GM. 2012. Natural products as sources of new drugs over the 30 years from 1981 to 2010. *Journal of natural products* **75(3)**, 311-335.

<https://doi.org/10.1021/acs.jnatprod.5b01055>