



Solid state fermentation of *Lathyrus sativus* and sugarcane bagasse by *Pleurotus sajor-caju*

Md. Rezaul Karim¹, Zinat Mahal², Safia Iqbal¹, Harun or Rashid², M. Abu Hena Mostofa Jamal¹, Md. Azizul Islam¹, Md. Mafizur Rahman^{1*}

¹Department of Biotechnology and Genetic Engineering, Islamic University, Kushtia, Bangladesh

²Institute of Food and Radiation biology, Atomic Energy Commission, Bangladesh

Article published on May 15, 2014

Key words: *Pleurotus sajor-caju*, lignocellulosic, Sugarcane bagasse, Fermentation, *Lathyrus sativus*.

Abstract

Lathyrus sativus (Khesari plant) and sugarcane bagasse are considered as agro wastes. Khesari plants are generally grown in fields as weeds and they have to be removed prior to cultivation. Taking this view in mind, we have investigated the conversion of these lignocellulosic agro-waste as an enriched feed stock for cattle via by solid state fermentation using a cellulolytic fungus, *Pleurotus sajor-caju*. The strain required 8 weeks to complete the fermentation on both the untreated and treated (with alkali, lime and presoaked) substrates at 30°C. Higher amounts of reducing sugar and soluble protein were found in each of the lime treated substrates than untreated substrates. Results also indicated that presoaked substrate contain higher amounts of reducing sugar and soluble protein than unsoaked substrate. Among the substrates, mixed substrate (khesari plant + sugarcane bagasse) was found to accumulate higher amount of sugar, 22.15 mg/g and protein, 22.80 mg/g than those of khesari plant in the 5th week of fermentation. The treatments that augmented the level of sugar and protein were also found to enhance the cellobiase, carboxy methyl cellulase and avicelase activity of crude culture extracts. These results suggest that lime treatment and presoaking seem to increase the digestibility of the substrates by the fungal cellulolytic enzymes. During eight weeks of fermentation, relatively higher cellobiase activity was found as compared to that of carboxymethylcellulase and avicelase at 30°C for the fungul strain. The results of the present study clearly indicate that fungal conversion with pretreatment transform these lignocellulosic agro-wastes to a nutritionally enriched animal feed.

*Corresponding Author: Md. Mafizur Rahman ✉ shilon83@gmail.com

Introduction

In developing agro-based countries like Bangladesh, farmers use cattle for farming. Due to financial constraints, farmers are not always able to provide their cattle with high quality feed. Consequently, most of them suffer from nutritional deficiency that give rise to different diseases. To tackle this issue it is imperative to transform a cattle feed into rich nutrients source (Hossain *et al.*, 2009).

Agricultural crop residues and/or byproducts are staple energy substrate of ruminant livestock in developing countries (Tripathi *et al.*, 2008). In Bangladesh, *Lathyrus sativus* (Khesari plant) and sugarcane bagasse are two agricultural wastes mainly consisting of lignin, cellulose and hemicellulose. Even though these two crop wastes contain enough cellulose to make them excellent sources of energy for ruminants, they are poor quality feeds due to low digestibility, poor palatability, low protein content and bulkiness (Umasaravanan *et al.*, 2011). Sugarcane bagasse and Khesari plant are mainly utilized as fuel. Conversion of these wastes by biotechnological approaches seems to be practical and promising alternative for transforming them into animal feed, increasing their nutritional value and thus producing a value-added product.

Some bacteria (such as *Clostridium thermocellum*) and fungi (like *Trichoderma*) are among the mostly utilized sources of cellulase enzyme which convert cellulose to glucose. Among these, fungi have been studied extensively. *Pleurotus sajor-caju* (edible fungi) obtain their energy by decomposing wood especially sapwood and heartwood of broad-leaved or rarely coniferous trees (Hossain *et al.*, 2009).

Making cellulose accessible to the enzymes is an essential factor to increase the rate of hydrolysis. Chemical pretreatment like alkaline pretreatment can increase the enzymatic effect on cellulose. This pretreatment not only removes lignin but also acts as a swelling agent, which enhances surface area of the substrate accessible for enzymatic activity (Rodhe *et al.*, 2011).

Solid state fermentation of lignocellulose materials by fungi has received attention, primarily because of low energy consumption, process simplicity, superior enzyme productivity, low capital investment, negligible liquid waste product and ease in product recovery (Reid and Seifert, 1982; Mukhopadhyey and Nandi, 1999). There is little literature on the use of fermented compound feed manufactured according to solid-state fermentation. This type of feed may have the potential to be efficient in promoting animal health and growth (Hu *et al.*, 2008). This process can greatly improve the nutritional value of cellulosic substrates like straw, bagasse etc (Hossain *et al.*, 2009).

The objective of the study is to the pretreatment affect the efficiency of solid state fermentation of khesari plant and mixed substrate (mixture of sugarcane bagasse and khesari plant) by *P. sajor-caju* (increased cellulolytic enzyme activity) for production of nutritionally enriched animal feed (increased amount of sugar and protein content).

Materials and methods

This study was done in the research laboratories of the Department of Biotechnology and Genetic Engineering, Islamic University and Institute of Food and Radiation biology, Atomic Energy Commission; Bangladesh from January 2011 to December 2011.

Biological Samples and Treatment

Two natural cellulosic agro-wastes were selected for the present study. Khesari plant alone or in combination with sugarcane bagasse were used as a substrate. Khesari plant and sugarcane bagasse were collected from local area of Navaron near Jessore district of Bangladesh. Collected cellulosic materials were first cleaned off from dirt and unwanted materials. Then, these were cut manually into tiny pieces (1.5cm), washed manually with water to clean dirt and unwanted materials, sun dried and then grounded by passing through a pulverizer (BICO Inc., USA).

The pure *P. sajor-caju* culture was used for fermentation. *P. sajor-caju* was routinely subcultured and maintained on potato dextrose agar (PDA) slants and was stored at 4°C.

Experimental design

In this study mixed substrate (mixture of sugarcane bagasse and khesari plant) are compared with khesari plant. Initially the grounded khesari plant and the mixture of sugarcane bagasse and khesari plant (50:50) were divided into three groups. One of them was lime treated, another group was alkali treated and the remained one was kept untreated. Now each of the three groups was further divided into two subgroups. One of the each subgroup was soaked overnight in distilled water (100ml/25gm) at room temperature and the other one kept unsoaked. Soaking was done to find out how this water treatment affects the final result. All the six subgroups were subjected to fermentation by *P. sajor-caju* followed by determination of reducing sugars, soluble proteins and enzyme activity.

Lime pretreatment

500g of untreated substrate were taken and then soaked with a lime solution (10g CaCO₃ in 3.75 liter distilled water). The substrates were left in soaking condition overnight. Then the lime solution was drained out by tap water (Awang *et al.*, 1994). This process was repeated until the pH of the final wash water reached 7.2). Treated substrate was then spread over aluminum foil and allowed to dry overnight at 60°C.

Alkali pretreatment

Untreated substrates were treated with sodium hydroxide (0.1 g/g of substrate) (Varga *et al.*, 2002). Substrate (50 g) was taken into a two liter conical flask containing 5 g sodium hydroxide in one liter distilled water. The flask containing substrate and sodium hydroxide solution was heated in boiling water (100°C) bath for 1hour. The flask was cooled under running tap water. The alkali treated substrate was separated from the solution by filtering with nylon cloth. Excess alkali was removed by repeated

washing with tap water until the pH was neutral. Complete removal of alkali was checked by pH paper. The nylon cloth containing substrate was squeezed to remove excess water and washed substrate spread over aluminum foil and allowed to dry overnight at 60°C.

Fermentation of substrates

P. sajor-caju was subcultured from stock PDA slant to PDA plate. After two weeks of incubation at 30°C, three pieces of mycelial growth (about 1 cm in diameter) were taken with two loops full of a hollow borer for inoculation in the fermentation medium in 100 ml conical flask in each containing 50 ml PDA broth. Then each flask was shaken in shaker for one week. Each 50 ml inoculums was then transferred to presoaked substrates (i.e. 20g substrate soaked in 110 ml H₂O for overnight) and unsoaked substrates (i.e. 20g substrate) and incubated in thermostat incubator at 30°C for 8 weeks. Controls for both presoaked and unsoaked substrate groups were maintained at same incubation conditions without *P. sajor-caju* inoculums. Both treated and untreated substrates were inoculated by fungi *P. sajor-caju* and incubated at temperature 30°C for 8 weeks as stated above.

Determination of sugar, soluble protein and enzyme activity

The substrates were investigated for the determination of their percentage of soluble protein, reducing sugar and enzyme activity. The amount of Protein was determined by the standard method developed Lowry *et al.*, 1951, reducing sugar was determined by method developed by Miller,1959 and enzyme assays were done according to Mandels and Sternberg, 1976.

Results

Effect of lime, alkali and presoaking treatment in accumulation of reducing sugar and protein

Before pretreatment, sugar (reducing sugars constituted of both cellulose and hemicellulose) and protein content were 5.12 mg/g and 0.9 mg/g respectively in mixed substrate. After alkali treatment, sugar and protein content increased

slightly (5.50 mg/g and 0.93 mg/g respectively). After lime treatment, sugar and protein content increased

up to 5.89 mg/g and 1.06 mg/g respectively.




Fig. 1. A. Reducing sugar accumulation pattern in differently treated khesari plant during 8 weeks of fermentation. B. Reducing sugar accumulation pattern in differently treated mixed substrate (khesari plant + sugarcane bagasse) during 8 weeks of fermentation. C. Soluble protein accumulation pattern in differently treated khesari plant during 8 weeks of fermentation. D. Soluble protein accumulation pattern in differently treated mixed substrate (khesari plant+ sugarcane bagasse) during 8 weeks of fermentation (RUS=Raw unsoaked, RS=Raw soaked, AUS=Alkali treated unsoaked, AS=Alkali treated soaked, LUS=Lime treated Unsoaked and LS=Lime treated soaked mixed substrate).

When, khesari plant was fermented, the highest amount of reducing sugar was found at 6th week of fermentation in presoaked lime treated substrate (17.05 mg/g) and in the following weeks, it was decreased. In lime treated unsoaked khesari plant, the highest amount of sugar was found at 6th week of fermentation (13.01 mg/g). In the case of lime treated presoaked alkali, the amount of the reducing sugar was found at 5th week of fermentation 12.26 mg/g while in case of unsoaked alkali treated substrate the

amount of reducing sugar was found 12.68 mg/g at 6th week of fermentation, and for presoaked untreated it was 11.74 mg/g. In addition, unsoaked untreated substrate had 10.87 mg/g of reducing sugar at 6th week of fermentation (Figure 1A). Similar pattern can be observed when the substrate was mixed (khesari plant + sugarcane bagasse), the highest amount of reducing sugar was found at 6th week of fermentation in presoaked lime treated substrate (22.15 mg/g) and gradually decreased in unsoaked lime treated (17.84

mg/g), presoaked alkali treated (15.91 mg/g), unsoaked alkali treated (15.16 mg/g), presoaked untreated (raw) (14.83 mg/g) and unsoaked untreated (13.06 mg/g) substrates respectively (Figure 1B).

In case of khesari plant, the highest amount of soluble protein was found 16.42 mg/g at 6th week of fermentation in presoaked lime treated substrate and gradually it was decreased in presoaked alkali treated (13.92 mg/g), unsoaked alkali treated (11.81 mg/g), presoaked untreated (raw) (11.68 mg/g), unsoaked untreated (10.73 mg/g) substrates respectively and the highest amount of soluble protein was reported in

unsoaked lime treated at 6th week (13.83 mg/g) (Figure 1C).

Therefore, the same pattern has been observed when the substrate was mixed (khesari plant + sugarcane bagasse), the highest amount of soluble protein was found at 6th week of fermentation in presoaked lime treated substrate (22.80 mg/g) and gradually decreased in unsoaked lime treated (18.87 mg/g), presoaked alkali treated (17.35 mg/g), unsoaked alkali treated (16.23 mg/g), presoaked untreated (raw) (15.99 mg/g) and unsoaked untreated (15.96 mg/g) substrates respectively (Figure 1D).




Fig. 2. A. Activity of cellulase of differently treated khesari plant during 8 weeks of fermentation. B. Activity of cellulase of differently treated mixed substrate (khesari plant+ sugarcane bagasse) during 8 weeks of fermentation. C. Activity of CMCase of differently treated khesari plant during 8 weeks of fermentation. D. Activity of CMCase of differently treated mixed substrate (khesari plant+ sugarcane bagasse) during 8 weeks of fermentation. E. Activity of avicelase of differently treated khesari plant during 8 weeks of fermentation. F. Activity of avicelase of differently treated mixed substrate (khesari plant+ sugarcane bagasse) during 8 weeks of fermentation (RUS=Raw unsoaked, RS=Raw soaked, AUS=Alkali treated unsoaked, AS=Alkali treated soaked, LUS=Lime treated unsoaked and LS=Lime treated soaked mixed substrate).

Activity of crude cellulolytic enzymes in the fermented substrates

Activity of cellobiase: Cellobiase activity was found to increase swiftly from 1st to 5th week of incubation and then tend to decrease. Cellobiase activity in presoaked lime treated khesari plant was found highest (2.10 units/ml), and a gradual decrease was observed in other substrates. The activity of unsoaked lime treated, soaked alkali treated, unsoaked alkali treated, soaked untreated, and raw unsoaked khesari plant were 1.96 units/ml, 1.58 units/ml, 1.49 units/ml, 1.36 units/ml and 1.12 units/ml respectively.(Figure 2A).

In case of mixed substrate (khesari plant + sugarcane bagasse), the activity was increased in a very similar pattern. Highest cellobiase activity was observed in 5th week, was found in presoaked lime treated mixed substrate (2.22 units/ml) and gradually decreased. The enzymatic activity for unsoaked lime treated on mixed substrate was found to be 1.98 units/ml, while soaked alkali treated, unsoaked alkali treated, soaked untreated substrate activity was found to be 1.68 units/ml, 1.61 units/ml, 1.54 units/ml respectively and for untreated unsoaked the activity was 1.29 units/ml.(Figure 2B).

Activity of Carboxymethylcellulase (CMCase)

Activity of CMCase also increased gradually from 1st to 5th week and then decreased slightly. In case of khesari plant substrate, CMCase activity in presoaked lime treated was found highest, 1.30 units/ml and a gradual decrease was recorded in other substrates. In case of unsoaked lime treated, soaked alkali treated, unsoaked alkali treated, soaked untreated khesari plant the values were 1.14 units/ml, 1.04 units/ml, 0.97 units/ml, 0.94 units/ml and the activity of raw unsoaked was 0.89 units/ml (Figure 2C). In addition, the activity of mixed substrate (khesari plant + sugarcane bagasse) was increased in a very similar pattern. Highest CMCase activity was at 5th week mostly. Best CMCase activity was found in presoaked lime treated mixed substrate, 1.41 units/ml and a gradual decline was found in other substrates. Unsoaked lime treated, soaked alkali treated, unsoaked alkali treated, soaked untreated mixed

substrate presented 1.21 units/ml, 1.10 units/ml, 0.99 units/ml, 0.98 units/ml CMCase activity for untreated unsoaked mixed substrate was 0.94 units/ml (Figure 2D).

Activity of Avicelase

Activity of avicelase also rose slowly from 1st week to the 5th week of solid-state fermentation of differently pretreated cellulosic agro-wastes and then decreased slightly. The avicelase activity was found to shoot up drastically up to 5th week of incubation then decreased slightly by 8th week when khesari plant was used as substrate for solid-state fermentation. All the unsoaked and presoaked differently treated substrates followed similar pattern but a higher activity was observed in case of soaked lime treated khesari plant. Highest avicelase activity was 1.10 units/ml, in presoaked lime treated khesari plant which was greater than others. There were some results for unsoaked lime treated khesari plant, 0.99 units/ml, soaked alkali treated, 0.91 units/ml, unsoaked alkali treated, 0.87units/ml, soaked untreated, 0.82 units/ml and for raw unsoaked substrate, the activity was 0.75 units/ml (Figure 2E).

In case of mixed substrate (khesari plant+ sugarcane bagasse), the activity was found in a very similar pattern. Highest avicelase activity was at 5th week mostly during the 8 weeks of fermentation. The best avicelase activity was found in presoaked lime treated mixed substrate, 1.37 units/ml and there was a gradual decrease in other substrates. For unsoaked lime treated, soaked alkali treated, unsoaked alkali treated, soaked untreated mixed substrate those were 1.11 units/ml, 0.98 units/ml, 0.92 units/ml, 0.90 units/ml respectively. Finally, the activity for untreated unsoaked was 0.84 units/ml (Figure 2F).

Discussion

The increase in reducing sugar and soluble protein content and the activities of the cellulolytic enzymes namely cellobiase, CMCase and avicelase were determined for a period of eight weeks. Initially, the raw substrates contained small amount of reducing sugars and soluble proteins. At different stages of

solid state fermentation, reducing sugar and protein content of each of the treated substrate gradually increased with the duration of fermentation by *P. sajor-caju*. The solid state fermentation was carried out for 8 weeks. It was observed that the reducing sugar content increase rapidly up to 5th week in all substrates and after that tend to slow down. It could be due to the fact that the rate and extent of biomass hydrolysis are inextricably linked to biomass structural characteristics (Zhu, 2005).

In the present study, reducing sugar level concentration has decreased at the end of the fermentation process. It was found that the excess fungal growth lead to the consumption of some of these sugars as their carbon source for metabolic activity (Qiu *et al.*, 2003).

The lime treated khesari plant and mixed substrate (khesari plant + sugarcane bagasse) contained higher amount of reducing sugar than untreated and alkali treated substrates (Figure 1A and 1B). Among them the highest amount of reducing sugar was found in lime treated presoaked mixed substrate. Thus, it can be said that lime treatment is responsible for structural modification of the substrates leading to their increased digestibility which in turn increases the reducing sugar content. Pretreatment may reduce the indigenous microflora which could positively influence solid state fermentation and fungal growth (Tengerdy and Szakacs, 2003). Rani *et al.* (2006) reported that pretreatment of substrate increased the yields of cellulase upto 33%. The presence of lignin in cellulosic substrates hinders the saccharification into monomeric sugars. Alkali treatment dissolves the lignin by breaking the linkages (Lee, 1997). Sodium hydroxide is a typical alkali used in alkaline pretreatment as used in this study (Varga *et al.*, 2002). It has been established by Valchev *et al.*, (2009) that the enzyme treatment on raw material as first stage treatment can be inefficient, probably the impeded access of the enzyme molecules to the amorphous sections of the carbohydrate chains.

The pattern for soluble proteins in all the treated substrates were found very similar as the reducing sugars (Figure 1C and 1D). Both presoaked and lime treated khesari plant and mixed substrate (khesari plant + sugarcane bagasse) contained higher amounts of soluble proteins compared to alkali treated, untreated and unsoaked substrates. Similar results were also obtained in case of enzyme activity. da Silva *et al.*, (2005) showed a very low CMCase activities when substrate grown on liquid medium. However, when fungus was grown by solid state fermentation this enzyme production increased. The findings in this paper justify the increase of CMCase activity at 5th week of fermentation. This may be due solid state fermentation provides the fungus with an environment closer to its natural habitat (wood and decayed organic matter), which stimulates this strain to produce more cellulolytic enzymes. In this study best CMCase activity was found in presoaked lime treated mixed substrate, 1.41units/ml and a gradual decrease in other substrates after 5th week. Similar pattern were obtained for the maximum cellulases productivity by *P. ostreatus* and *P. sajor-caju* cultivated on leaf biomass and pseudo-stem biomass of banana (Reddy *et al.*, 2003). Alkali treatment might be another cause of higher enzymatic activity because alkaline condition might increase accessibility of the enzyme to the substrate for enzymatic attack by removal of lignin seal (Ahmed *et al.*, 2012). Boonmee (2009) showed some fungal isolates produced either CMCase or cellobiase, when other produces both enzymes at the same level. Our study showed different level of CMCase or cellobiase activity (1.41 units/ml and 2.22 units/ml respectively). These extracellular enzymes produced by the *P. sajor-caju* could degrade the cell wall components of the substrates (Naraian *et al.*, 2010).

Conclusion

Finally, it can be said that, khesari plant and mixed substrate (khesari plant + sugarcane bagasse) can be upgraded to enriched animal feed by chemical pretreatments especially lime treatment and solid-state fermentation using the edible fungus, *P. sajor-*

caju, as the fermented products contained high amount of reducing sugar and soluble protein.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgements

We thank staffs and kindness and financial support of the Department of Biotechnology and Genetic Engineering, Islamic University, Bangladesh.

References

Ahmed FM, Rahman SR, Gomes DJ. 2012. Saccharification of Sugarcane Bagasse by Enzymatic Treatment for bioethanol production. *Malaysian Journal of Microbiology* **8**, 97-103.

Awang MR, Husin WBW, Osman T, Mahmud MS, Zainal N. 1994. Evaluation palm empty fruit bunch and its fermented products as for ruminant animal by nutritional values characterization and in-vitro dry matter digestibility. *Proceedings of the Seminar at Malaysian Institute for Nuclear Technology*. 7-9 Nov

Hossain S, Khalil MI, Alam MK, Khan MA, Alam N. 2009. Upgrading of Animal Feed by Solid State Fermentation by *Pleurotus sajor-caju*. *European Journal of Applied Sciences* **1** (4), 53-58.

Hu J, Lu W, Wang C, Zhu R, Qiao J. 2008. Characteristics of solid-state fermented feed and its effects on performance and nutrient digestibility in growing-finishing pigs. *Asian-Australasian Journal of Animal Science* **21**, 1635-1641.

Lee J. 1997. Biological conversion of lignocellulosic biomass to ethanol. *Journal of Biotechnology* **56**, 1-24.

Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951. Protein measurement with the Folin-phenol reagent. *Journal of Biological Chemistry* **193**, 265-275.

Mandels M, Sternberg D. 1976. Recent advances in cellulase technology. *Journal of Fermentation Technology* **54**, 267-286.

Miller GL. 1959. Use of dinitrosalicylic acid for determination of reducing sugar. *Annual Biochemistry* **31**, 426-428.

Mukhopadhyey S, Nandi B. 1999. Optimization of cellulose production by *Trichoderma reesei* ATCC 26921 using a simplified medium on water hyacinth biomass. *Journal of Scientific and Industrial Research* **58**, 107-111.

Naraian R, Singh D, Verma A, Garg SK. 2010. Studies on in vitro degradability of mixed crude enzyme extracts produced from *Pleurotus* spp. *Journal of Environmental Biology* **31**(6), 945-951.

Qiu L, Zhao M, Li F, Qi W, Zhang W, Yue X, Cui J. 2003. Changes in biological activity during artificial fermentation of flue-cured tobacco. *Tob Sci* **46**, 24-27.

Reddy GV, Babu PR, Komaraiah P, Roy KRRM, Kothari IL. 2003. Utilization of banana waste for the production of lignolytic and cellulolytic enzymes by solid substrate fermentation using two *Pleurotus* species (*P. ostreatus* and *P. sajor-caju*). *Process Biochemistry* **38**, 1457-1462.

Reid ID, Seifert KA. 1982. Effect of an atmosphere of oxygen on growth respiration and lignin degradation by white rot fungi. *Canadian Journal of Botany* **60**, 252-260.

Rani SR, Sukumaran RK, Pillai A, Prema P, Szakacs G, Pandey A. 2006. Solid state fermentation of lignocellulosic substrates for cellulase production by *Trichoderma reesei* NRRL 11460. *Indian Journal of Biotechnology* **5**, 332-336.

Rodhe AV, Sateesh L, Sridevi J, Venkateswarlu B, Rao LV. 2011. Enzymatic hydrolysis of sorghum straw using native cellulose produced by *T. reesei* NCIM 992 under solid state fermentation using rice straw. *3 Biotech* **1**, 207–215.

Silva R, Lago ES, Merheb CW, Macchione MM, Park YK, Gomes E. 2005. Production of xylanase and cmcase on solid state fermentation in different residues by *Thermoascus aurantiacus* miehe. *Brazilian Journal of Microbiology* **36**, 235-241.

Tengerdy RP, Szakacs G. 2003. Bioconversion of lignocellulose in solid substrate fermentation. *Biochemical Engineering Journal* **13**, 169–179.

Tripathi MK, Mishra AS, Misra AK, Vaithyanathan S, Prasad R, Jakhmola RC. 2008. Selection of white-rot basidiomycetes for bioconversion of mustard (*Brassica compestris*) straw under solid-state fermentation into energy substrate for rumen micro-organism. *Letter in Applied Microbiology* **46**, 364–370.

Umasaravanan D, Jayapriya J, Rajendran RB. 2011. Comparison of lignocellulose biodegradation in solid state fermentation of sugarcane bagasse and rice straw by *Aspergillus tamari*. *Ceylon Journal of Science (Biological Science)* **40 (1)**, 65-68.

Valchev I, Nenkova S, Tsekova PP, Lasheva V. 2009. Use of enzymes in hydrolysis of maize stalks. *Bioresources* **4**, 285–291.

Varga E, Szengyel Z, Reczey K. 2002. Chemical pretreatment of corn stover for enhancing enzymatic hydrolysis. *Applied Biochemistry and Biotechnology* **98**, 73–87.

Zhu L. 2005. Fundamental study of structural features affecting enzymatic hydrolysis of lignocellulosic biomass. *Electronic Thesis*, A&M University, Texas.