



## Tolerance potential of different species of *Aspergillus* as bioremediation tool - comparative analysis

Shazia Iram\*, Kanwal Waqar, Naila Shuja, Kausar Perveen, Ijaz Akhtar, Iftikhar Ahmad

<sup>1</sup>Department of Environmental Sciences, Fatima Jinnah Women University, Rawalpindi, Pakistan

<sup>2</sup>National Agricultural Research Center, Islamabad, Pakistan

Article published on April 25, 2013

**Key words:** Heavy metals, Tolerance potential, *Aspergillus*, MIC.

### Abstract

Due to increase of industrialization in public and private sectors along with urbanization, are reflected in varying degree of pollution in the different compartment (water, soil and air) of environment. The present study has been carried out to understand the tolerance potential of different species of *Aspergillus* (*Aspergillus niger*, *Aspergillus flavus*, *Aspergillus fumigatus*) isolated from agriculture soil of Kasur irrigated with water contaminated with sewage and industrial effluent. The degree of tolerance of fungi was measured by minimum inhibitory concentration in the presence of different concentrations of metals (Cr and Pb) and compared to control sample. Results are shown the variation in the tolerance level of different isolates of *Aspergillus*. Few isolates are tolerant, moderately tolerant and some are sensitive.

\*Corresponding Author: Shazia Iram ✉ [iram.shazia@gmail.com](mailto:iram.shazia@gmail.com)

## Introduction

Contamination of sediments and natural aquatic receptors with different pollutant is a major environmental problem all over the world (Baldrian and Gabriel, 2002; Gavrilesca, 2004; Malik, 2004). Industrialization and urbanization especially in developing countries have led to the accumulation of heavy metals and petroleum hydrocarbons in the environment (Adveniyi, 1996; Bamgbose and Sibanjo, 1998; Byomi et al., 1999; Manay et al., 1999; Ngodigha et al., 1999; Yamasoet et al., 2000; Adedniyi and Folabi, 2002). Metals that are released into the environment tend to persist indefinitely, accumulating in to living tissues and posing a serious threat to the environment and public health. These harmful substances accumulate in crops via food chain; find their way into our bodies, where they can cause a variety of illness.

They are cytogenic, mutagenic, and carcinogenic in nature and are posing threats to the urban population, which rely on vegetables and foliage crops grown in pre-urban lands (Kumar et al., 1991; Fargosova, 1994). Introduction of heavy metal in environment generally induces morphological and physiological changes in microbial communities (Vadkertiova and Slavikova, 2006), hence exerting a selective pressure on the microbiota (Verma et al., 2001). These contaminated sites are source of metal resistance microorganisms (Gadd, 1993).

Among microorganisms fungi are very important microorganism; it can tolerate heavy metals to a limit and can also help to remove heavy metals from contaminated soil. Fungi and yeast biomasses are known to tolerate heavy metals (Khan, 2001, Baldrian, 2003; Gavrilesca, 2004). They are a versatile group, as they can adapt and grow under various extreme conditions of pH, temperature and nutrient availability, as well as high metal concentrations (Anand et al., 2006). They offer the advantage of having cell wall material which shows excellent metal-binding properties (Gupta et al., 2000). They are used for bioremediation because of

their mycelia nature and ability to accumulate metals of all families.

Many fungal species such as *Rhizopus arrhizus*, *Penicillium spinulum* and *Aspergillus niger* have been extensively studied for heavy metals biosorption and the process mechanism seems to be dependent upon species (Zhou and Kiff, 1991; Hafez et al., 1997 ; Kapoor and Viraraghavan, 1997). The accumulation of heavy metals in agricultural soils is of increasing concern due to food safety issues and potential health risks as well as its determinate effects on soil ecosystem. Farmers in big cities where water from natural surface rain is not available use sewerage water and water of natural drains for crop production, due their being less expensive.

In most parts of Pakistan, untreated city effluent is utilized for growing vegetables around large urban settlements such as Kasur. Farmers use it as a source of irrigation water and plant nutrients. However, its continuous use may have serious environmental implications, since it also contains heavy metals. Use of untreated city effluent for irrigation without risk assessment and management could be a serious hazard, impacting soil and crop quality and ultimately human health. Heavy metals often lead to the degradation of soil health and contamination of food chain mainly through the vegetables grown on such soil( Jackson and Alloway 1992; Rattan et al., 2002). In Pakistan, heavy metals contaminated land is becoming an environmental and economic issue. Combination of inadequately plan effluent disposal techniques and a rapidly growing population has lead to gradual accumulation of heavy metals in the soils and water of Pakistan. According to EPA, the industrial effluents being dumped into the land, by electroplating, tanner industries and textile mills and contains toxic metals such as Cr, Ni, As, Hg, Cu, and Pb, which have contaminated the soil and the biota residing in and around soils in Pakistan (EPA, 1990). Over the years there have been increasing concerns over the rising level of heavy metal contamination, as they tend to persist in the environment,

accumulating in the biota, thereby entering the food chain causing deleterious harm to higher animals and humans (Zhang et al., 1998). Harmful substances accumulate in crops and via food chain, find their way into our bodies, where they can cause a variety of illness.

The aim of present study was to check the tolerance potential of different species of *Aspergillus* was investigated against heavy metals. For the present study soil samples were collected from peri-urban agricultural areas of Kasur which is irrigated by the contaminated industrial water and containing heavy metals and toxic chemicals. Fungi were isolated from the contaminated soil and studied for their tolerance analysis.

## Materials and methods

### *Sampling and Sampling site*

The main purpose of the present study was to evaluate the tolerance potential of different isolated fungal strain towards heavy metals (Pb, Cr). For this purpose, soil samples were collected from peri-urban agricultural areas of Kasur (Figure 1). The agriculture land of kasur is contaminated by sewage and industrial effluents from the nearby industries and contains heavy metals and toxic chemicals. From the contaminated soil fungi were isolated and preserved for further detail investigation of heavy metal tolerance.

### *Sterilization of Apparatus*

Petri plates, media bottles, distilled water, McCartney bottles and syringes were sterilized in autoclave. For sterilization purpose all apparatus were autoclaved for 40 minutes at 121°C. After autoclaving all sterilized material dried in oven at 95°C.

### *Media Preparation and Fungal Isolation*

Potato Dextrose Agar (PDA) media was used for the isolation of fungi. For the preparation of PDA, potatoes (200g) were peeled, sliced and boiled, and then sieved through a clean muslin cloth to get a

broth in which agar (7.5g) and dextrose sugar (7.5g) was added. The media was then autoclaved for 30 minutes at 121°C (Martin, 1995). Fungi were isolated on Potato Dextrose Agar (PDA) by soil dilution method (Razak et al., 1999).

### *Preparation of Plates*

Fungi were isolated on Potato Dextrose Agar (PDA) by soil dilution method (Razak et al., 1999). Poured the media in Petri-dishes and allowed to solidify for 24 hour. To suppress the bacterial growth, 30 mg/lit of streptomycin was added in the medium (Martin, 1950). Once the agar was solidified, and then put plates in an inverted position for 24 hours at room temperature (Martin, 1995).

### *Identification of Fungi*

The fungal cultures were identified on the basis of macroscopic (colonial morphology, color, texture, shape, diameter and appearance of colony) and microscopic (septation in mycelium, presence of specific reproductive structures, shape and structure of conidia, and presence of sterile mycelium) characteristics (Table, 1) (Zafar et al., 2006). Pure cultures of fungi isolates were identified with the help of literature (Domsch et al., 1980; Barnett and Hunter 1999).

### *Heavy Metal Tolerance Test Experiment*

For the evaluation of tolerance potential among isolated fungal strains, Potato Dextrose Agar (PDA) medium was prepared and amended with various concentrations (0, 200, 400, 600, 800 and 1000 mg/l) of heavy metal {Cr(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O and Pb(NO<sub>3</sub>)<sub>2</sub>} and 5.6 pH was maintained by adding 5 molar solution of NaOH. Media was autoclaved for 20 minutes at 121°C and poured in Petri plates. The plates were incubated at 29°C for 7 days. The growth of fungi was monitored from the point of inoculation or centre of the colony. Tolerance is measured by observing minimum inhibitory concentration (MIC) and tolerance index.

### *Heavy metals analysis of soil*

Each soil sample (1g) was taken in the conical flask (50ml), added 10ml of HNO<sub>3</sub>:HClO<sub>4</sub> (1:2) solution (50ml) and heated for half an hour. Solutions were filtered through Whatman 1 filter paper and volume was made to 50ml by adding distilled water. Soil samples were digested in triplicates and analyzed for Zn, Cd, Cr, Cu, Ni and Pb. The blank was prepared for quality assurance of samples. The blank sample contained 10ml of HNO<sub>3</sub>:HClO<sub>4</sub> (1:2) solution and heated for half an hour and volume was made 50ml by adding distilled water. For the determination of heavy metals the atomic absorption spectrophotometer was powered on and warmed up for 30 minutes. After the heating of cathode lamp, the air acetylene flame was ignited and instrument was calibrated or standardized with different working standards (Vanloon and Lichwa, 1973)

*Statistical Analysis*

The experiments were set up with three replicates. Analysis of variance was performed by using statistical software (SPSS 17) to compare resistance to metal among individual isolates.

**Results and discussion**

Heavy metals released to the environment have been increasing continuously as a result of the industrial activities and technological development and poses significant threat. Application of domestic and industrial effluents to nearby agricultural fields is a common practice of irrigation and wastewater disposal in Pakistan. When wastewater is applied to agricultural fields, heavy metals enter the soil and get fixed into the soil components. Thus continuous application of wastewater tends to accumulate large quantities of heavy metals in soil, which persists there for an indefinite period to have long lasting effects in the soil.

With increasing environmental awareness and legal constraints being imposed on discharge of effluents, different type of techniques presently in existence for removal of heavy metals from contaminated waters including reverse osmosis, electro dialysis, ion-

exchange, chemical precipitation, phytoremediation, etc. However, all these methods have disadvantages like incomplete metal removal, high reagent and energy requirements generation of toxic sludge or other waste products that require careful disposal (Ahalya et al., 2003). Recently research interest has diverted towards use of biomaterials like microbial biomass has emerged as an alternative technique.

**Table 1.** List of Fungal isolates.

Isolated fungus	Sample code
Aspergillus fumigatus	K3, K4, K12, K18, K26, K27, K28, K29, K30
Aspergillus niger	K 14, K 16, K 22, K 23, K 24
Aspergillus flavus	K 5, K 7, K 17, K 33, K 34, K 36, K 35, K 37, K 41

Fungi are natural inhabitants of soil and have greater potential for remediation by virtue of their aggressive growth; greater biomass, production and extensive hyphal reach in soil. They are a versatile group, as they can adapt and grow under various extreme conditions of pH, temperature and nutrient availability, as well as high metal concentrations (Anand et al., 2006). They offer the advantage of having cell wall material which shows excellent metal-binding properties (Gupta et al., 2000). Describing the ability to grow at high metal concentrations, distinguished fungi tolerant to heavy metals. Generally, the contaminated sites are the sources of metal resistant micro-organism (Gadd, 1993).

The results of present study depicted that all tested isolates shows different tolerance behavior for different metals. Some isolates were sensitive, moderately tolerant and tolerant (table 2). Most of the isolates were sensitive and very few were tolerant. Among all tested isolates of *Aspergillus*, *A.flavus* was more tolerant specie while *A.niger* was moderately tolerant. As Baldrian and Gabriel, 2002 reported that various genera and also isolates of the

same genus did not necessarily have the same heavy metal tolerance.

It is apparent that these fungal strains exhibit considerable tolerance towards Cr and Pb and can

become dominant microorganisms in some polluted habitats. This response (i.e. resistance to Cr and Pb) of *Aspergillus flavus* can be exploited for the remediation of heavy metal contaminated soil (McGrath, 2001).

**Tables.** Tolerance index of fungal isolates against lead.

Isolated fungus	Pb (TI)	Isolated fungus	Pb TI
<i>Aspergillus fumigatus</i> (K3)	0.29	<i>Aspergillus fumigatus</i> (K26)	0.28
<i>Aspergillus fumigatus</i> (K4)	0.30	<i>Aspergillus fumigatus</i> (K27)	0.17
<i>Aspergillus flavus</i> (K 5)	0.47	<i>Aspergillus fumigatus</i> (K28)	0.23
<i>Aspergillus flavus</i> (K 7)	1.09	<i>Aspergillus fumigatus</i> (K29)	0.28
<i>Aspergillus niger</i> (K 14)	0.71	<i>Aspergillus fumigatus</i> (K30)	0.25
<i>Aspergillus niger</i> (K 16)	0.48	<i>Aspergillus flavus</i> (K 33)	0.89
<i>Aspergillus flavus</i> (K 17)	1.18	<i>Aspergillus flavus</i> (K 34)	1.26
<i>Aspergillus niger</i> (K 22)	0.62	<i>Aspergillus flavus</i> (K 36)	0.12
<i>Aspergillus niger</i> (K 23)	0.54	<i>Aspergillus flavus</i> (K 35)	0.14
<i>Aspergillus niger</i> (K 24)	0.63	<i>Aspergillus flavus</i> (K 37)	0.39
		<i>Aspergillus flavus</i> (K 41)	0.55

**Tables.** Tolerance index of fungal isolates against chromium.

Isolated fungus	Cr (TI)	Isolated fungus	Cr TI
<i>Aspergillus fumigatus</i> (K3)	0.16	<i>Aspergillus fumigatus</i> (K26)	0.40
<i>Aspergillus fumigatus</i> (K4)	0.16	<i>Aspergillus fumigatus</i> (K27)	0.19
<i>Aspergillus flavus</i> (K 5)	0	<i>Aspergillus fumigatus</i> (K28)	0.26
<i>Aspergillus flavus</i> (K 7)	0.14	<i>Aspergillus fumigatus</i> (K29)	0.44
<i>Aspergillus niger</i> (K 14)	0.47	<i>Aspergillus fumigatus</i> (K30)	1.37
<i>Aspergillus niger</i> (K 16)	0.09	<i>Aspergillus flavus</i> (K 33)	0.39
<i>Aspergillus flavus</i> (K 17)	1.09	<i>Aspergillus flavus</i> (K 34)	0.26
<i>Aspergillus niger</i> (K 22)	0.17	<i>Aspergillus flavus</i> (K 36)	0.20
<i>Aspergillus niger</i> (K 23)	0.39	<i>Aspergillus flavus</i> (K 35)	0.12
<i>Aspergillus niger</i> (K 24)	0.47	<i>Aspergillus flavus</i> (K 37)	0.33
		<i>Aspergillus flavus</i> (K 41)	0.12

*Aspergillus flavus* and *Aspergillus niger* showed tolerance against Pb but it showed sensitiveness for Chromium metal. On the other hand, *A.fumigtus* shows exactly contrasting behavior. The variation in


the metal tolerance may be due to the presence of one or more strategies of tolerance or resistance mechanisms exhibited by fungi (Ezzouhri et al., 2009; Babich and Stotzky, 1979). Ezzouhri et al.,

2009 also screened fungi (*Fusarium sp*) for their resistance to heavy metals. The results revealed that the majority of the isolates were resistant to Cr, Cu and Zn. The level of resistance depended on the

**Table 2.** Concentration of Heavy Metals in Kasur Soil (mg/kg)

	Pb	Cd	Cr
Min	42.1	2.0	54.1
Max	70.2	3.4	210.2



Fungi growth phase, which was observed from the tolerance index of the fungi with time upon exposure to heavy metals, was found to be a predictable pattern for each strain and metal concentrations. The determination of Minimum inhibitory concentration (MIC) of metals suggests that the resistance level against individual metals dependent on different fungal isolates. MIC was defined as the minimum inhibitory concentration of the heavy metal that inhibited visible growth of test fungi. MIC of Cr and Pb is determined. It ranged from 800mg/l to 1000mg/l for chromium and lead for all tested isolates (figure. 3). The growth phase was characterized by a lag, retarded, similar and enhanced rates compared to control which appear to reflect the tolerance development or adaptation of fungi in the presence of heavy metals (Hashem et al., 2000).



**Fig. 1.** Map showing sampling site.


isolate tested, as well as the site of its isolation. The statistical analysis showed the diverseness in heavy metal tolerance of different isolates (Figures. 2, 3 and 4).

Mean	50.6	2.6	99.6	44
Median	47.2	2.6	93.2	43
S.D	±10.0	±0.5	±49.4	±8
Kurtosis	0.852364	-0.95102	4.182443	2.3
Skewness	1.285644	0.437593	1.824419	0.5
	60.8	56.6	156.7	




**Fig. 2.** Tolerance index of isolates of *A.fumigatus* against (a) chromium and (b) lead.

The results revealed that growth rate of all tested isolates at lower concentration was higher but with but with passage of time and exposure of higher concentration density and growth of species were reduced leading to no growth. It was of interest to establish the effect of type of heavy metal, the metal concentration and the strain on this adaptive behavior. The same increment and reduction in growth was observed during study on filamentous fungi, which belonged to the genera *Aspergillus*, were more resistant to Cr at higher metal concentrations and suddenly the growth pattern changes (Wainwright et al., 1997).



**Fig. 3.** Tolerance index of isolates of *A.flavus* against (a) chromium and (b) lead.


**El-Morsy (2004)** studied 32 fungal species isolated from polluted water in Egypt for their resistance to metals and found that *Cunninghamella echinulata* biomass could be employed as a biosorbent of metal ions in wastewater. However, although some authors found that micro-organisms isolated from contaminated sites were more tolerant than those from natural environments (Ahalya et al., 2003).



**Fig. 4.** Tolerance index of isolates of *A.niger* against (a) chromium and (b) lead.

Our preliminary findings indicate that fungi from soil contaminated with heavy metals have tolerance potential due to their physiological adaptation and they have greater potential for remediation of metals.

*A.flavus* and *A.niger* showed higher tolerance potential than other tested fungal strain which makes them more attractive potential candidates for further investigations regarding their ability to remove metals from contaminated sites.



**Fig. 5.** Minimum inhibitory concentration of different species of *Aspergillus* against metals.

**Conclusion**

The present study concludes that soil irrigated with waste water contains tolerant fungus. The findings are *A.flavus* and *A.niger* showed higher tolerance potential than other tested fungal strain which makes them more attractive potential. The results obtained affirmed that the response of the isolates to heavy metals depended on the metal tested; its concentration in the medium and on the isolate considered and can be used further for the purposes of bioremediation.

**References**

**Adveniyi AA.** 1996. Determination of cadmium, copper, iron, lead, manganese and zinc in water leaf (talinum triangular) in dumpsite, Journal of Environment International **22**, 259-262.

**Adeniyi AA, Folabi JA.** 2002. Determination of total petroleum hydrocarbons and heavy metals in soils with in the vicinity of facilities handling refined petroleum products in logos metropolis, Journal of Environment International **28**, 79-82.

**Anand P, Isar J, Saran S, Saxena RK.** 2006. Bioaccumulation of copper by *Trichoderma viride*, Bioresoure Technology **97**, 1018-1025.

- Ahalya N, Ramachandra TV, Kanamadi RD.** 2003. Biosorption of heavy metals. Res. J. Chem. Environ 7, 71-78.
- Babich H, Stozky G.** 1985. Heavy metal toxicity to microbe mediated ecological processes: a review and potential application to regulatory policies. Environment Resource 36, 111-137.
- Bamgbose O, Sibanjo O.** 1998. Comparative assessment of three methods for chlorinated hydrocarbon determination in fish, *Mig. Journal of Science* 32, 169-174.
- Byomi AM, Hassan MM, Saleh RS, Yousaf MA, Aamer AA.** 1999. Population of waste and animals feed with some heavy metals and their implication in raw buffalo milk. In proceeding of the 5<sup>th</sup> scientific congress Egyptian soc, For cattle diseases Egypt 28, 226-230.
- Barnett HL, Hunter BB.** 1999. Illustrated genera of imperfect fungi, fourth edition. Prentice Hall Inc.
- Baldrian P, Gabriel J.** 2002. Intraspecific variability in growth response to cadmium of the wood-rotting fungus *Piptoporus betulinus*. Mycologia 66, 428-436.
- Baldrian P.** 2003. Interactions of heavy metals with white-rot fungi. Enzyme Microbiological Technology 32, 78-91.
- Baldrian P, Gabriel J.** 2002. Intraspecific variability in growth response to cadmium of the wood-rotting fungus *Piptoporus betulinus*, Mycologia 94, 428-436.
- Domsch KH, Gams W, Anderson TH.** 1980. Compendium of soil fungi. London, England: Academic Press.
- EPA.** 1990. Drafts report Govt. of Punjab, Lahore. Environment Protection Agency, Govt. of Punjab, Lahore, Pakistan, 51.
- El- Morsy EM.** 2004. *Cunninghamella echinulata* a new biosorbent of metal ions from polluted water in Egypt. Mycologia 96, 1183-1189.
- Ensink JHJ, Simmons RW, Vander Hoek W.** 2004. Wastewater use in Pakistan: the cases of Haroonabad and Faisalabad. In: Wastewater Use in Irrigated Agriculture: Confronting the Livelihood and Environmental Realities (Eds. C.A. Scott, N.I. Faruqi and L. Raschid-Sally). CAB International, Wallingford. 91-99.
- Ezzouhri L, Castro E, Moya M, Espinola F, Lairini K.** 2009. Heavy metal tolerance of filamentous fungi isolated from polluted sites in Tangier, Morocco. *African Journal of Microbiology Research* Vol. 3(2), 035-048.
- Fargosova A.** 1994. Effect of Pb, Cd, Hg, As and Cr on germination and root growth of *sinapsis alba* seeds, Journal of Environment Quality 26, 639-646.
- Gadd GM.** 1993. Interaction of fungi with toxic metals. New Phytol. 124- 24.
- Gavrilesca M.** 2004. Removal of heavy metals from the environment by biosorption. Eng. Life Sci. 4, 219-232.
- Gupta R, Ahuja P, Khan S, Saxena R.K, Mohapatra H.** 2000. Microbial biosorbents meeting challenges of heavy metal pollution in aqueous solutions, Current Sciences 78, 967-973.
- Gavrilesca M.** 2004. Removal of heavy metals by the environment Biosorption. Life Sciences 4, 219-232.
- Hafez N, Abdel-Razek AS, Hafez MB.** 1997. Accumulation of some heavy metals on *Aspergillus flavus*. J. Chem. Technol. Biotechnol 68, 19-22.

**Hashem AR.** 1995. Effect of Heavy Metal Ions on the Mycelial Growth of Some Fungi Isolated from the Soil of Al-Jubail Industrial city, Saudi Arabia, Department of Botany and Microbiology, College of Science. King Saud University.

**Jackson, Alloway.** 1992. Heavy Metal Load of Soil, Water and Vegetables in Peri-Urban Delhi, Environmental Monitoring and Assessment **120**, 1-3.

**Kumar G, Singh RP, Sushila.** 1991. Nitrate assimilation and biomass production in *Sesamum indicum* L. seedlings in a lead enriched environment, Water Air and Soil pollution **66**, 163-171.

**Khan AG.** 2001. Relationship between chromium biomagnification ratio accumulation factor and mycorrhizal in plants growing on tannery effluents polluted soils, Journal of Environment International **26**, 417-423.

**Kapoor A, Viraraghavan T.** 1997. Heavy metal biosorption sites in *Aspergillus niger*, Bioresource. Technology **61**, 221-227.

**Lone MI, Aslam R, Khan KS.** 2000. Water quality and soil contamination in some industrial areas of Pakistan. Journal of Soil sciences **18**, 1-6.

**Manay N, Perera L, Cousillus Z.** 1999. Lead contamination in Uruguay, Review of environmental contamination toxicology **159**, 25-39.

**Martin JH.** 1995. On the origin of colonies of fungi developing on soil dilution plates. Trans British Mycological Society **38**, 298-301.

**Malik A.** 2004. Metal bioremediation through growing cells. International journal of Environment **30**, 261-278.

**Massaccesi G, Romero MC, Cazau MC, Bucsinszky AM.** 2002. Cadmium removal capacities

of filamentous soil fungi isolated from industrially polluted sediments, in La Plata (Argentina). Journal of Microbiology and Biotechnology **18(4)**, 817-820.

**Malik A.** 2004. Metal bioremediation through growing cells, Environment International **30**, 261-278.

**Ngodigha EN, Olayimiko FO, Oruwari BM, Ekweozor EKE, Whhekhe SN.** 1999. Toxic effects of crude oil in organs and blood cells of West Africa dwarf goat, Nigeria Veterinary Journal **20**, 82-91.

**Razak AA, Bachman G, Farrag R.** 1999. Activities of microflora in soils of upper and lower Egypt. The African Journal of Mycology and Biotechnology **7**, 1-19.

**Vanloon JC, Lichwa J.** 1973. A study of atomic absorption determination of some important heavy metals in fertilizers and domestic sewage plant sludges. Environ Letter **4**, 1-8.

**Verma T, Srinath T, Gadpayle RU, Ramteke PW, Hans RK, Garg S.K.** 2001. Chromate tolerant bacteria isolated from tannery effluent. Bioresour. Technol **78**, 31-35.

**Vadkertiova R, Slavikova E.** 2006. Metal tolerance of yeasts isolated from water, soil and plant environments. Journal of Basic Microbiology **46**, 145-152.

**Wainwright M, Gadd GM.** 1997. Fungi and Industrial Pollution. In: Wicklow DT, Soderstrom BE (eds) The Mycota, IV, Environmental and microbial relationships. Springer-Verlag, Berlin Heidelberg, 85-97.

**Yamasoet MA, Artaxo P, Miguel A, Allen AG.** 2000. Chemical composition of aerosol particle from direct emission of vegetation fire in Amazon basin.

Water soluble spices and trace elements, Atmospheric Environment **34**, 1641-1653.

**Zhou JL, Kiff RJ, Biomass J.** 1991. The uptake of copper from aqueous solution by immobilized fungal biomass, Journal of chemical technology biotechnology **52**, 317-330.

**Zhang L, Zhao L, Yu Y, Chen C.** 1998. Removal of lead from aqueous solution by non-living N pzipous Nigricans, Water Rea **32**, 1437-1444

**Zafar S, Aqil F, Ahmad I.** 2006. Metal tolerance and biosorption potential of filamentous fungi isolated from metal contaminated agricultural soil. Bioresource Technology **98**, 2557-2561.