



## RESEARCH PAPER

## OPEN ACCESS

## Comparative study determining the impacts of forest harvesting on hydrology and water yield of broadleaved and needle leaved forests: State of knowledge and research outlook

Tanzeel Javaid Aini Farooqi\*, Umer Hayat, Muhammad Roman

*College of Forest Science, Beijing Forestry University, Beijing 100083, China*

**Key words:** Deforestation, Streamflow, Broadleaved, Needle leaved, Hydrology

<http://dx.doi.org/10.12692/ijb/16.2.507-515>

Article published on February 28, 2020

### Abstract

Water availability both at present and in the future is of primary importance for sustaining life on earth. The forest role in carbon sequestration and their impacts on water yield and the hydrological cycle is currently a burning issue within the scientific community. As a general concern, different forest types and cover may vary in their effects on water interception and discharge due to various structural and growth attributes, with implications to the water supply to downstream ecosystems. Changes in forest cover impose either positive or negative impacts on catchment hydrology depending on the direction and degree of changes. To reconcile the forest carbon sequestration and water conservation, there is a dire need to understand the mechanistic linkage between the two services. By conducting literature, we examine the impacts of forest cover change in the shape of deforestation on catchment water yield for contrasting forest types: coniferous forests and broadleaved forests. The research found that after deforestation, the percent increase in streamflow in two different forest types was 16% and 27% for broadleaved and needle-leaved forests, respectively. This research provides a scientific insight about when where and why to cut/plant the trees with a particular focus on sustainable forest management practices to get maximum and long term socio-economic benefits.

\* **Corresponding Author:** Tanzeel Javaid Aini Farooqi ✉ [tanzeelfarooqi\\_21@yahoo.com](mailto:tanzeelfarooqi_21@yahoo.com)

## Introduction

Forests covered around 30% of the land surface (Bonan, 2008) and considered as the promising role in many ways by providing a vast quantity of ecosystem services (Costanza *et al.*, 1997), like NTFP's, Carbon sequestration, clean drinking water, nutrient recycling as well as biodiversity conservation. It is evident in understanding that among all of these ecosystem services, carbon sequestration and water are the primary services linked with the provision of many other functions of the forest ecosystem. Forest on one side by sequestering carbon increase the productivity as well as considered as proposed strategy to mitigate the global warming issues (Krankina *et al.*, 1997; Ruddell *et al.*, 2007); similarly, forests are considered as provision of rainfall (van der Ent *et al.*, 2010) and clean drinking water.

Therefore, because of the escalating surge in the world's population, resulting in a higher demand for water and land resources for food and fiber products at the expense of forests. The primary cause of the drastic increase of CO<sub>2</sub> concentration in the atmosphere is due to anthropogenic disturbance along with Land use and cover changes and over-exploitation of forest resources in the shape of deforestation (Houghton, 2007; Houghton, 2012). The demand for wood and fiber products for household and industry, enhance the increase of afforestation schemes have launched for the last few decades in several parts of the world (FAO, 2010; Payn *et al.*, 2015). For example: from 1990 to 2015 the global forest cover falling from 31.85% to 30.85%, but on same time an increase from 4.06% to 6.95% of total afforested area. This upsurge was most speedy in the temperate zone on the regional level of East Asia than by Europe, North America, and Southern and Southeast Asia (Payn *et al.*, 2015). These expansions accompanied by a parallel increase in concerns regarding water losses (Andréassian, 2004; Jackson *et al.*, 2005) in shape of annual streamflow reduction (Trabucco *et al.*, 2008) consequently, the decline in water availability to downstream users (van Dijk and Keenan, 2007). On the other hand, afforestation schemes triggered a reduction of up to 52%, among


which 13% of the complete drying of streams on a global level (Jackson *et al.*, 2005). Especially the drylands areas, those might be more vulnerable to climate extremes due to its high ecohydrological sensitivity (Matyas and Sun, 2014) accompanied by an increase in streamflow following forest cutting (Bosch and Hewlett, 1982; Jones and Post, 2004).

The world's forest-growing stock is composed of two-thirds of broadleaf trees and one-third of needle leaf trees (FAO, 2010). There are many studies have been reported the interrelationship between deforestation and their influences on water yield very globally depending on different forest types (Hornbeck *et al.*, 1993; Vildan and Michael, 1996; Vazken Andreassian, 2003; Komatsu *et al.*, 2011; Troendle *et al.*, 2001; Pike and Rob, 2003; Adams and Flower, 2006).

The deforestation on one side enhances stream water level, but on the other hand has many adverse impacts on whole ecosystem, e.g., land degradation, biodiversity disturbances, flood, disturb the chemistry of carbon water and nitrogen cycle, as well as other environmental problems which may collectively intensify the global warming. A piece of explicit knowledge on how forest types would shape the tradeoff between forest productivity and ecosystem water loss is a pre-requisite for assisting regional forestry planning and forest management in the context of carbon sequestration and water conservation. In this fight between forest carbon and water, the negative influences of forest on streamflow might be to control the proportion of forest cover at the catchment scale, which has the potential to modify the streamflow regime (Zhang *et al.*, 2012). For this purpose, it is dire need to understand better the affiliation between runoff concerning forest cover proportion (Brown *et al.*, 2013).

This research highlighted the research gaps of deforestation schemes, and their prone and corns for society and how to tackle these issues to gain and maintain maximum availability of forest ecosystem services on sustainable basis (Fig. 1) as carbon-water interrelation entirely depends upon many factors, the most dominant factors and forgiven climate, forest

types, and structure, age as well as soil. The objective of the study is to simulate the hydrological effects of different deforestation scenarios of broadleaved and needle-leaved forests on a global scale.



**Fig. 1.** Conceptual framework showing the core issue during deforestation scheme and their prone and corns.

## Materials and methods

### Data collection and processing

We have compiled this large data set of deforestation studies and their impacts on water yield from research articles published peer review journals; research from the World's forest, consisting total of 71 watershed sites of Needle leaved forest stand and 103 sites belongs to broadleaved forest stand with totaling of 310 observations from all over the world. We compiled this data sets from 22 peer-reviewed journals as well as reports from governmental and nongovernmental research institutes, representing many parts of the world (Supplementary info.). The forest types were classified into needle leaved and broadleaved for this study. All the information gathering about different deforestation activities on water yield was compiled, with mean annual streamflow before and after treatment.

### Testing of Significance

First we have performed the Normality test i.e Shaprio-wilk test, this test showed that the conditions of normality and homogeneity of variance were not met. Later nonparametric Kruskal–Wallis tests were applied. In each case, the dependent variable was either the proportional change in water yield following change in factors of evaluation i.e deforestation percentage. The significance test suggests that the water yield rate isn't the same in each of the two or more region ( $p < 0.001$ ).

## Results and discussion


The results of Shaprio-wilk test rejected normality at  $P < 0.0001$  as shown in Fig. 2. Similarly, the Kruskal-Wallis test showed significant increase in streamflow after deforestation in broadleaved forests of the world with  $P < 0.0001$ . The summery of test statistic is mentioned in Table 1. The results also indicated that before treatment annual streamflow of broadleaved forests was 775mm per annum whereas, after treatment an additional of 150mm of water added into streamflow with total of 925mm, leading to annual increase of 16% in the streamflow of broadleaved forests in the world as shown in Fig. 3(a). Regression analysis also demonstrate that correlation between deforestation and change in water yield is consistent, with increasing trend along with the increase of deforestation percentage with  $R^2 = 0.21$ ,  $p < 0.001$  as shown in Fig. 4(a). Broadleaved are mainly known as nutrient-rich area species, warm weather, and continuous availability of sunshine in the spring season that actually boost photosynthesis process with their extensive set of leaves. Especially in the winter season, deciduous species which are in the majority of broadleaves leaves e.g., Oak, maple, and elm, etc. which lose chloroplast cells that are the main drivers for capturing sunlight and ultimately leaves changes their color and finally fall to the ground. Movement of energy into roots, tree enters into dormancy condition. However, there are few exceptions like live Oak, eucalyptus, and the majority of the tropical forest belongs to a broadleaved group, which is evergreen species. They do not lose their leave in the dormancy period. Our results are primarily following the findings of many regional

watershed studies. The results reveals that on global level the impact of deforestation on streamflow also depend on the forest subtype e.g., in broadleaved evergreen Eucalyptus species on streamflow was positive with higher increase in streamflow reported by many previous research investigations (Watson *et al.*, 2001; Cornish, 1993; Stoneman, 1993; David *et al.*, 1994; Lane and Mackay, 2001).

**Table 1.** Representing the summary of test statistics for Pre and Post treatment water yield in broadleaved and needle leaved forest.


Treatments	n	Chi-square	DF	P-value
Pre-NL vs Post NL	188	102.33	01	<0.0001
Pre-BL vs Post BL	188	165.14	01	<0.0001

Similarly, another study deciduous broadleaf forest catchments in North America showed a significant increase in the water quantity after fifteen years of clear-felling (Lynch and Corbett, 1990; Hornbeck *et al.*, 1993). In another regional collection of research investigations of streamflow showed highly significant results with a consistent increase in water yield after harvesting of broadleaved deciduous forest stands (Bosch and Hewlett, 1982; Stednick, 1996; Fahey and Jackson, 1997). Similar results have reported by (Watson *et al.*, 2001; Cornish, 1993; Stoneman, 1993; Lane and Mackay, 2001) the same in broadleaved evergreen which is mainly in line with our research findings. The results of Shaprio-wilk test rejected normality at  $P < 0.0001$  as shown in Fig. 2. Similarly, the Kruskal-Wallis test showed significant increase in streamflow after deforestation in needle leaved forests of the world with  $P < 0.0001$ . The summary of test statistic is mentioned in Table.1. The results clearly reveal that before treatment annual streamflow of needle leaved forests was 410mm per annum whereas, after treatment an additional of 111mm of water added into streamflow with total of 521mm with annual increase of 27% in the streamflow of needle leaved forests in the world as shown in Fig. 3(b). Regression analysis also demonstrate that correlation between deforestation of needle leaved forest and change in water yield is also consistent, showing increasing trend along with the increase of percentage cut area of forest with  $R^2 = 0.23$ ,  $p < 0.001$  as shown in Fig. 4(b).



**Fig. 2.** Shaprio-wilk test rejected normality at  $P < 0.0001$  in pre and post deforestation of (a) Broadleaved and (b) Needle leaved forests in the global dataset.

\*Pre-BL: Pretreatment streamflow of Broadleaved forest, Post-BL: Post treatment Increased in streamflow of Broadleaved forest, Pre-NL: Pretreatment streamflow of needle leaved forest, Post-NL: Post treatment Increased in streamflow of needle leaved forest.




**Fig. 3.** Illustrating the Post treatment increase in water yield (mm) in (a) Broadleaved and (b) Needle leaved forests on global scale catchment studies.

\*Pre-BL: Pretreatment streamflow of Broadleaved forest, Post-BL: Post treatment Increased in streamflow of Broadleaved forest, Pre-NL: Pretreatment streamflow of needle leaved forest, Post-NL: Post treatment Increased in streamflow of needle leaved forest.

Needle leaved or conifers are consisting of long thin leaves. The majority of needle-leaved are evergreen species, so they remain on tree whole year, and the replacement process is slow and continuous, there are few exceptions of Tamarack and larch are needle-leaved deciduous tree species.

Their leaves are usually smaller, rough structure, tighter needles having spatiality of wind and pest resistant as well as waterproofing than broadleaved and deciduous species. They can withstand harsh weather of spring season with poor nutrient soils. In many of previous big data sets, it indicated that needle-leaved forests, consumption, and utilization of water are relatively higher and deforestation treatment caused a significant increase in water yield (Anderson *et al.*, 1976; Bosch and Hewlett, 1982; John, 1996; Scott *et al.*, 2000; Adams and Flower, 2006; Komatsu *et al.*, 2011). Similarly, the clear-felling in large watersheds (>2000ha) showed a significant upsurge in water yield with around 25% cut resulted in a 52% increase in water yield of that particular area (Burton, 1997).



**Fig. 4.** Showing the consistent increase in water yield (mm) after deforestation in (a) Broadleaved and (b) Needle leaved forests of dataset.

\*Pre-BL: Pretreatment streamflow of Broadleaved forest, Post-BL: Post treatment Increased in streamflow of Broadleaved forest, Pre-NL: Pretreatment streamflow of needle leaved forest, Post-NL: Post treatment Increased in streamflow of needle leaved forest.

These results are unique of its type because, in previous studies, it was only compared deforestation cause an increase or decrease in streamflow between both forest types after treatment without knowing, that before and after treatment, what actual increase in streamflow is and in various forest types?

In this study, we have also come to know that in broadleaved forests, the majority of the area seems sufficient water area, whereas needle-leaved forests lie in water-deficient zones, as shown in Fig.3(a & b). The reason for higher water consumption of majority of the needle-leaved forest is due to their evergreen in

nature. They consume water whole year, whereas in broadleaved because of the presence of a majority of deciduous species might be the prominent reason for their efficient utilization of water during the dormant season of the year. There is a common belief that broadleaved having more runoff because of less annual evapotranspiration than needle-leaved forests. This is mostly known belief that, in broadleaved deciduous forest with higher winter precipitation (e.g., the United States) (Bosch and Hewlett, 1982; Landsberg and Gower, 1997; Komatsu *et al.*, 2011), broadleaf forests are usually evergreen like in New Zealand (Landsberg and Gower, 1997) and their winter precipitation is typically low in Japan (Komatsu *et al.*, 2010). In our dataset consisting of both broadleaved deciduous in the majority as well as evergreen species of the forest. Therefore, we can say that this might be one of the reasons for having a more significant percent change in annual water yield than needle-leaved harvesting. Experiments in small and large areas represent broadleaved and needle-leaved forests trees. In this range, the relationship between pre-harvest water yield and water yield increase after harvesting varied greatly in both of the forest types of the world. However, we have also suggested that there might be many other factors involved in the variation of streamflow response to deforestation of different forest types e.g., elevation, slope, area, soil, climate, are the prime factors influencing the water yield of a particular forest stand.

## Conclusion

This study showed that water yield response to cover changes is unpredictable among conifers and deciduous. However, some general conclusions can be justified. Deforestation of conifers showed a more considerable annual increase in water yield than deciduous forest. However, this decrease is dependent on its associated factors, especially climate, topography, growth stages, and soil. In most cases, the reduction of cover less than 20% did not project a clear picture of the increase of annual water yield. The catchment studies results are complicated to express very precisely due to the variation in experimental conditions as well as different ways of conducting research trails.

However, the available data representing valuable information about the role of forest structure on hydrology especially in high and low rainfall area, that will be helpful for decision-makers to take a practical step for modifying the current Carbon sequestration policy.

#### Acknowledgement

TJA Farooqi acknowledges the financial support of the China Scholarship Council and Ministry of Science and Technology of China (Grant No. 2016YFC0502104).

#### Authors' contributions

TJA Farooqi conceived the idea and designed the conceptual framework, conducted literature search and wrote the manuscript. U Hayat assisted and contributed in data analysis. R Muhammad assisted in building conceptual framework. All authors read and approved the final manuscript.

#### Abbreviation

FAO- Food and Agricultural Organization, NTFP's- Non-timber forest products, USA-United State of America

#### References

- Adams KN, Fowler AM.** 2006. Improving empirical relationships for predicting the effect of vegetation change on annual water yield. *Journal of hydrology* **321**, 90-115.
- Andréassian V.** 2004. Waters and forests: from historical controversy to scientific debate. *Journal of hydrology* **291**, 1-27.
- Bonan GB.** 2008. Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science* **320**, 1444-1449.
- Bosch JM, Hewlett JD.** 1982. A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration. *Journal of hydrology* **55**, 3-23.
- Brown AE, Western AW, McMahon TA, Zhang L.** 2013. Impact of forest cover changes on annual streamflow and flow duration curves. *Journal of Hydrology* **483**, 39-50.
- Burton TA.** 1997. Effects of basin-scale timber harvest on water yield and peak streamflow 1. *JAWRA Journal of the American Water Resources Association* **33**, 1187-1196.
- Cornish PM.** 1993. The effects of logging and forest regeneration on water yields in a moist eucalypt forest in New South Wales, Australia. *Journal of Hydrology* **150**, 301-322.
- Costanza R, d'Arge R, de Groot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem S, O'Neil RV, Paruelo J, Raskin RG, Sutton P, van den Belt M.** 1997. The value of the world's ecosystem services and natural capital. *Nature* **387**, 253-260.
- David JS, Henriques MO, David TS, Tomé J, Ledger DC.** 1994. Clearcutting effects on streamflow in coppiced *Eucalyptus globulus* stands in Portugal. *Journal of Hydrology* **162**, 143-154.
- Ellison D, Morris CE, Locatelli B, Sheil D, Cohen J, Murdiyarto D, Gutierrez V, Van Noordwijk M, Creed IF, Pokorny J, Gaveau D.** 2017. Trees, forests and water: Cool insights for a hot world. *Global Environmental Change* **43**, 51-61.
- Fahey B, Jackson R.** 1997. Hydrological impacts of converting native forests and grasslands to pine plantations, South Island, New Zealand. *Agricultural and Forest Meteorology* **84**, 69-82.
- Hornbeck JW, Adams MB, Corbett ES, Verry ES, Lynch JA.** 1993. Long-term impacts of forest treatments on water yield: a summary for Northeastern USA. *Journal of Hydrology* **150**, 323-344. doi:10.1016/0022-1694(93)90115-P
- Houghton RA, House JI, Pongratz J, Van Der Werf GR, DeFries RS, Hansen MC, Quéré CL, Ramankutty N.** 2012. Carbon emissions from land use and land-cover change. *Biogeosciences* **9**, 5125-5142.
- Jackson RB, Jobbágy EG, Avissar R, Roy SB, Barrett DJ, Cook CW, Farley KA, le Maitre DC, McCarl BA, Murray BC.** 2005. Trading water for carbon with biological carbon sequestration. *Science* **310**, 1944-1947.

- Jones HS, Beets PN, Kimberley MO, Garrett LG.** 2011. Harvest residue management and fertilisation effects on soil carbon and nitrogen in a 15-year old *Pinus radiata* plantation forest. *Forest Ecology Management* **262**, 339-347.
- Kabeya N, Chappell NA, Tych W, Shimizu A, Asano S, Hagino H.** 2016. Quantification of the effect of forest harvesting versus climate on streamflow cycles and trends in an evergreen broadleaf catchment. *Hydrological Sciences Journal* **61**, 1716-1727.
- Komatsu H, Kume T, Otsuki K.** 2010. Water resource management in Japan forest management or dam reservoirs? *Journal of Environmental Management* **91**, 814-823. DOI: 10.1016/j.jenvman.2009.10.011.
- Komatsu H, Kume T, Otsuki K.** 2011. Increasing annual runoff—broadleaf or coniferous forests? *Hydrol Process* **25**, 302-318.
- Krankina ON, Dixon RK, Kirilenko AP, Kobak KI.** 1997. Global climate change adaptation: examples from Russian boreal forests. *Climatic Change* **36**, 197-216.
- Landsberg JJ, Gower ST.** 1997. Applications of Physiological Ecology to Forest Management. Academic Press: San Diego, CA 354.
- Lane PNJ, Mackay SM.** 2001. Streamflow response of mixed-species eucalypt forests to patch cutting and thinning treatments. *Forest Ecology and Management* **143**, 131-142.
- Lynch JA, Corbett ES, Sopper WE.** 1980. Impact of forest cover removal on water quality. Pennsylvania State University. Research Paper 23–604. University Park, PA: Institute for Research on Land and Water Resource p.91.
- Lynch JA, Corbett ES.** 1990. Evaluation of best management practices for controlling nonpoint pollution from silvicultural operations. *Journal of the American Water Resources Association* **26**, 41-52. DOI: 10.1111/j.1752-1688.1990.tb01349.x
- Matyas C, Sun G.** 2014. Forests in a water limited world under climate change. *Environmental Research Letters* **9**, 085001.
- Payn T, Carnus JM, Freer-Smith P, Kimberley M, Kollert W, Liu S, Orazio C, Rodriguez L, Silva LN, Wingfield MJ.** 2015. Changes in planted forests and future global implications. *Forest Ecology and Management* **352**, 57-67.
- Pike R, Scherer R.** 2003. Overview of the potential effects of forest management on low flows in snowmelt-dominated hydrologic regimes. *Journal of Ecosystems and Management* **3**.
- Ruddell S, Sampson R, Smith M, Giffen R, Cathcart J, Hagan J, Sosland D, Godbee, J, Heissenbuttel J, Lovett S, Helms J, Price W, Simpson R.** 2007. The role for sustainably managed forests in climate change mitigation. *Journal of Forestry* **105**, 314-319.
- Sahin V, Hall MJ.** 1996. The effects of afforestation and deforestation on water yields. *Journal of hydrology* **178**, 293-309.
- Stednick JD.** 1996. Monitoring the effects of timber harvest on annual water yield. *Journal of Hydrology* **176**, 79-95.
- Stoneman GL.** 1993. Hydrological response to thinning a small jarrah (*Eucalyptus marginata*) forest catchment. *Journal of Hydrology* **150**, 393-407.
- Trabucco A, Zomer RJ, Bossio DA, van Straaten O, Verchot LV.** 2008. Climate change mitigation through afforestation/reforestation: a global analysis of hydrologic impacts with four case studies. *Agriculture, ecosystems and environment* **126**, 81-97.
- Troendle CA, Wilcox MS, Bevenger GS, Porth LS.** 2001. The Coon Creek water yield augmentation project: Implementation of timber harvesting technology to increase streamflow. *Forest Ecology and Management* **143**, 179-187.

**Van Dijk AIJM, Keenan RJ.** 2007a. Planted forests and water in perspective. *Forest Ecology and Management* **251**, 1-9.

**Watson F, Vertessy RA, McMahon TA, Rhodes B, Watson I.** 2001. Improved methods to assess water yield changes from paired-catchment studies: application to the Maroondah catchments. *Forest Ecology and Management* **143**, 189-204.

**Zhang L, Zhao FF, Brown AE.** 2012. Predicting effects of plantation expansion on streamflow regime for catchments in Australia. *Hydrology and Earth System Sciences*, **16**, 2109-2121.

### Supplementary Information

The following list of publications used in the data collection of this study:

**Anderson HW, Hoover MD, Reinhart KG.** 1976. Forests and water: effects of forest management on floods, sedimentation, and water supply. General Technical Report PSW-018. Berkeley, CA: US Department of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station **115**, 18.

**Andréassian V.** 2004. Waters and forests: from historical controversy to scientific debate. *Journal of hydrology* **291**, 1-27.

**Bent GC.** 2001. Effects of forest-management activities on runoff components and ground-water recharge to Quabbin reservoir, central Massachusetts. *Forest Ecology and Management* **143**, 115-129.

**Bosch JM, Hewlett JD.** 1982. A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration. *Journal of hydrology* **55**, 3-23.

**Brechtel HM, Fuhrer HW.** 1991. Water yield control in beech forest. A paired watershed study in the Krofedorf forest research area. 20<sup>th</sup> General Assembly of the International Union of Geodesy and Geophysics, Vienna, Austria **08**, 477-84.

**Cheng JD.** 1989. Streamflow changes after clear-cut logging of a pine beetle-infested watershed in southern British Columbia, Canada. *Water Resources Research* **25**, 449-56.

**Cosandey C, Andréassian V, Martin C, Didon-Lescot JF, Lavabre J, Folton N, Mathys N, Richard D.** 2005. The hydrological impact of the Mediterranean forest: a review of French research. *Journal of Hydrology* **301**, 235-49.

**Cosandey C, Andréassian V, Martin C, Didon-Lescot JF, Lavabre J, Folton N, Mathys N, Richard D.** 2005. The hydrological impact of the Mediterranean forest: a review of French research. *Journal of Hydrology* **301**, 235-49.

**Fahey B, Jackson R.** 1997. Hydrological impacts of converting native forests and grasslands to pine plantations, South Island, New Zealand. *Agricultural and Forest Meteorology* **84**, 69-82.

**Fritsch JM.** 1990. Les effets du défrichement de la forêt amazonienne et de la mise en culture sur l'hydrologie des petits bassins versants. PhD Thesis. Université des Sciences et Techniques du Languedoc, Montpellier, pp392.

**Hornbeck JW, Adams MB, Corbett ES, Verry ES, Lynch JA.** 1993. Long-term impacts of forest treatments on water yield: a summary for northeastern USA. *Journal of Hydrology* **150**, 323-44.

**Kabeya N, Chappell NA, Tych W, Shimizu A, Asano S, Hagino H.** 2016. Quantification of the effect of forest harvesting versus climate on streamflow cycles and trends in an evergreen broadleaf catchment. *Hydrological Sciences Journal*. **61**, 1716-27.

**Komatsu H, Kume T, Otsuki K.** 2011. Increasing annual runoff—broadleaf or coniferous forests?. *Hydrological processes* **25**, 302-18.

**Pearce AJ, Rowe LK, O'Loughlin CL.** 1980. Effects of clearfelling and slash-burning on water yield and storm hydrographs in evergreen mixed forests, western New Zealand. IAHS-AISH Publication (IAHS) p130.

**Pike R, Scherer R.** 2003. Overview of the potential effects of forest management on low flows in snowmelt-dominated hydrologic regimes. *Journal of Ecosystems and Management* **3**, 3.



**Rowe LK, Jackson R, Fahey B.** 2002. Land use and water resources: hydrological effects of different vegetation covers **SMF2167**, Report. 5.

**Sahin V, Hall MJ.** 1996. The effects of afforestation and deforestation on water yields. *Journal of hydrology*. **178**, 293-309.

**Scott DF, Scott DF.** 2000. A re-analysis of the South African catchment afforestation experimental data. Pretoria: Water Research Commission.

**Stednick JD.** 1996. Monitoring the effects of timber water yield harvest on annual. *Journal of Hydrology* **176**, 79-95.

**Swift Jr LW, Swank WT.** Long term responses of streamflow following clearcutting and regrowth/Réactions à long terme du débit des cours d'eau après coupe et repeuplement. *Hydrological Sciences Journal* **26**, 245-56.

**Troendle CA, Wilcox MS, Bevenger GS, Porth LS.** 2001. The Coon Creek water yield augmentation project: Implementation of timber harvesting technology to increase streamflow. *Forest Ecology and Management* **143**, 179-87.

**Webb AA.** 2009. Streamflow response to Pinus plantation harvesting: Canobolas State forest, southeastern Australia. *Hydrological Processes* **23**, 1679-89.