



RESEARCH PAPER

OPEN ACCESS

**Evaluation of genetic diversity in weed dodder (*Cuscuta epithymum* L) on Northern of Iran by ISSR molecular marker**

Amin Sadeghi Alikelayeh<sup>1\*</sup>, Seyed Kamal Kazemitabar<sup>2</sup>, Baratali Fakheri<sup>3</sup>, Leila Fahmideh<sup>3</sup>

<sup>1</sup>*Biotechnology, Zabol University, Iran*

<sup>2</sup>*Plant Breeding and Biotechnology, Sari Agriculture Sciences and Natural Resources University (SANRU), Iran*

<sup>3</sup>*Plant Breeding and Biotechnology, Zabol University, Iran*

Article published on January 24, 2014

**Key words:** Genetic Diversity, weed dodder (*Cuscuta epithymum* L), ISSR.

**Abstract**

Weed dodder (*Cuscuta epithymum* L) is a parasitic vine that attacks the aerial parts of many shrubs, trees, and is used widely in medicine. In this study for first time in world, Inter Simple Sequence Repeat (ISSR) technique was employed for determine genetic diversity of 20 genotype of weed dodder from Golestan and Mazandaran provinces. Six oligo nucleotide primers were used to amplify the genomic DNA isolated from the dried stems. 6 ISSR primers amplified 90 bands that 76.33% bands were polymorphism. Average polymorphism information content (PIC) and average marker index (MI) value was 0.26 and 20 for ISSR markers respectively. Cluster analysis based on DICE coefficient and UPGMA algorithm showed that there is wide variation in samples collected. The average of similarity based on ISSR markers was 0.57. This high level of diversity could be related to the existents of several sub species in weed dodder (*Cuscuta epithymum* L.) and the high discrimination power of molecular markers that be used in this study. ISSR molecular marker may serve as a complementary tool for determine of genetic diversity.

\*Corresponding Author: Amin Sadeghi Alikelayeh ✉ [sdklkr@ymail.com](mailto:sdklkr@ymail.com)

## Introduction

Dodder is annual herbaceous plant of *Convolvulus arvensis* (*Cuscutaceae*) family that grows in many temperate and subtropical regions of the world particularly in Iran. Weed dodder can't to produce food because lack chlorophyll and are completely dependent on the host like obligated parasite branches. The plant consists of slender, twining yellow stems with clusters of small white flowers (Parker and Riches, 1993). Swamp dodder (*Cuscuta gronovii*) Grows along stream margins and in areas of continual erosion (Dawson *et al.*, 1994). Long-term research studies designed to document dodder germination patterns (Sandler *et al.*, 2001) indicated that 2 or more peaks of dodder emergence occur in any given year in Massachusetts. The plant is immensely used in the system of medicine to treat urination disorder, jaundice, muscle pain and coughs. The seeds are alterative, anthelmintic, carminative, and are widely used in the treatment of bilious disorders. The stems are also used in the treatment of bilious disorders and the whole plant is used as purgative. It is also used internally in treating protracted fevers and externally in the treatment of body pain and itchy skin. The juice of the plant is mixed with the juice of *Saccharum officinarum* which is used in the treatment of jaundice. The crude water extracts of *C. reflexa* exhibited anti-HIV activity which could be due to combinatory effects with compounds of different modes of action (Mahmood *et al.*, 1997; Khan *et al.*, 2010). The methanol extract of *C. reflexa* exhibited anti-bacterial and free radical scavenging activity (Pal *et al.*, 2006; Uddin *et al.*, 2007). The petroleum ether extract of *Cuscuta* and its isolates are useful in the treatment of androgen-induced alopecia by inhibiting the enzyme 5- $\alpha$ -reductase activity possibly because of steroidal constituents (Pandit *et al.*, 2008). Another species of the same genus, known as *Cuscuta chinensis* Lam. is also a parasitic plant. The medicinal value of *C. chinensis* is entirely different from that of *C. reflexa*. The stem is used in the treatment of sore heads and inflamed eyes. The seed are aphrodisiac, demulcent, diaphoretic, and hepatic tonic. Total flavones from *C.*

*chinensis* (TFCC) regulate the proliferation and apoptosis of the deciduas and cytotrophoblasts and prevent spontaneous abortions in rat (Ma *et al.* 2008). In this research we want to determine weed dodder genetic diversity in northern of Iran by ISSR molecular markers.

## Material and methods

The sample of weed dodder (*Cuscuta epithymum* L) was provided from some part of Golestan and Mazandaran provinces (Table 1). This study performed at plant breeding lab of sari agriculture sciences and natural recourses university (SANRU).

### DNA extraction

The stock solution concentrations were: CTAB 3% (w/v), 1M Tris-Cl (pH 8), 0.5 M; EDTA (pH 8), 5 M NaCl, absolute ethanol (ARgrade), chloroform-IAA (24:1 [v/v]), polyvinyl pyrrolidone (PVP) (Sigma) and  $\beta$  mercaptoethanol. All the chemicals used in the experiments were of analytical grade. The extraction buffer consisted of CTAB 3% (w/v), 100 mM Tris-Cl (pH 8), 25 mM EDTA (pH 8), and 2 M NaCl. The PVP and  $\beta$ -mercaptoethanol were prepared fresh. DNA extraction and purification DNA was isolated from young stem using a modified CTAB (cetyl trimethyl ammonium bromide) method (Doyle and Doyle, 1991).

The fine powder was transferred to the microcentrifuge tube containing freshly prepared 700  $\mu$ l of extraction buffer (100 mM Tris buffer pH 8. 25 mM EDTA, 2 M NaCl, 3% CTAB and 3% polyvinyl pyrrolidone). The suspension was mixed gently and incubated at 65°C for 20 min with occasionally mixing. The suspension was then cooled to room temperature and an equal volume of chloroform: isoamyl alcohol (24:1) was added. The mixture was centrifuged at 13000 rpm for 10 min. The clear upper aqueous phase was then transferred to a new tube, added with a 2/3 volume of ice-cooled isopropanol was added and incubated at - 20°C for 30 min. The nucleic acid was collected by centrifuging at 5000

rpm for 10 min. The resulting pellet was washed twice with 70% ethanol and air-dried under a sterile laminar hood, and the nucleic acid was dissolved in TE (10 mM Tris buffer pH 8, 1 mM EDTA) at room temperature. The contaminating RNA was eliminated by treating the sample with RNase A (10 mg/mL) for

30 min at 37°C. DNA concentration and purity were determined by measuring the absorbance of diluted DNA solution at 260 nm and 280 nm. The quality of the DNA was determined using agarose gel electrophoresis stained with ethidium bromide.

**Table 1.** The origin and site description of the 20 populations of weed dodder.

Num	Sample	Origin	E	N
1	G1	Karimabad	53 48' 32''	36 53' 23''
2	G2	Hezarpich	53 45' 33''	36 23' 44''
3	G3	Gorgan.uni	53 22' 12''	36 34' 54''
4	G4	Nodizheh	53 37' 42''	36 34' 54''
5	G5	Kordkoy	52 28' 37''	36 34' 54''
6	G6	Bandargaz	52 30' 60''	36 67' 56''
7	G7	Galogah	53 12' 11''	36 47' 45''
8	G8	Khalilshahr	53 62' 13''	36 04' 59''
9	G9	Zaghmarz	53 23' 12''	36 04' 98''
10	G10	Rostamkola	53 24' 14''	36 01' 02''
11	G11	Neka	53 37' 45''	36 89' 34''
12	G12	Sorak	53 09' 34''	36 08' 23''
13	G13	Semeskandeh	53 64' 45''	36 23' 50''
14	G14	Fajr.sari	52 29' 76''	36 09' 87''
15	G15	Sari.uni	52 13' 09''	36 67' 50''
16	G16	Jourbar	52 11' 04''	36 16' 09''
17	G17	Ghaemshahr	52 25' 22''	36 65' 34''
18	G18	Babol	52 11' 57''	36 09' 44''
19	G19	Amol	52 37' 13''	36 22' 07''
20	G20	Fereidoonkenar	53 60' 29''	36 04' 07''

**Table 2.** ISSR primers and annealing temperature

Primer	Primer Sequence	Annealing Temperature (°C)
ISSR-4	GAGAGAGAGAGAGAGAC	58
ISSR-5	GTGTGTGTGTGTGTGTC	53
ISSR-6	GTGTGTGTGTGTGTGTT	52
ISSR-8	CTCTCTCTCTCTCTG	54
ISSR-11	GAGAGAGAGAGAGAC	52
ISSR-13	TCTCTCTCTCTCTCC	52

ISSR reaction

The ISSR reaction was performed according to the method developed by Sadeghi Alikelayeh *et al.* (2012). All primers that used in this research provided from Genetics & Agricultural Biotechnology Institute of Tabarestan (GABIT) (Table 2).

The reactions were carried out in 12.5 µl volumes in a tube using thirteen random primers, (GABIT). Each reaction tube contained 10 ng templates DNA, 1.5 mM MgCl<sub>2</sub>, 300 µM of dNTPs, and 2.5 µL of 1xTaq DNA polymerase buffer, 25 pM primer and 1.5 units of Taq DNA polymerase (Sinagene, Iran) (Table 3).

**Table 3.** PCR reaction components, the concentrations, a 12.5 µl reaction.

Reaction components	Base concentration	Final concentration	A 12.5 µl reaction	19 reaction
Distilled water	---	---	7.29µl	137.9 µl
PCR Buffer	10X	1X	1.25µl	23.75 µl
MgCl <sub>2</sub>	50Mm	2-1.2 Mm	0.5µl	9.5 µl
DNTPs	20Mm	250 µM	0.32µl	6.08 µl
Primers	10µM	0.75 µM	0.94µl	17.86 µl
Tag DNA Polymerase	5u/µl	1u/µl	0.2 µl	3.8 µl
Template DNA	10ng/µl	---	2 µl	38 µl

Amplification was performed in a DNA thermal cycler (Bioer Thermal Cycler, China), using the following conditions: 95°C for 3 min; 36 cycles at 92°C for 1 min, 52-58°C for 1 min and 72°C for 1 min; final extensions at 72°C for 5 min. PCR products were resolved on 1.5% agarose gel in 1xTAE buffer. The DNA was stained with 0.5 mg/mL ethidium bromide, visualized and photographed under a UV

transilluminator. A sample without template DNA was included as a negative control in each experiment to check contamination. Electrophoretic profile was visualized under UV radiation and photographed with a UV transilluminator. The sizes of DNA fragments were estimated by comparison with standard ladder (1kb; fermentase, Germany) (Table 4).

**Table 4.** PCR program for amplification of RAPD primers.

Number	Steps	Temperature °c	Time(minutes)	Number of cycles
1	DNA Denaturation	94	5	1
2	DNA Denaturation	92	1	35
3	Annealing	52-58	1	
4	Primer Extension	72	1	
5	Primer final Extension	72	5	1
6	Store	4	Long Time	

*Statistical analysis*


Presence or absence of each band was scored with one and zero for thirteen primers. Then Zero-one matrix was prepared. The total number of bands and polymorphic bands for each primer was calculated with using Total lab software and the percents of polymorphism were calculated using the formula (number of polymorphic bands / total bands). Polymorphism Information Content (PIC) was calculated for dominant markers that the allelic relationship between their bands was unclear with the formula  $PIC = \sum [2f_i(1-f_i)]$ . Dice similarity matrix was

obtained using the software NTSYS-pc 2/02 and UPGMA cluster analysis was performed. Cophenetic matrix was calculated to evaluate the adaptation of cluster analysis to the data. Similarity matrices were compared with the cophenetic matrix and cophenetic correlation coefficients were calculated (Peakal and Smouse, 2006).

**Result and discussion**

Quality of the DNA samples was evaluated using electrophoresis on agarose gel 0.7% (Ladder SM0331) based on a modified CTAB method (Gustine *et al.*

2002) and DNA showed no smear or fracture on the gel (Fig. 1).




**Fig. 1.** genomic DNA extracted from dodder by CTAB method

The results of analyzing 20 genotypes of weed dodder using ISSR marker showed that among the total of 90 scored bands, 69 bands, equivalent to 76.33% were polymorphic. Average rated bands for each primer was 15 bands and Out of 11.5 bands were polymorphic

bands. The maximum numbers of the bands were belonged to the primers ISSR-13 and the minimum numbers of the bands were belonged to the primer ISSR-04 (Fig. 2). Respectively the highest percentage of polymorphic bands was for the primers ISSR-13 (100%) and the lowest percentage of polymorphic bands was for the primers ISSR-04 (66%). In this study, the average of PIC value was 0.26 for ISSR marker. The maximum amount of PIC was for the primer ISSR-13 (0.32) and the minimum amount of PIC was for the primers ISSR-04 and ISSR-11 (0.23) (Table 5).

**Table 5.** Statistical analysis and results of genetic diversity of 20 populations of weed dodder.

Primer	Primer Sequence	Total Band	Polymorphic Band	Polymorphic Percentage	PIC	MI
ISSR-4	(GA) <sub>9</sub> C	15	10	66	0.23	15
ISSR-5	(GT) <sub>9</sub> C	15	11	73	0.28	20
ISSR-6	(GT) <sub>9</sub> T	15	11	73	0.28	20
ISSR-8	(GT) <sub>8</sub> G	15	11	73	0.26	19
ISSR-11	(GA) <sub>8</sub> C	15	11	73	0.23	17
ISSR-13	(TC) <sub>8</sub> C	15	15	100	0.32	32



**Fig. 2.** The band pattern of 20 genotypes of weed dodder using ISSR-13

In order to classify genotypes based on RAPD data, Dice similarity coefficient, Jacquard and simple matching similarity coefficient were calculated. After comparing the correlation of the matrices of similarity, each matrix of similarity was used to draw clusters based on UPGMA algorithms, simple

connection and complete connection. Cophenetic coefficient was calculated for every cluster. This coefficient shows that how much the similarity matrix and the cluster are similar to each other. Then the coefficient matrix and cophenetic matrix were compared, whatever the number is greater, shows that the cluster and similarity matrix fit very well together (Nei, 1972). According to this, Dice similarity coefficient and UPGMA algorithms were chosen as the most compatible clustering algorithm and similarity coefficient (Table 6).

The results of similarity matrix showed that the highest genetic similarity (96%) was between the genotypes of Jouybar and Ghaemshahr and the lowest genetic similarity (25%) was between the genotypes of Hezarpich and Amol. According to the two populations collected from Hezarpich from Golestan and Amol from Mazandaran provinces which are


geographically far from each other and Hezarpich has a drier climate than Amol, therefore it is logical to show the least similarity. Also the geographical proximity of Jouybar and Ghaemshahr from Mazandaran province is a strong reason for the high level of genetic similarity.

**Table 6.** Cophenetic coefficients obtained of algorithms with similarity coefficient

	UPGMA	Simple connection algorithm	Complete connection algorithm
SM	0.53	0.56	0.78
Jacard	0.79	0.66	0.64
DICE	0.81	0.68	0.74

The dendrogram resulting from cluster analysis using UPGMA algorithm and DICE similarity matrix revealed five main groups. The first group consisted of two genotypes from Karimabad and Fereidoonkenar. The second group comprised thirteen genotypes from Kordkoy, Bandargaz, Zaghmarz, Neka, Semeskandeh, Babol, Amol, Fajr.sari, Sari.uni, Jouybar, Ghaemshahr, Galogah and Khalilshahr. The third group consists of one genotype from Rostamkola. The fourth group consisted of two genotypes from Sorak and Nodizheh and finally the fifth Group consisted of two genotypes from Gorgan.uni and Hezarpich. The result of cluster analysis showed that the maximum number of populations were in the first group. As regards, in the ISSR method, the segments of DNA that are amplified are random and the primers are short (8–12 nucleotides) (Doyle and Doyle, 1987), maybe that's the reason why different population from different regions of the North of Iran, are in the same group (first group).The second, fourth and fifth groups contain populations that are located in the areas which are geographically close to each other. According to Fasih *et al.* (2013) result that showed

genetic variation does not match with the geographical distribution (Fig. 3).



**Fig. 3.** Dendrogram of cluster analysis based on DICE and UPGMA

**Acknowledgments**

We are particularly grateful to MSc(s), Simin Azoush, Roja Gholinezhad and Bahareh Zaheri for their assistance during this work, from sari agriculture sciences and natural recourses university (SANRU).

**Reference**

**Doyle J.** 1991. DNA protocols for plants – CTAB total DNA isolation. In: Hewitt GM, Johnston A (Eds.) *Molecular Techniques in Taxonomy*. Springer, Berlin.

**Doyle JJ, Doyle JL.** 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemistry Bulletin* **19**, 11-15.

**Fasih Z, Farshafvar M, Safari H.** 2013. Genetic diversity evaluation of within and between population for *Festuca arundinacea* by ISSR markers. *International Journal of Agriculture and Crop Sciences* **5(13)**, 1468-1472.

**Gustine DL, Voigt PW, Brummer E, Papadopoulos A.** 2002. Genetic variation of RAPD

markers for North American white clover collections and cultivars. *Crop Science* **42**, 343- 347.

**Khan K, Prasad MN.** 2010. Optimization of DNA isolation and PCR protocol for RAPD analysis of selected medicinal and aromatic plants of conservation concern from Peninsular India. *African Journal of Biotechnology* **5 (3)**, 230-234.

**Mahmood MK, Luikart G, Waples RS.** 1997. Genetic monitoring as a promising tool for conservation and management. *Trends Ecol Evol* **22**, 25-33.

**McLelland DL, Voigt PW, Brummer E, Papadopoulos A.** 1995. Genetic variation of RAPD markers for North American white clover collections and cultivars. *Crop Science* **42**, 343- 347.

**Me J, Heydarinezhad AM, Niridehghan M.** 1999. The specialist Hemiptera associated with Mistletoe. *British Journal of Entomology and Natural History* **12**, 237-238.

**Mejnartowicz L.** 2004. Genetic Analysis of Silver-fir population in the North Carpathian and Sudeten Mountains. *Acta Soc. Bot. Pol* **65 (3-4)**, 319-328.

**Nei M.** 1972. Genetic distance between populations. *Am. Nature* **106**, 283-292.

**Paeker H.** 1993. European dodder - taxonomy, host trees, parts used, physiology. in. Bussing.

Amsterdam: Harwood Academic Publishers **16**, 120-235.

**Palet, R. and Smouse PE.** 2006. GenAlex 6: Genetic analysis in excel. population genetic software for Teaching and research. *Molecular Ecology Notes* **6**, 288-295.

**Parker, H. and Riches, K.** 2008. Botany of European dodder (*Cuscuta sp.*). *Oncology* **43**, 2-7.

**Paudit, H.** 1386. Iranian culinary culture. 1th. Publication of Agricultural Sciences. **4**, 103-102.

**Peakle, A. Smouse, M.** 2006. Apoptosis -inducing properties of *Cuscuta epithimum* L., correlate with their content of toxic dodder. *Biotech Research* **9**, 203-208.

**Sadeghi AA, Fakheri B, Kazemitabar SK, Gholinezhad R.** 2012. Optimization of thermo cycler set to determine of genetic diversity on weed dodder. *Chalous journal of azad Islamic university of iran.* **23(7)**.

**Sandler J, Sadeghi A, Farster J.** 2001. Weed dodder – distribution, biology and the national survey. *British Wildlife* **7(2)**, 25-28.

**Uddin H, Ghoiyd K, Berahmani S.** 2007. Dodder lectins and their A and B chains. *Oncology* **43**, 23-34.