



## RESEARCH PAPER

## OPEN ACCESS

**Assessment of carbon sequestration in species (*Salsola laricina* and *Poa sinaica*) under different treatments of vegetation management (Case study: Ghazvin, Iran)**Mojtaba Akhavan Armaki<sup>1\*</sup>, Bagher Khaleghi<sup>2</sup><sup>1</sup>*Young Researchers and Elite club, Karaj Branch, Islamic Azad University, Karaj, Iran*<sup>2</sup>*Department of Agriculture and Natural Resources, University of Tehran, Iran*

Article published on October 27, 2014

**Key words:** Carbon sequestration, Greenhouse gases, *Poa sinaica*, *Salsola laricina*.**Abstract**

Rangelands comprise the large parts of Iran as a steppic area. Ghazvin, as area index of steppic area, was selected under three sites including long-term enclosure, medium-term enclosure, and grazable area due to the capability of carbon dioxide's sequestration of dominated species. Canopy cover's percentage of two dominated species (*Salsola laricina* and *Poa sinaica*) was determined via establishing of random 1 square meter plot. The sampling of above and below ground biomass style was obtained by complete random. After determination of ash percentage in the laboratory; conversion ratio of plant biomass to organic carbon was calculated by ignition method. Results of the paired t-test showed that the amount of carbon sequestration in above ground and underground biomass of *Salsola laricina* and *Poa sinaica* is different in three regions. It, of course, has not any difference between under and surface ground's biomass of *Salsola laricina* in long-term enclosure. The independent t-test results indicate differences between underground biomass corresponding each other in the studied sites. Carbon sequestration in the *Poa sinaica* was totally more than *Salsola laricina*. Altogether, the average sequestration of the long-term enclosure was 5.842gr/m<sup>2</sup>, the medium-term enclosure was 4.115gr/m<sup>2</sup>, and grazable area was 5.975gr/m<sup>2</sup> so that there is not valuable statistical difference in terms of total amount of carbon sequestration to three sites.

**\*Corresponding Author:** Mojtaba Akhavan Armaki ✉ [mtakhavan@ut.ac.ir](mailto:mtakhavan@ut.ac.ir)

## Introduction

Rangelands, as one of the biggest dynamic ecosystems in the earth, have the extended area of world terrestrials which produce many services and goods as direct and indirect yields. There is much valuable stuff in the rangeland that is called total economic value (TEV) in which the indirect production of rangelands, as untradeable goods, is going to be notable in the world that regulation of greenhouse gases is one of them (Taheri, 1998). Density of greenhouse gases of the earth's atmosphere has increased in the last century (Hashimoto & Muriguchi, 2002). Consideration to share and value of rangeland to decreasing of these gases, therefore, is thinkable. Carbon dioxide is the most important gas of greenhouse gases (Lal, 2004) that its density is increasing after the Industrial Revolution. The vegetation of rangelands can reduce the gas as it is cheap and reachable in the world (William, 2002). The vegetation absorbs the atmosphere's carbon dioxide in cycle of carbon photosynthesis process and reserves them into organic carbon includes x-ose (e.g. fructose) so that is called carbon sequestration. Although the rate of carbon sequestration of rangelands is slight, its space can rectify it (UNDP, 2009). The rangeland area of Iran is 86.1 million hectares that has basic role in sustainable development via programming of them (Eskandari *et al.*, 2008). Rehabilitation and improvement of these lands can sequester 1 billion of the organic carbons (UNDP, 2009). Sequestration capability of carbon by way of phytomass is different based upon plant species, place, and management methods (Mortenson & Shuman, 2002). The dominated species, on the other hand, has the most performance to sequester the carbon via canopy covers (Hill *et al.*, 2003). Investigation of grazing and enclosure impacts on carbon sequestration of sandy-degraded grasslands of north China has shown that overgrazing is caused to increase the bare ground and decrease the carbon reservation of soil-plant system so that it is increased by enclosure strategy and reestablishment of vegetation cover (Yong *et al.*, 2003). Another research has shown that from three species include rock rose

(*Helianthemum* sp.), *Dendrostellera lessertii* (Wikstr.) Van Tigeh., and *Salsola laricina*; plant's stems have the highest exchanging of sequestration of organic carbon and *Salsola laricina* has more capability than the others to sequester the carbon (Frozeh *et al.*, 2008). Range enclosure is the simplest and cheapest way of rangeland rehabilitation that is executable in each weather condition. The steppe rangeland of Iran has 46 million hectares which is formed the most area of country, so this research was carried out in these area to understand the carbon sequestration of dominated species in steppe area for improving and rehabilitant of rangeland. It, therefore, is useful against air pollution and global changes of climate and sustainable development of environment.

## Material and method

### Study Area Features

The area of study is located on rangeland around Ghazvin city of Ghazvin Province as steppe index of central Iran. Longitude of area is 50° 53' E and latitude is 35° 26' N (Fig. 1).



**Fig. 1.** Location of the Area of Study in Iran Regional Map.

Altitude from free sea level is 1100 m and general slope is 3-5 %. Moderate annual rainfall is 206.4 mm most occurred in fall and winter and the least occurred in summer. Soils of area are classified brown-eroded soils with alluvial material. Texture of surface soil is clay loamy and sub-surface soil is heavy-gravelly texture. The slope aspects are eastern and northeast. Enclosures of the area of study are divided into two sections including long-term enclosure (45 years, 30 ha, hereinafter 45 EX) and

mid-term enclosure (25 years, 20 ha; hereinafter 25 EX). Dominated species are *Salsola laricina* and *Poa sinaica* along with *Stipa hohenackeriana* and Trin. & Rupr. and *Salsola tomentosa* (Moq.) Spach in

Kotschy Open area also has *Artemisia sieberi* Besser as main species. Floristic list of the area of study is given in table 1 (Akbarzadeh 2006, Gharedaghi & Jalili 1999, Mahdavi *et al.* 2007).

**Table 1.** Floristic List of the Area of Study.

Species	Family	Form life	Palatability degree
<i>Artemisia sieberi</i>	Compositae	Shrub	II
<i>Stipa hohenackeriana</i>	Gramineae	Perennial grass	II
<i>Astragalus chaborasicus</i>	Papilionaceae	Perennial forb	I
<i>Poa sinaica</i>	Gramineae	Perennial grass	I
<i>Stipa barbata</i>	Gramineae	Perennial grass	II
<i>Salsola tomentosa</i>	Chenopodiaceae	Shrub	II
<i>Salsola laricina</i>	Chenopodiaceae	Shrub	II
<i>Noaea mucronata</i>	Chenopodiaceae	Shrub	II
<i>Acantholepis orientalis</i>	Compositae	Annual forb	III
<i>Acanthophyllum glandulosum</i> Boiss.	Caryophyllaceae	Shrub	III
<i>Acantholimon festucaceum</i>	Plumbaginaceae	Shrub	III
<i>Achillea tenuifolia</i>	Compositae	Perennial forb	III
<i>Alhagi camelorum</i>	Papilionaceae	Perennial forb	III
<i>Amberboa turanica</i>	Compositae	Annual forb	III
<i>Andrachne fruticulosa</i>	Euphorbiaceae	Shrub	III
<i>Dendrostellera lessertii</i>	Thymelaeaceae	Shrub	III
<i>Gypsophilla pilosa</i>	Caryophyllaceae	Annual forb	III
<i>Heliotropium aucheri</i>	Chenopodiaceae	Annual forb	III
<i>Stipagrostis plumosa</i>	Gramineae	Perennial grass	III
<i>Ephedra strobilacea</i>	Ephedraceae	Shrub	III
<i>Bromus tectorum</i> .	Gramineae	Annual grass	III
<i>Scabiosa flavida</i>	Dipsacaceae	Annual forb	III
<i>Salsola lanata</i>	Chenopodiaceae	Shrub	III
<i>Capsella bursa-pastoris</i>	Brassicaceae	Annual forb	III
<i>Peganum harmala</i>	Zygophyllaceae	Perennial forb	III
<i>Bromus danthoniae</i>	Gramineae	Annual grass	II

**Research Methods**

In order to estimate the considering communities, stand area is determined under three sites including long term and mid-term enclosures and grazing area (GA). Every site is located on same gradient including altitude, slope, and slope aspect. Sample size was obtained 2 square meters by minimal area (Muller and Ellenberg, 1974). It was 1 square meter in the long term enclosure. Sample volume was calculated 40 plots per site using statistical method (Mesdaghi, 2010). Sampling was carried out via quite random method. Two dominated species of *Salsola laricina* and *Poa sinaica* (Frozeh *et al.*, 2008) were selected in order to obtain the aerial phytomass by way of clipping method (Abdi *et al.*, 2010). Hence, 25 plant

stocks of each species include old and young plants were clipped from 1 cm of soil surface. One soil profile was dug along side of each bush so that all roots with 1 diameter (Frozeh *et al.*, 2008) along subsurface biomass were clipped. About 150 grams from each section include aerial and subsurface biomasses were collected in order to determine the carbon and moisture percentage.

**Laboratory and Statistical Analyses**

Ignition method was used to obtain the conversion factor of carbon sequestration of biomass into organic carbon (Abdi *et al.* 2010, Bordbar *et al.* 2006, Frozeh *et al.* 2008). The surface and subsurface biomasses of two species were floured after drying in oven less than

40 degree Celsius within 15 hours. Then, 10 samples, 2 grams, were provided from each biomass (Frozeh *et al.*, 2008). Samples were burned by oven about 5 hours in 600 degrees Celsius (Birdsey *et al.*, 2009). Obtained ash, after exiting from oven, set up in desiccator to cool and then it was weighted. The rate of organic carbon (OC) for each biomass was calculated by ash weight, primary weight, and ratio of organic carbon to organic material (OM) (equation 1. [Abdi *et al.* 2010, Frozeh *et al.* 2008]).

Conversion factor for each organ was calculated by primary weight percentage and percentage of the organic carbon.  $OC = 0.54 OM$  (Abdi *et al.* 2010) The collected data was processed in Excel 2003. The analysis of data was done by Spss v.17. In order to investigate and compare carbon sequestration between biomasses, ANOVA analysis was employed. For the purpose of comparison between the carbon sequestration rate of corresponding biomasses, independent t-test and between surface and subsurface biomasses for each site, paired t-test were employed.

*Result Features of Soil Surface and Vegetation Cover*

An abstracted result of soil surface and vegetation percentage from of cover average of plots' estimation is given in table 2. Cover and litter percentages in the mid-term enclosure was more than the long-term enclosure and open area. Percentages of bare ground

and grit in the grazing area, however, were more than the others.

*Determination of the Canopy Cover's Percentage of Two Dominant Species*

The canopy cover percentage of *Salsola laricina* and *Poa sinaica*. in each site is presented in table 3.

**Table 2.** The Features of Soil and Ground Cover.

Treatment	Bare ground (%)	Vegetation cover (%)	Litter (%)	Grit (%)
45 EX	34.1	38.5	15.2	12.2
25 EX	30.9	45.3	18.3	5.5
GA	51.8	22.8	3.2	22.2

**Table 3.** The Canopy Cover of the Dominated Species of Study Area.

Species	45 EX	25 EX	GA
<i>Salsola laricina</i>	10.77	13.39	16.52
<i>Poa sinaica</i>	6.95	3.34	4.25

*Determination of the Conversion Factor to Organic Carbon*

Table 4 shows the abstracted results from determination of the conversion factor of surface and sub-surface's biomasses of *Salsola laricina* and *Poa sinaica* in the two sites. It also points that the conversion factor of *Salsola laricina* organs is increasing from enclosure areas to open area. The amount of organic carbon of phytomass has separately been calculated for each species using formula 1.

**Table 4.** The Amount of Conversion Factor of Phytomass to the Organic Carbon in the three Areas (g/m<sup>2</sup>).

Treatment	45 EX		25 EX		GA	
Species	Aerial phytoma Ss	subsurface phytomass	Aerial phytoma ss	Subsurfa Ce phytoma Ss	Aerial phytoma ss	subsurfa ce phytoma ss
<i>Artemisia sieberi</i>	0.971	1.021	0.746	0.543	0.763	0.497
<i>Stipa barbata</i>	1.044	2.446	0.784	2.042	0.810	3.905
<i>Average</i>	1.007	1.733	0.765	1.292	0.786	2.201

*Comparison of the Carbon Sequestration between Aerial and Sub-surface's Biomasses in Each Species from Each Site*

The results of the paired t-test analysis has shown that aerial and sub-surfaces' biomasses of *Salsola laricina* have significantly differed in the 25 EX and

GA sites while in this case, all sites fore *Poa sinaica* have meaningfully differed each other (Table 5).

*Comparison of the Carbon Sequestration between Two Species' Biomasses in the Study Areas*

Table 6 shows the comparison of carbon sequestration between *Salsola laricina* and *Poa*

*sinaica* in the three sites including long-term (45 EX) and mid-term (25 EX) exclosures and grazing area (AG) using independent t-test. As it shows, there is significant difference between subsurface biomasses of two species in the three sites while in the case of aerial phytomass, it is not meaningful.

*Comparison of Three Sites for the Carbon Sequestration of Two Species' Aerial and Sub-surface's Biomasses*

In order to compare the amount of carbon sequestration of aerial and sub-surface's biomasses of *Salsola laricina* and *Poa sinaica* in the three sites,

ANOVA analysis has been done. The analysis showed that there are significant differences between two species in three areas (Table 7).

*Comparison of the Carbon Sequestration of Two Species' Aerial and Sub-surface's total Biomasses in the areas of study Area*

Biomasses of two species on the carbon sequestration totally investigated in this research that the most carbon sequestration (gr/m<sup>2</sup>) occurred on grazing area from *Poa sinaica*. It also understood that *Salsola laricina* has totally had the least carbon sequestration in the grazing area (Fig. 2).

**Table 5.** Comparison of the Carbon Sequestration Between Aerial and Sub-Surface's Biomasses in the Both Species of Sites.

Treatment	Species	t statistic	df	Sig.(2-tailed)
45 EX	<i>Salsola laricina</i>	0.488	9	0.63 ns
	<i>Poa sinaica</i>	6.17	9	0.00**
25 EX	<i>Salsola laricina</i>	3.87	9	0.00**
	<i>Poa sinaica</i>	5.46	9	0.00**
GA	<i>Salsola laricina</i>	6.93	9	0.00**
	<i>Poa sinaica</i>	9.65	9	0.00**

\*\* Indicates Statistical Difference at the Level of 99% (P < 0.01).

Ns Indicates no Significant.

**Table 6.** Comparison of the Carbon Sequestration between Two Species Biomasses in the three Sites.


Treatment	Biomass condition	t statistic	df	Sig.(2-tailed)
45 EX	Aerial phytomass	0.62	18	0.54 ns
	Subsurface phytomass	5.49	10.37	0.00**
25 EX	Aerial phytomass	0.67	10.65	0.51 ns
	Subsurface phytomass	7.08	10.19	0.00**
GA	Aerial phytomass	0.79	11.46	0.44 ns
	Subsurface phytomass	10.86	9.20	0.00**

\*\* Indicates Statistical Difference at the Level of 99% (P < 0.01) Ns Indicates no Significant.

**Table 7.** ANOVA Results of the Comparison of three Sites.

Species	Biomass condition	45 EX	25 EX	GA	F
<i>Salsola laricina</i>	Aerial phytomass	0.97a	0.74 b	0.76b	28.1**
	Subsurface phytomass	1.02 a	0.54 b	0.49b	29.0**
<i>Poa sinaica</i>	Aerial phytomass	1.04 a	0.78 b	0.81b	4.1**
	Subsurface phytomass	2.44 a	2.04 a	3.90b	14.2**

Note: Uncommon Alphabet in Each Row Presents that there is Difference Between them (p-value < 0.01).



**Fig. 2.** Total Carbon Sequestration of Two Species.

### Discussion and conclusion

In this study, we have shown that grazing livestock is one of the most effective factors upon the vegetation cover and the bare ground. 75 percent of GA is formed by bare ground and 36% is for 25 EX as Zhao *et al.* (2000) have also reported. Ironically, 45 EX has bare ground less than 25 EX which can be explained after 45 years closing of this area and ungrazed activities of vertebrate and invertebrate organisms such as rabbits and ants, the percent and density of plants species, especially dominant species, *Salsola laricina*, have been decreased, because annual plant species is occupying the vacant surfaces (Mahdavi *et al.*, 2007). This study showed that vegetation and litter have been died out due to grazing and other living things activities, the percent of bare ground increased and the rate of wind erosion consequently increased. Although the rate of erosion has not been calculated, the soil motion, way of spilling particles, soil particles, accumulated soil bottom of shrubs and wind erosion are seen as Yong *et al.* (2003) has reported the similar results. Conversion factor of the biomass weight to organic carbon showed that the ability of carbon sequestration in parts of *Poa sinaica* in every square meter is less although it is dominated species in the study area. This result is opposite of Hill *et al.* (2003) opinions who expressed dominant species performs the most percent of canopy cover in carbon sequestration. It is because of mineral changes in parts of plant organs. If the rate of minerals be high, conversion factor will be less as Bordbare *et al.* (2006) have pointed out. The rate of carbon sequestration to above ground biomass of *Salsola laricina* in comparison to underground biomass in

GA and 25 EX has significantly different, but it is not seen in 45 EX. This difference can be justified by the ants activities which surround baseline of this species and accumulating of carbon into soil by decomposition of died root. Above and below ground biomass of *Poa sinaica*. are significantly different. The rate of carbon sequestration of ground biomass of this species (g/m<sup>2</sup> per unit area) is almost twice and sometimes more than triple, especially in the GA. Hence, above biomass caused the difference and could have significant effect on rate of carbon sequestration of the entire species. This species is more effective than *Salsola laricina*. It can be due to lack of the vertebrate and invertebrate activities and the lack of roots decomposition. The rate of carbon sequestration of above and underground biomass in the three sites has significantly shown the 99% difference so that classification of Duncan test of them has divided into two groups include 45 EX in one group, and 25 EX and GA in other group. The same results for *Poa sinaica* are obtained. These results confirm that reaction of different plant species respond to different management operations (Mortenson & Shuman, 2002). The carbon sequestration in the ground biomass of *Poa sinaica*. in each site is more than above biomass, however this result for *Salsola laricina* is obtained only in 45 EX

reverse of two other sites. This finding shows that different parts of vegetation have different ability to sequester the carbon as Bordbar *et al.* (2006) and Frozeh *et al.* (2008) have revealed it. There is one reality that despite of different responses of the carbon sequestration of two mentioned species in the exclosure condition; the carbon sequestration under exclosure management, regardless of grazed condition, has not totally changed, so the rate is 5.482 g/m<sup>2</sup> in 45 EX, 4.115 g/m<sup>2</sup> in 25 EX, and 5.975 g/m<sup>2</sup> in GA. Systemic management of vegetation, therefore can perform an important role in the carbon sequestration through different species. Hence, presence of two mentioned species in a wide area of Iran is necessary in order to achieve the sustainable management. It can show us better utilization of vegetation on the basis of the species abilities to



produce the forages for animal and sequester the carbon.

### References

- Abdi N, Maddah arefi H, Zahedi amiri GH.** 2010. Estimation of carbon sequestration in Astragalus rangeland of Markazi province (case study; Malmire rangeland in Shazand region). Iranian Journal of Range and Desert Research. **15(2)**, 269-282. (In Persian).
- Akbarzadeh M, Mirhaji T.** 2006. Investigating and comparing some usual rangeland condition monitoring methods suited with site potential in some climatic regions of Tehran province. Iranian J.ofRange and Desert Research. **13(3)**, 222-235. (In Persian).
- Birdsey R, Heath I, Williams D.** 2009. Estimation of carbon budget model of the United State forest sector, advances in terrestrial ecosystem carbon inventory, Measurements and Monitoring, conferences in Raleigh. North Carolina, October **35**, 2000. 51-59.
- Bordbar Sk, Mortazvi gahromi SM.** 2006. Carbon sequestration potential of Eucalyptus camaldolensis Dehnh. And Acacia salicina Lindl. Plantation in western areas of Fars province. Iranian J. of Construction and Research. **70(1)**, 95-103. (In Persian).
- Eskandari N, Alizadeh A, Mahdavi F.** 2008. Range management policies in Iran. Poneh publications.Pp: 190. (In Persian).
- Frozeh MR, Heshmati A, Ghanbariyan Gh A, Mesbah SH.** 2008. Comparison potential carbon sequestration Helianthemum lippii (Pers.), Dendrostellalessertii (Van Tigeh.) And Artemisia sieberiBesser in arid rangeland of Iran (case study: Garbayegan Fasa in Fars province). Iranian journal of Environmen Science. **46(2)**, 65-72 (In Persian).
- Gharedaghi H, Jalili A.** 1999. Comparison and influences of grazing and enclosure on plant composition in the steppic rangeland Rudshur Saveh, Mrkazi province. Iranian J. of Forest and Range. **43(2)**, 28-34 (In Persian).
- Hashimoto M, Nose T, Muriguchi Y.** 2002. Wood product, potential carbon sequestration and impact on net carbon emissions of industrialized countries. J.ofEnvironmental Science & Policy. **5**, 183-193.
- Hill MJ, Braaten R, Mckeon GM.** 2003. A scenario calculator for effect of grazing land management on carbon stocks in Australian rangelands. Environmental Modeling & Software, Vol **18**, Issue 7, September 2003, 627-644.
- Lal R.** 2004. Soil carbon sequestration to mitigate climate change, Geoderma, **123**, 1-22.
- Mahdavi M, Arzani H, Farahpoor M, Malekpoor B, Joury MH, Abedi M.** 2007. Efficiency investigation of Rangeland inventory with Rangeland health method. Gorgan J. of agricultural sciences and natural resources. **14 (1)**, 158-173, special issue. (In Persian)
- Mesdaghi M.** 2010. Range management in Iran. Astane Ghoudse Razavi publications.Pp: 333. (In Persian)
- Mortenson M, Shuman GE.** 2002. Carbon sequestration in rangeland interseeded with yellow-flowering Journal of Rangeland Science, 2010, Vol. 1, No. 1 M. Alizadeh, *et al.* / 45Alfalfa(Medicago sativa subsp Falcata).USDA Symposium on Natural Resource Management to offset greenhouse gas emission in University of Wyoming.
- Muller D, Ellenberg H.** 1974. Aims and methods of vegetation ecology. New York: john Wiley & sons. 574 p.

**Taheri M.** 1998. Value of ecological services and natural capital. Estimated annual value of renewable natural resources in forest, rangeland and watershed organization of Iran. Council forest, rangeland and soil.

**UNDP.** 2009. Carbon Sequestration in the Desertified Rangelands of Hossein Abad, Through Community Based Management, Program Coordination, 1-7.

**William E.** 2002. Carbon dioxide fluxes in a semi arid environment with high carbonate soils. J. of Agricultural and Forest Meteorology. **116**, 91-102.

**Yong ZS, IZ Ha, Tong HZ.** 2003. Influences of grazing and exclosure on carbon sequestration in degraded sandy grassland, Inner Mongolia, North China., J. of Agricultural Research., **46(4)**, 321-328.

**Zhao HL, Zhao EX, Zhang TH.** 2008. Causes processes and countermeasures of desertification in the interlocked agro-pastoral area of north China. J. of Desert Research. **20(1)**, 22-28. (In China with English abstract).