



## RESEARCH PAPER

## OPEN ACCESS

## The effect of polyacryl amide a 200 polymer on optimization of moisture and rooting indices of *Rosa sp.*

Kamran Parvanak<sup>1\*</sup>, Hadi Chamheidar<sup>2</sup>

<sup>1</sup>Department of Soil Science, Yadegar-E- Imam Khomeini(RAH) Branch, Islamic Azad University, Tehran, Iran

<sup>2</sup>Department of Soil Science, Shoushtar branch, Islamic Azad University, Shoushtar, Iran

**Key words:** Polyacryl amide A 200, optimization, irrigation interval, *Rosa Sp.*

<http://dx.doi.org/10.12692/ijb/5.6.119-127>

Article published on September 20, 2014

### Abstract

This study aimed to evaluate the impact of four Polyacryl amide A 200 polymer rates (zero, 0.15, 0.3 and 0.6% w/w) and four irrigation intervals (1, 2, 3 and 4 days) on the amount of available water and some rooting indices of *Rosa Sp.* The experimental layout consisted of factorial arranged in a complete block randomized design with three replicates. The results of variance analysis showed that various polymers rates, irrigation interval and their interactions had significant effects on the amount of available water and rooting indices (rooting percentage, fresh and dry weight, length, diameter and number of root) ( $p \leq 0.01$ ). Based on the mean comparison, the applying polymers independently as much as 0.3% increased rooting indices of cuttings significantly, while the irrigation interval had negative effect on the rooting indices of the plant. Interaction of irrigation interval and the use of polymers showed the highest rooting percentage (83.3%), root fresh and dry weight (4.1 and 3.2 g respectively), root elongation and diameter (5.7 and 2.1 cm respectively) in treatment 0.3% polymer and 2 days irrigation interval that had significant difference relative to control and other treatments ( $p \leq 0.01$ ). Application of polymers in 0.3 and 0.6%, increased the amount of available water 2 to 2.75 times respectively. According to these results, applying of 0.3% polymer with 2 days irrigation intervals also produced cuttings with good rooting and saving 50% irrigation water resources costs as compared to the daily irrigation.

\*Corresponding Author: Kamran Parvanak ✉ [kparvanak@iausr.ac.ir](mailto:kparvanak@iausr.ac.ir)

## Introduction

The shortage of fresh water and growing competition for clear water makes less water available for agriculture production. The great challenge for the coming decades will be the task of increasing food production with less water, particularly in countries with limited water and land resources. Iran is one of the driest countries in the world that the shortage of water in its agriculture is the most important development bottleneck. Also, since the era of cheap water harvesting and water potential of the country can no longer meet the needs of the growing demand for water, particularly in the agricultural sector, so shall be forced to optimize water use. Although all flowers are beautiful, but *Rosa Sp* flowers are interested in by many people. The culturing of *Rosa Sp* is an economic job today and culturing of it is customary in cities (Malkeshi and Mojib Hagh Ghadam, 2009). Some factors such as long period dormancy of seeds and high amount of hollow and dead seeds reduce germination percentage of the different *Rosa Sp* so its vegetative propagation has particular importance for seedling production of the plant (Merkle and Nairn, 2005). Replication media of nurseries includes a large coarse minerals portion to raise the proportion of large pores filled with air, but existence of coarse minerals (such as sand and gravels), which have low water holding capacity and high deep water percolation, cause moisture which is the most important factor in rooting of cuttings, easily be drained. Water depletion could loss irrigation water resources and also reduces the physiological and metabolic processes which are effective in rooting of cuttings (Akhter *et al.*, 2004). The usage of hydrogel to attract and increase the water holding capacity of the seedling rooting media is considered recently (Peterson, 2002). Their application in seedling rooting media could eliminate moisture stress and helping adaption of planted crops with environment. The hydrogel also raised the water efficiency and therefore could increase the irrigation interval (Huttermann *et al.*, 1999). Jandagheyan, 2001 investigated the effect of polyacrylamide hydrogel and irrigation interval on the rooting of geranium and philodendron. The results showed that with the

increasing usage of hydrogels from zero to 50 percent by volume, rooting percentage, plant height, fresh and dry root weight increased. The maximum value of these indices were observed in geranium in 0.3% hydrogel with 6 days irrigation intervals and in philodendron in 0.5% hydrogel with 3 days irrigation intervals. Ghasemi and Khoshkhoe, 2007 studied the effect of superabsorbent polymers on irrigation interval and growth and development of *Chrysanthemum morifoliumramat*. The results showed that using the hydrogel had a significant effect on vegetative parameters. All vegetative mean parameters except the ratio of root to shoot did not significantly differ in 0.6% hydrogel in 4 days irrigation intervals treatments with control treatment (no hydrogel) in the 2 days irrigation intervals. Assareh *et al.*, 2007 investigated yew rooting and causes of their hard and late rooting of cuttings. Results showed that the highest rooting was in pure sand and absorbent hydrogel in addition of indole butyric acid hormone treatment. Farokhzad *et al.*, 2009 studied the effect of polyacrylamide polymer and indole butyric acid treatments on rooting of cuttings of *Magnolia soulougenae*. The results showed that the percentage of rooting of cuttings increased from 26.66% in control to 70% in the 3000 ppm indole butyric acid with 0.8% super absorbent treatment. Meghani *et al.*, 2010 investigated the effect of media culture on rooting of *Bougainvillea glabra* cuttings. Their results indicated that the highest rooting percentage with an average of 75% was obtained from 50% sand with 0.5% moisture absorbent material. Research conducted by Huttermann *et al.*, 1999 indicated that with addition of stockosorb hydrogels to soil to 0.4% w/w after 17 days, all 45 seeds germinated. Jalili *et al.*, 2011 studied the effect of four different levels of hydrogel and irrigation interval on the growth of rose seedlings. Their finding indicated that the number of branches, branch height, canopy diameter, 10 days irrigation interval with using 0.4% hydrogel treatments were most appropriate. This study aimed to evaluate the application of the hydrogel to supply optimal moisture needed for rooting of *Rosa Sp* cuttings and adjust the irrigation interval to result in the saving of irrigation water.

## Materials and methods

In order to investigate the effect of Polyacryl A/mide A 200 polymer and irrigation interval on available water and some rooting properties of cutting of *Rosa Sp*, an experimental layout consisted of factorial arranged in a complete block randomized design with three replicates and each replication consist of six cuttings was carried out in the greenhouse in Isfahan city in 2012-2013. Treatments were four hydrogel levels consists of 0 (control), 0.15, 0.3, 0.6% w/w and four irrigation intervals (1, 2, 3 and 4 days). At the bottom of each pot (10 × 10 cm), 1 cm thick of bran was added to helping drainage. Then, the hydrogel treatments and sand mixtures which mix to a depth of 8 cm of pots was poured into the pots. Taking of cuttings of the young shoots were performed in early autumn and about 3 to 4 inches away from the bottom of the cuttings immediately entered the 2/1000 concentration of Benomyl fungicides for disinfection and then were planted in pots cultures. After establishment of cuttings the amounts of irrigation water by 40% allowable depletion of soil moisture for the sand that used in the rooting context, were calculated by the moisture content by volume of the sand field capacity, permanent wilting point and pot sizes and were distributed for each pot and each irrigation interval on the pot surface. While careful monitoring of pots during rooting, factors such as rooted cutting percentage in each treatment, the root fresh and dry weight, root diameter, root length and number of roots in the final of rooting procedure were recorded. Also at the end of the experiment, samples

were taken from the rooting media and the amount of available water was determined from the difference between water content at field capacity and permanent wilting point measured by pressure plates. Statistical analysis of the data for the hydrogel rates and irrigation interval was performed by analysis of variance (ANOVA), comparing the mean by the LSD method with using MSTAT-C and drawing diagrams by EXCEL.

## Results

### *Polymer application rates and irrigation interval effect on the amount of available water*

The results (Table 1) showed that the main effect of hydrogel application, irrigation interval and interaction of hydrogel application and irrigation interval had a significant effect on available water ( $p \leq 0.01$ ). The results of mean comparison (Fig. 1) showed that different irrigation interval had significant difference about this property, so that the maximum amount of available water related to 1 day irrigation interval and with increasing irrigation interval significantly decreased. Based on the comparison of different hydrogel levels impact on the amount of available water (Fig. 1), it was seen that the greatest amount of available water is related to 0.6% hydrogel treatment. Interaction of hydrogel application and irrigation interval on available water showed that the 0.3% hydrogel in 1 day irrigation interval has the highest mean. The minimum amount of available water was relevant to control treatment in 3 and 4 days irrigation interval (Fig. 1).

**Table 1.** Data analysis and statistics of the measurement indices of *Rosa Sp* cuttings under the experimental treatments.

SOV	Df	Mean of square						
		Amount of available water (%)	Percentage of rooted cuttings	Dry weight of cuttings root (g)	Fresh weight of cuttings root (g)	Root length (cm)	Root diameter (cm)	The number of roots per cuttings
Polymer rate	3	21.015**	2704.572**	3.139**	4.639**	8.173**	0.844**	48.188**
Irrigation interval	3	43.905**	6040.466**	14.346**	17.737*	66.198**	3.396**	93.688**
Polymer rate * Irrigation interval	9	4.250**	326.643**	0.292**	0.465**	1.678**	0.161*	6.688**
Error	32	0.099	1.847	0.035	0.051	0.035	0.029	0.783

\*, \*\* Significant at  $P=0.05$  and  $P=0.01$  levels, respectively.


### *Percentage of rooted cuttings*

Analysis of variance of irrigation interval and hydrogel

rates on percentage of rooted cuttings of *Rosa Sp* are included in table (1). As the results show, the main

effects of the use of hydrogels, irrigation interval and interactions effects of hydrogels and irrigation interval applications on percentage of rooted cuttings was significant ( $p \leq 0.01$ ). A mean comparison of hydrogel rates and irrigation interval application effects on percentage of rooted cuttings is shown in fig. (2). As it can be seen, the different irrigation interval had a significant effect on the average percentage of rooted cuttings and the highest percentage of rooted cuttings relate to 1 day irrigation interval. Hydrogel application in culture media have a significant effect on average of percentage of rooted cuttings, so that with increasing hydrogel rates from

zero to 0.3%, the percentage of rooted cuttings has increased. With increasing hydrogels from 0.3 to 0.6%, the percentage of rooted cuttings was significantly decreased (Fig. 2). The interaction between the hydrogel and of irrigation interval on the mean of percentage of rooted cuttings showed that the highest mean of rooted cuttings (83.3%) is relevant to 0.3% hydrogel with 1 day irrigation interval which with the same amount of hydrogel and 2 days irrigation interval not be significant. The lowest mean of rooted cuttings (15%) was related to control treatment (without hydrogel) in irrigation interval of 3 and 4 days (Fig. 2).



**Fig. 1.** Mean comparison of the different polymer rates and irrigation interval on the available water percentage (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).

#### *The number of roots per cuttings*

The analysis of variance and mean comparison results of number of roots per cuttings grown indifferent percentages of hydrogels are presented respectively in table (1) and Fig. 3. As shown in table (1), the effect of different percentages of hydrogel, irrigation interval, as well as their interaction on the number of roots per cuttings of *Rosa Sp* was statistically significant ( $p \leq 0.01$ ). The highest mean number of roots per cuttings related to 1 day irrigation interval and with increasing of irrigation interval this index was significantly decreased (Fig. 3). Application of hydrogel effect on the number of roots per cuttings and the highest number was related to application of 0.3% hydrogel and the lowest was related to control and 0.15% of the hydrogel. Interaction of hydrogel application and irrigation interval showed that the highest mean number of roots per cuttings was related to the use of 0.3% of the hydrogel in the 1 day irrigation interval


which had not significant difference with 0.3% hydrogel in 2 days irrigation interval. The lowest mean number of roots per cuttings was relevant to control with irrigation interval of 3 and 4 days and 0.15% of the hydrogel with 4 days irrigation interval (Fig. 3).

#### *Fresh weight of cuttings root*


The results of analysis of variance (Table 1) showed that the main effects of the application of Super AB A 200 hydrogel, irrigation interval and also interaction of them had significant effect on root fresh weight of cuttings of *Rosa Sp* ( $p \leq 0.01$ ). The mean comparison results (Fig. 4) showed that different irrigation interval had significant difference about fresh weight of root of cuttings, so that the highest root fresh weight was associated with 1 day irrigation interval and showed significant decrease with increasing of irrigation intervals. According to mean comparison of different

rates impact on root fresh weight (Fig. 3), it is understandable that the highest root fresh weight is related to 0.3% hydrogel treatment. Interaction between application of the hydrogel and irrigation interval of the mean weight of roots per cuttings showed that 0.3% hydrogel in 2 days irrigation

interval had the maximum mean that did not have significant difference in 0.3% hydrogel in 1 day irrigation interval. The lowest root fresh weight was in control in 3 and 4 days irrigation interval and also 0.15, 0.3 and 0.6% hydrogel rates for 4 days irrigation interval (Fig. 4).



**Fig. 2.** Mean comparison of the different polymer rates and irrigation interval on the percentage of rooted cuttings (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).



**Fig. 3.** Mean comparison of the different polymer rates and irrigation interval on the number of roots per cuttings (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).

#### *Dry weight of cuttings root*

The results showed that root dry weight was affected by irrigation interval (Table 1), and based on mean comparison the highest cuttings root dry weight was related to 1 day irrigation interval and root dry weight decreased significantly with increasing irrigation interval (Fig. 5). The hydrogel rates had a significant effect ( $p \leq 0.01$ ) on cuttings root dry weight (Table 1). With increasing hydrogels up to 0.3% the root dry weight was significantly increased and the highest root dry weight relevant to 0.3% hydrogel treatment (Fig. 5). The results of the mean comparison relating to the interaction of the hydrogel rates and irrigation interval indicated that the highest root dry weight was in 0.3% hydrogel with 1 day

irrigation interval which were not significantly different in same amount of hydrogel in 2 days irrigation interval. The lowest root dry weight was related to the control in 3 and 4 days interval irrigation that did not show significant difference with 0.15, 0.3 and 0.6% hydrogel rates for 4 days irrigation interval (Fig. 5).

#### *Root diameter growth*

The analysis of variance and mean comparison results of diameter of root growth in cuttings grown in media impregnated with different percentages of hydrogel rates respectively presented in table (1) and fig. 6. As shown in table (1), the effect of different percentages of hydrogel and irrigation interval as well as their interactions on

root diameter growth of *Rosa Sp* cuttings was significant ( $p \leq 0.01$ ). According to the mean comparison (Fig. 6), the mean diameter of roots per cuttings in 1 to 2 days irrigation interval were not significantly difference and this index decreased significantly with increase of the irrigation interval. The maximum diameter of the roots was related to 0.3% hydrogel treatment and minimum was related to control and

0.15% of the hydrogels rates. Interaction of irrigation interval and hydrogel rates had significant effect on root diameter and showed that the greatest root diameter were in 0.3% hydrogel in 2 days irrigation interval and so that was not significantly different with this and 0.6% hydrogel rates in 1 day irrigation interval (Fig. 6).




Fig. 4. Mean comparison of the different polymer rates and irrigation interval on the fresh weight of cuttings root (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).





Fig. 5. Mean comparison of the different polymer rates and irrigation interval on the dry weight of cuttings root (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).

#### Root length

The analysis of variance results (Table 1) showed that the main effects of hydrogel application, irrigation interval and also interaction of hydrogel application and irrigation interval had significant effect on root elongation of *Rosa Sp* cuttings ( $p \leq 0.01$ ). The mean comparison results (Fig. 7) showed the greatest elongation of roots were related to 1 and 2 days irrigation interval and this index significantly decreased with increasing of irrigation interval. Application of hydrogel effect on root elongation so that the maximum root length was related to 0.6% hydrogel treatment which was not significantly

different with 0.3% treatment but had significantly different with control and other values. The mean comparison of interaction between irrigation interval and hydrogel rates (Fig. 7) showed the greatest root length was related to 0.6% hydrogel with 1 day irrigation interval which there was no significant difference with the same amount of hydrogel in 2 days irrigation interval and 0.3% hydrogel in 1 to 2 days irrigation interval. The lowest mean was relevant to control in 3 and 4 days irrigation interval and also 0.15 and 0.6% hydrogel in irrigation interval of 4 days.





**Fig. 6.** Mean comparison of the different polymer rates and irrigation interval on the root diameter growth (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).


## Discussion

Rooting of *Rosa Sp* cuttings is an important step in the reproduction of the plant and is sensitive to the lack of moisture. Effect of moisture stress can cause failure or delay rooting that is different based on degree of moisture lack. Studies of hydrophilic hydrogel in rooting media of plants indicated that with quick absorption of water and keeping it, could decrease moisture stress and increase the water absorption efficiency. The amount of this increasing depending on the application of the hydrogel and irrigation interval (Huttermann *et al.*, 1999). The results of this study showed that, in the absence of hydrogel, with increasing irrigation interval from 1 day to 4 days all parameters measured included the percentage of rooted cuttings, fresh and dry root weight, root length, root diameter growth, number of roots per cuttings showed a significant reduction in each cuttings (Fig. 2 to 7). In fact, the reduction in these parameters with increasing irrigation interval is due to the lack of available water in the bed of cuttings. The results of present study is corresponding to results of Tavakoli *et al.*, 2011, Abedi and Asad Kazemi, 2006 and Taylor and Halfacre, 1986 that rooting index decreased with increasing of irrigation interval. With increasing hydrogel rates from zero up to 0.3%, rooting indices showed a significant increase and reached its maximum value in the 0.3% hydrogel concentration (Fig. 2 to 7). The reasons of increasing of rooting indices due to applying hydrogels up to 0.3% could be attributed to potentially high moisture absorption of hydrogel and storage of water in its network and due to increasing of available water 2 times versus control

(Fig. 1). With increasing of hydrogels up to 0.6%, the percentage of rooting and fresh and dry weight of cuttings decreased significantly. It appears that the reason for this decline is that with the large amounts of hydrogel, the water amount in the rooting media of cuttings too (2.75 times compared to no use of hydrogels, Fig. 1) increases. This increase is primarily a result of reduced air-filled pore space for cuttings media because of hydrogel swelling by water (Still, 1976). Second, excessive moisture can cause the slime of rooting media and resulting of decrease of physiological and metabolic processes of root. These results are compatible with the results of Assareh *et al.*, 2007, Meghani *et al.*, 2009, Farokhzad *et al.*, 2009 and Huttermann *et al.*, 1999. Based on the mean comparison results of interaction between irrigation interval and hydrogel rates at each irrigation interval (Fig. 2 to 7), it was found that with the increasing use of hydrogels up to 0.3%, the measured rooting's cuttings indices increased and the maximum value was related to 2 days irrigation interval. The reason of increasing of these indices in low irrigation interval (2 days compared with the control and 1 day) could be that the addition of hydrogel to media culture of cuttings could act like a storage water tank during the dry period and increased rapidly the moisture retention period of cuttings media and wetting and redistribution of water in the rooting zone of cuttings after irrigation. The results of this study is coincide with results of Jandagheyan *et al.*, 2001 about rooting of *Phylodendron scandens* and *Pelargonium hortorum*, Ghasemi and Khoshkhoe, 2007 of *Chrysanthemum*

*morifoliumramat*, Gehring and Lewis, 1980 of *Zinnia elegans* and *Callendula officinalis*, El-Hady and Wanas, 2006 of greenhouse cucumbers and Woodhouse and

Johnson, 1991 of *Cupressus arizonica* that the improved rooting index quantities derived from the interaction of the hydrogel and irrigation interval.



**Fig. 7.** Mean comparison of the different polymer rates and irrigation interval on the Root length (Bars within a polymer class having the same letter are not different at  $P = 0.01$ ).

### Conclusion

According to the present results can be stated that the use of hydrogels based on rates of 0.3 and 0.6, the amount of available water increased 2 to 2.75 times respectively. The independently hydrogels usage as much as 0.3% increased rooting of *Rosa Sp* cuttings and improved root morphological characteristics, while irrigation interval alone has had a negative effect on rooting of cuttings of this plant. Interaction between the hydrogel and irrigation interval in the presence of 0.15, 0.3 and 0.6% hydrogel and 2 days irrigation interval improved rooting and root cuttings of *Rosa Sp* and morphological characteristics compared to the control treatment. But the highest rooting percentage, root fresh and dry weight and length and diameter of roots were observed in the 0.3% hydrogel and 2 days irrigation interval. According to these results, applying of 0.3% polyacryl amide A 200 polymer with 2 days irrigation interval can be suggested for optimum rooting of cuttings of *Rosa Sp* plant.

### References

**Abedi-Koupai J, Asadkazemi J.** 2006. Effects of hydrophilic polymer on the field performance of an ornamental plant (*Cupressus arizonica*) under reduced irrigation regimes. *Iranian Polymer Journal* **15**, 715-725.

<http://dx.doi.org/org/10.1016/j.scienta.2009.12.031>

**Akhter J, Mahmood K, Malik Mardan KA, Ahmad AM, Iqbal MM.** 2004. Effects of hydrogel amendment on water storage of sandy loam and loam soils and seedling growth of barley, wheat and chickpea. *Plant soil environment* **50(10)**, 463- 469.

**Assareh MH, Nikvash N, Ghorbanli M, Ghamarizare A.** 2007. Investigation on anatomical structure of yew cuttings rooting and reasons of their hard and late rooting. *Journal Pajouhesh and Sazandegi* **74(2)**, 114-123.

**El-Hady OA, Wanas SHA.** 2006. Water and fertilizer use efficiency by cucumber grow under stress on sandy soil treated with acrylamide hydrogels. *Journal of Applied Sciences Research* **2(12)**, 1293-1297.

**Farjon A.** 1998. *World Checklist and Bibliography of Conifers*, Royal Botanic Gardens, Kew.

**Farokhzad A, Asghari M, Ghasemzadeh R, Asadzadeh A, Hojattei Y.** 2009. Effect of polyacrylamide polymer and indole butyric acid treatments on rooting of cuttings of *Magnolia soulougeana*. cv. Violet. *Journal Pajouhesh and Sazandegi* **83**, 2-15.

**Gehring JM, Lewis AJ.** 1980. Effect of hydrogel on wilting and moisture stress of bedding plants. *Journal of*



the American Society for Horticultural Science **150** (4), 511-513.

**Ghasemi M, Khoshkhoue M.** 2007. Effect of Superabsorbent polymers on irrigation interval and growth and development of *Chrysanthemum morifolium* ramat. Journal of Horticultural Science and Technology **8**(2), 65-82.

**Hüttermann A, Zomporodi M, Reise K.** 1999. Addition of hydrogels to soil for prolonging the survival of *Pinus helvensis* seedlings subjected to drought. Soil and Tillage Research **50**, 295-304.  
[http://dx.doi.org/10.1016/S0167-1987\(99\)00023-9](http://dx.doi.org/10.1016/S0167-1987(99)00023-9)

**Jalili K, Jalili J, Sohrabi H.** 2011. Effect of Super Absorbent Polymer (Tarawat A 200) and irrigation interval on growth of Almond Sapling. Journal of Soil and Water **21**(2), 121-134.

**Jandagheyan M.** 2001. Investigation of effect Polyacrilamid on *Phylodendron scandens* S and *Pelargonium hortorum* L rotting. MS Thesis, Islamic Azad University, Shiraz Branch, Iran, 81-92.

**Malkeshi SH, Mojib Hagh Ghadam Z.** 2009. The review of important rose pests and their integrated pest management in landscape. 3<sup>rd</sup> National Symposium of Green Area, Iran.

**Meghani H, Solimane A, Askarei N.** 2010. Effect of substrate on rooting of *Bougainvillea glabra*

cuttings. 6<sup>th</sup> Iranian Horticultural Science Congress. 345-350.

**Merkle SA, Nairn CJ.** 2005. Hardwood tree biotechnology. Journal of In Vitro Cellular and Developmental Biology-Plant **41**, 602-619.  
<http://dx.doi.org/10.1079/IVP2005687>

**Peterson D.** 2002. Hydraulic polymers—effects and uses in the landscape. Restoration and reclamation review. University of Minnesota. St. Paul, Minnesota. 13-16.

**Still SM.** 1976. Growth of 'Sunny Mandalay chrysanthemums' in hardwood-bark amended media as affected by insolubilized poly (ethylene oxide). HortScience **11**, 483-484.

**Tavakoli A, Purreza M, Khodakarami Y.** 2011. Preliminary investigation on possibility of rooting of Manna oak by layering in Zagros forests. Iranian Journal of Forest and Poplar Research **19**(3), 432-440.

**Taylor KC, Halfacre RG.** 1986. The effect of hydrophilic polymer on media water retention and nutrient availability to *Ligustum lucidum*. Horticultural Science **21**(5), 1159-1161.

**Woodhouse J, Johnson MS.** 1991. Effect of superabsorbent polymers on survival and growth of crop seeding. Agriculture Water Management **20**, 63-70.  
[http://dx.doi.org/10.1016/0378-3774\(91\)90035-H](http://dx.doi.org/10.1016/0378-3774(91)90035-H)